

Field Performance of Seedlings and Micropropagated Plantlets of Carob Tree

M.L. Osório*, J. Osório, M.M. David and A. Romano

Faculdade de Engenharia de Recursos Naturais, Universidade do Algarve, Campus de Gambelas, 8005-139 Faro, Portugal. Fax: + 351 289 818419

*Corresponding author: e-mail: mlosorio@ualg.pt

Keywords: Photosynthesis, photoinhibition, photochemistry, survival

Abstract

Micropropagated plantlets of *cv. Mulata* and *cv. Galhosa* growing in pots, after acclimatization in the glasshouse and growth for several months under natural conditions, were transferred to a field trial. Carob seedlings of 'Mulata' were also transferred under the same conditions. The field trial has been established with 100% of success in micropropagated plants and 97% in seedlings. Three months after transfer to the field, plants showed good growth but micropropagated plants of 'Mulata' exhibited more vigour than 'Galhosa' and than seedlings, with the greatest number of branches and a larger length increase in the main stem. Before transference to the field, net photosynthetic rates (P_N), water use efficiency (P_N/g_s) and quantum yield of PSII (ϕ_{PSII}) of the potted plants were higher in seedlings than in micropropagated plants. Three months after transplantation all gas exchange parameters were improved for all types of plants, particularly net photosynthetic rate, and no significant differences were observed between plants either micropropagated or seedlings. On the other hand, ϕ_{PSII} decreased significantly in seedlings. The results suggest that the pre-acclimation, in pots, to external environmental conditions might have improved field survival, plant vigour and enhanced the endurance to adverse conditions of micropropagated carob trees.

INTRODUCTION

Carob tree (*Ceratonia siliqua* L.) is a large (10-20 m) evergreen polygamotrioecious tree of slow growth and great longevity. Due to its interesting agroecological features, such as resistance to drought and salinity, adaptation to poor soils, and minimal cultural requirements, it has been introduced into the whole Mediterranean basin and other Mediterranean-like regions (Battle and Tous, 1997). Traditional carob propagation has been achieved by grafting saplings with female buds of chosen productive trees. This traditional method of propagation has failed to meet the market demand for new, selected plant material, required to establish the new early-producing orchards on a large scale. Thus, the use of micropropagation techniques seems to be appropriate to fulfill the increased demand for propagating this tree (Romano *et al.* 2002). The transplantation of micropropagated plants into the field is usually a critical step and substantial numbers of plants die after transfer. Although carob tree is well adapted to the Mediterranean climates, the performance of micropropagated plants can be affected when moved to the field, since they will be coping with substantial lower relative humidity, higher light and higher or lower temperature which are potential

stressful conditions to them. It has been demonstrated that the combination of these factors, that predispose plants to photoinhibition or down-regulation process, contributes to the reduction in carbon assimilation that will further affect the young plants ability for growth and survival (Chaves *et al*, 2002). In particular, a CO₂ deprivation at the chloroplast level by stomatal closure could enhance the sensitivity of the photosynthetic apparatus to high irradiance (Flexas *et al.* 1998). Protection mechanisms against excess light are thus an important strategy under Mediterranean conditions and may be achieved by several mechanisms such as the regulated thermal dissipation at the light harvesting complexes (Demmig-Adams and Adams 1996). This photoprotective mechanism competes with photochemistry for the absorbed energy, leading to a decrease in quantum yield of PSII (Genty *et al.* 1989).

As it is still a matter of uncertainty how environmental constraint factors affect the field establishment of micropropagated plants of carob tree, the present research was undertaken in order to investigate how micropropagated plants of two Portuguese carob cultivars, Galhosa and Mulata, deal with the new environment in terms of survival, growth and photosynthetic performance. Differences between micropropagated plants and seedlings were also evaluated.

MATERIAL AND METHODS

Micropropagated plants of *Ceratonia siliqua* (L.) from cv Mulata and cv Galhosa growing in pots, after acclimatization in the glasshouse and growth for several months under natural conditions, were transferred to a field trial in the south of Portugal, where survival rate and physiological performance has been followed. Carob seedlings of 'Mulata' were also transferred under the same conditions. Before transplantation superphosphate and natural organic fertilizers were applied to the soil. Plants were irrigated twice a week during summer. Gas exchange and chlorophyll *a* fluorescence measurements were taken one week before and three months after transplantation. Gas exchange measurements were carried out in the youngest fully expanded leaf at the middle of the light period, using a portable Minicuvette System HCM 1000 (Walz, Effeltrich, Germany). Leaf net photosynthetic rate (P_N) and stomatal conductance (g_s) were calculated according to the equations of von Caemmerer and Farquhar (1981). Chlorophyll *a* fluorescence measurements were performed in the same leaves using a portable pulse amplitude modulation fluorometer (PAM-2000 system, Walz, Germany). The maximal photochemical efficiency of PSII (F_v/F_m) was estimated from F_o (basal fluorescence) and F_m (maximal fluorescence) which values were taken before dawn. The intrinsic efficiency of open PSII reaction centers (F'_v/F'_m) was calculated from basal and maximal fluorescence measured under natural irradiance (F'_o and F'_m) and the quantum yield of PSII in light-adapted leaves (ϕ_{PSII}) was evaluated by the $(F'_m - F'_s)/F'_m$ ratio (Genty *et al.* 1989). The photochemical quenching (q_p), which was used as an estimate of the fraction of open centres, was calculated as: $q_p = 1 - (F'_s - F'_o)/(F'_m - F'_o)$ (Bilger and Schreiber 1986) and the thermal energy dissipation at the PSII as: $NPQ = (F_m/F'_m) - 1$ (Cornic 1994). The electron transport rate (ETR) was estimated as described by Krall and Edwards (1992), by multiplying $\Delta F/F'_m \times PPF \times 0.5 \times 0.84$. Plant survival and some shoot growth parameters were also evaluated.

RESULTS AND DISCUSSION

Survival and Growth

Micropropagated plants survived transplanting in excellent conditions. The field trial was established with 100% of success in micropropagated plants and 97% in seedlings. In general, plants showed good growth three months after transfer to the field, but micropropagated plants of 'Mulata' exhibited more vigour than 'Galhosa' and than 'Mulata' seedlings, exhibiting the greatest number of branches. The shoot length increase was also higher in micropropagated 'Mulata' plants, although no statistical significant difference has been observed between plants (Table 1).

Gas Exchange and Chlorophyll Fluorescence

Before transplantation no significant differences were observed in net photosynthetic rate (P_N), stomatal conductance (g_s), and instantaneous water use efficiency (P_N/g_s), between 'Galhosa' and 'Mulata'. P_N and P_N/g_s were higher in seedlings than in micropropagated plants, but these differences were only significant relatively to 'Mulata' (Fig. 1A, B, C). Despite the reduction in P_N of micropropagated plants relative to seedlings, the maximal PSII photochemical efficiency (F_v/F_m) was similar and high (0.78-0.73) in the three types of plants, which indicates no PSII damage. No significant differences were observed in the intrinsic efficiency of open PSII reaction centers (F'_v/F'_m), the photochemical quenching (q_p) and non-photochemical quenching (NPQ) of chlorophyll fluorescence between the three types of plants (Fig. 2B, C, D). However, the quantum yield of PSII electron transport in the light (ϕ_{PSII}) was higher in seedlings (Fig. 2A), which agrees with the best photosynthetic capacity of these plants, and can be related with a greater concentration of chlorophyll.

After transplantation all gas exchange parameters were improved and no significant differences were observed between plants (Fig. 1D, E, F, G), despite ϕ_{PSII} has decreased in seedlings (Fig. 2 D). On the other hand, the (F_v/F_m) ratio declined significantly (0.57-0.63), which may indicate that some degree of photoinhibition occurred. The observed decrease on NPQ (Fig. 2H), also point out that regulated thermal dissipation in light harvesting complexes were not promoted suggesting that this mechanism could not be enough to avoid some photoinhibition. Carob tree has been considered the most sensitive species, among other Mediterranean sclerophyll species, to low temperatures (Larcher, 1981). Considering the low night temperatures that occurred in the week of the field measurements (about 6 °C at predawn), the authors ascribe the photoinhibitory effect to this particular environmental condition and not to transplantation *per se*.

CONCLUSIONS

This study shows that micropropagated plants of carob tree have excellent field survival. Results suggest that the pre-acclimation to external environmental conditions in pots might have improved field survival and plant vigour. The enhanced photosynthetic carbon assimilation and growth three months after transplantation are presumably related with adequate fertilization (enough nutrient reserves), and with the absence of a volumetric limitation to the root system development.

ACKNOWLEDGEMENTS

We thank financial support by INIAP - project AGRO 770 "Plantation of a clonal field of micropropagated plants of carob tree".

Literature Cited

- Battle, I. and Tous, J. 1997. Carob tree. *Ceratonia siliqua* L. Promoting the conservation and use of under-utilised and neglected crops. Plant Genetic Resource Institute, Gatersleben/International, Rome.
- Bilger, W. and Schreiber, U., 1986. Energy-dependent quenching of dark level chlorophyll fluorescence in intact leaves. *Photosynth. Res.* 10: 303-308.
- Chaves, M.M., Pereira, J.S., Maroco, J., Rodrigues, M.L., Ricardo C.P.P., Osório, M.L., Carvalho, I., Faria, T. And Pinheiro, C., 2002. How plants cope with water stress in the field. *Photosynthesis and growth. Ann. Bot.* 89: 907-916.
- Cornic, G., 1994. Drought stress and high light effects on leaf photosynthesis. p. 297-313. In: Baker NR, Bowyer JR (Eds.), *Photoinhibition of photosynthesis: From Molecular Mechanisms to the Field.* BIOS Scientific Publishers, Oxford.
- Demmig-Adams, B. and Adams, W.W.III., 1996. Xanthophyll cycle and light stress in nature: uniform response to excess direct sunlight among higher plant species. *Planta* 198: 460-470.
- Flexas, J., Escalona, J.M. and Medrano, H., 1998. Down-regulation of photosynthesis by drought under field conditions in grapevines leaves. *Aust. J. Plant Physiol.* 25: 893-900.
- Genty, B., Briantais, J.M. and Baker, N.R., 1989. The relationship between the quantum yield of photosynthetic electron transport and quenching of chlorophyll fluorescence. *Biochim. Biophys. Acta* 990: 87-92.
- Krall, J.P. and Edwards, G.E., 1992. Relationship between photosystem II activity and CO₂ fixation in leaves. *Physiol. Plant.* 86: 180-187.
- Larcher, W. 1981. Low temperature effects on Mediterranean sclerophylls: an unconventional viewpoint. In: Margaris & Mooney (Eds.), *Componentes of Productivity of Mediterranean-Climate Regions, Basic and Applied Aspects.* Martinus Nijhoff/Dr W. Junk Publishers, The Hague.
- Romano A., Barros S. and Martins-Loução M.A. 2002. Micropropagation of the mediterranean tree *Ceratonia siliqua* L. *Plant Cell, Tissue and Organ Culture*, 68: 35-41.
- Von Caemmerer, S. and Farquhar, G.D., 1981. Some relationships between the biochemistry of photosynthesis and the gas exchange of leaves. *Planta* 153: 376-387.

Tables

Table 1. Main shoot length increase and number of new leaves and lateral branches of micropropagated ‘Mulata and ‘Galhosa’ and seedlings of carob tree three months after transplanted at field.

Plant Type	Survival (%)	Increase in shoot length (cm)	N° new leaves on terminal shoot	N° of new lateral branches
‘Mulata’	100 a	6.4 ± 1.2 a	4.2 ± 0.5 ab	1.8 ± 0.3 a
‘Galhosa’	100 a	3.7 ± 1.1 a	5.5 ± 0.4 a	0.1 ± 0.2 ab
Seedling	97 a	3.9 ± 1.5 a	3.3 ± 0.5 b	0.0 ± 0.0 b

The values shown are means ± SE from five samples. Values followed by the same letter are not significantly different at $P \leq 0.05$ (one-way ANOVA, Bonferroni test).

Figures




Figure 1. Net photosynthesis rate (P_N), stomatal conductance (g_s) and instantaneous water use efficiency P_N/g_s of micropropagated plants and seedlings of carob tree before (Pots) and after (Field) transplanted. The values shown are means ± SE from five samples. Values followed by the same letter are not significantly different at $P \leq 0.05$ (one-way ANOVA, Bonferroni test).




Figure 2. Quantum yield of PSII (ϕ_{PSII}), intrinsic efficiency of open PSII reaction centers (F_v/F_m), photochemical quenching (q_p) and non-photochemical quenching (NPQ) of chlorophyll fluorescence of micropropagated plants and seedlings of carob tree before (Pots) and after (Field) transplantation. The values shown are means \pm SE from five samples. Values followed by the same letter are not significantly different at $P \leq 0.05$ (one-way ANOVA, Bonferroni test).