



Hypoxic Induced Decrease in Oxygen Consumption in Cuttlefish (*Sepia officinalis*) Is Associated with Minor Increases in Mantle Octopine but No Changes in Markers of Protein Turnover

Juan C. Capaz^{1†}, Louise Tunnah^{2†}, Tyson J. MacCormack², Simon G. Lamarre³, Antonio V. Sykes^{1*} and William R. Driedzic^{4*}

OPEN ACCESS

Edited by:

Graziano Fiorito,
Stazione Zoologica Anton Dohrn, Italy

Reviewed by:

Claudio Agnisola,
University of Naples Federico II, Italy
Panagiotis Grigoriou,
Hellenic Centre for Marine Research,
Greece

*Correspondence:

Antonio V. Sykes
asykes@ualg.pt
William R. Driedzic
wdriedzic@mun.ca

[†]These authors have contributed
equally to this work.

Specialty section:

This article was submitted to
Invertebrate Physiology,
a section of the journal
Frontiers in Physiology

Received: 05 January 2017

Accepted: 11 May 2017

Published: 26 May 2017

Citation:

Capaz JC, Tunnah L,
MacCormack TJ, Lamarre SG,
Sykes AV and Driedzic WR (2017)
Hypoxic Induced Decrease in Oxygen
Consumption in Cuttlefish (*Sepia
officinalis*) Is Associated with Minor
Increases in Mantle Octopine but No
Changes in Markers of Protein
Turnover. *Front. Physiol.* 8:344.
doi: 10.3389/fphys.2017.00344

¹ Centro de Ciências do Mar do Algarve, Universidade do Algarve, Faro, Portugal, ² Department of Chemistry and Biochemistry, Mount Allison University, Sackville, NB, Canada, ³ Département de Biologie, Université de Moncton, Moncton, NB, Canada, ⁴ Department of Ocean Sciences, Memorial University of Newfoundland, St. John's, NL, Canada

The common cuttlefish (*Sepia officinalis*), a dominant species in the north-east Atlantic ocean and Mediterranean Sea, is potentially subject to hypoxic conditions due to eutrophication of coastal waters and intensive aquaculture. Here we initiate studies on the biochemical response to an anticipated level of hypoxia. Cuttlefish challenged for 1 h at an oxygen level of 50% dissolved oxygen saturation showed a decrease in oxygen consumption of 37% associated with an 85% increase in ventilation rate. Octopine levels were increased to a small but significant level in mantle, whereas there was no change in gill or heart. There were no changes in mantle free glucose or glycogen levels. Similarly, the hypoxic period did not result in changes in HSP70 or polyubiquitinated protein levels in mantle, gill, or heart. As such, it appears that although there was a decrease in metabolic rate there was only a minor increase in anaerobic metabolism as evidenced by octopine accumulation and no biochemical changes that are hallmarks of alterations in protein trafficking. Experiments with isolated preparations of mantle, gill, and heart revealed that pharmacological inhibition of protein synthesis could decrease oxygen consumption by 32 to 42% or Na⁺/K⁺ ATPase activity by 24 to 54% dependent upon tissue type. We propose that the decrease in whole animal oxygen consumption was potentially the result of controlled decreases in the energy demanding processes of both protein synthesis and Na⁺/K⁺ ATPase activity.

Keywords: European cuttlefish, *Sepia officinalis*, HSP70, octopine, polyubiquitinated protein, ventilation frequency

INTRODUCTION

The European cuttlefish (*Sepia officinalis*) is a dominant species in the north-east Atlantic Ocean and Mediterranean Sea. It is a benthic species that occurs to depths of 200 m. Sexually mature and juvenile individuals migrate to coastal grounds in the spring; the latter in pursuit of abundant food sources (Pierce et al., 2008; Bloor et al., 2013). These geographic areas are at risk of becoming

hypoxic because of the increase of nutrients and organic matter into the coastal waters (Van Den Thillart et al., 1994; Grantham et al., 2004; Chan et al., 2008). Hypoxic conditions occur naturally in nature (Ekau et al., 2010); however, there are reports of a global oceanic oxygen decline (Schmidtko et al., 2017), which is becoming a problem for marine biodiversity (Vaquer-Sunyer and Duarte, 2008) and an important environment stressor to marine species (Cosme and Hauschild, 2016; Townhill et al., 2017). Hypoxia is also a potential challenge under intensive aquaculture conditions (Brett, 1979; Delaney and Klesius, 2004; Burt et al., 2013) due to the high density of animals and because of the costs associated with keeping oxygen saturation at normal levels. Routine oxygen consumptions of sepoids are within the same range of those of octopus species (Boyle, 1991) and optimal oxygen saturations for *Octopus vulgaris* are reported to be in between 100 and 65% dissolved oxygen saturation (DO₂) for proper food intake and growth, while suboptimal (modest hypoxia) values range from 65 to 35% DO₂ (Cerezo Valverde and García García, 2005). These suboptimal values may be reached with a combination of high densities and temperatures in Mediterranean countries, where cuttlefish culture will take place. This is relevant because oxygen availability limits thermal tolerance in *S. officinalis* (Melzner et al., 2006), O₂ consumption rates rise with increasing temperature up to a metabolic threshold (Melzner et al., 2007) related to the oxygen-binding properties of haemocyanin, which lowers with increasing temperature (Brix et al., 1994). As such, it is important to have an understanding of the response of cuttlefish to challenges of a suboptimal oxygen environment that is considered to be either ecologically and/or practically relevant.

S. officinalis, as with other cephalopods, exhibit a decrease in oxygen consumption below a critical environmental oxygen level (Houlihan et al., 1982; Johansen et al., 1982; De Wachter et al., 1988; Cerezo Valverde and García García, 2005; Seibel et al., 2014). Severe hypoxia induces a mobilization of mantle glycogen pools in this species with concomitant accumulation of the anaerobic end product octopine (Storey and Storey, 1979; Storey et al., 1979). The production of octopine is catalyzed by the enzyme octopine dehydrogenase (ODH) that occurs in particularly high levels in mantle, ventricle, and tentacle relative to other tissues (Storey, 1977). Despite this existing information, the biochemical responses of this species, at the tissue level, to environmentally relevant levels of hypoxia have not been fully characterized, so the importance of anaerobic octopine production under these conditions is unclear. In addition to octopine accumulation, more recent studies with the exceptionally hypoxia tolerant Humboldt (jumbo) squid, *Dosidicus gigas*, reveal other aspects of the response to low oxygen levels that include a decrease in mantle contraction frequency (i.e., gill ventilation) and increases in mantle heat shock protein 70 (HSP70) and ubiquitinated proteins (Trübenbach et al., 2013a,b, 2014). Here we question if these responses occur in *S. officinalis* as well, as it is considered a hypoxia-sensitive species (Storey et al., 1979).

Decreases in oxygen consumption that are not due to reduced spontaneous swimming activity must be related to decreases in other components of energy demand. Aside from muscle

contraction, the two most energy demanding processes in cells are the maintenance of ionic gradients via Na⁺/K⁺ ATPase and protein turnover (synthesis and degradation) (Wieser and Krumschnabel, 2001). The rapid growth rates of cephalopods have been suggested to occur due to high rates of protein synthesis and high efficiency of protein retention, which implies low protein degradation (Houlihan et al., 1990; Carter et al., 2009; Moltschanivskyj and Carter, 2010). It is possible to measure the relationship between these processes and oxygen consumption by pharmacological inhibition of Na⁺/K⁺ ATPase with ouabain and protein synthesis with cycloheximide (Wieser and Krumschnabel, 2001; Agin et al., 2003). Here we use these pharmacological tools on isolated preparations to assess the potential tissue specific contribution of Na⁺/K⁺ ATPase activity and protein turnover to the whole animal decrease in oxygen consumption.

Mantle is the largest tissue of *S. officinalis*, corresponding to approximately 30–40% of body mass (Castro et al., 1992; Speers-Roesch et al., 2016). We focused our attention on this tissue for assessment of biochemical changes and to determine its potential contribution to alterations in metabolic demand under hypoxia. Although gill and heart represent only about 2 and 0.1% of body mass, respectively (Speers-Roesch et al., 2016), these tissues were also included in our study as they are essential organs to supply oxygen to the rest of the animal, which must maintain function under hypoxia, and have been previously connected to high fractional rates of protein synthesis (Houlihan et al., 1990).

MATERIALS AND METHODS

Ethical Statement

All the procedures were approved by CCMAR Animal Welfare Committee (ORBEA CCMAR-CBMR) and Direção-Geral de Alimentação e Veterinária (DGAV) of the Portuguese Government, according to National (Decreto-Lei 113/2013) and EU legislation (Directive 2010/63/EU) on the protection of animals used for scientific purposes. In addition, protocols were approved by institutional Animal Care Committees at each of the Canadian Universities where authors of this paper are based. Procedures were only applied to live animals by authorized users.

Animals

Cuttlefish (*S. officinalis*) were reared according to the latest culture technology described in Sykes et al. (2014). All experiments used juveniles produced from eggs laid by a F6 captive stock. The mass of group 1 was 59.3 ± 5.4 g (*N* = 18) and of group 2, used only in mantle oxygen consumption experiments to avoid limiting oxygen diffusion across thick mantle tissue, was 1.90 ± 0.14 g (*N* = 7). Experiments were done during May 2016, at CCMAR's Ramalhete Aquaculture Station (Ria Formosa, Portugal—37°00′22.39″N; 7°58′02.69″W). Temperature, salinity, and DO₂ were measured daily, at 9 h 30, in the stock tank. Both temperature and DO₂ were measured with a VWR DO220 probe, while salinity was measured with a VWR EC300 salinity meter. Water temperature was 20.5 ± 1.27 (S.D.)°C, salinity was 35.9 ± 0.9 (S.D.) g L⁻¹ and DO₂ level

was 101.1 ± 1.2 (S.D.)%. Cuttlefish were fed frozen grass shrimp (*Palaemonetes varians*) *ad libitum* on a daily basis.

Oxygen Uptake and Ventilation Frequency in Whole Animals

Two closed respirometry (Lefevre et al., 2016) acrylic chambers ($48.5 \times 28.5 \times 29$ cm), each with an external recirculating pump (Maxi-Jet PH MP400) and a PASCO PasPort PPS-2169 multiparameter sensor were used. The recirculating system had a total volume of 41 L. Seawater entered and exited on opposite sides of the chamber at a flow of 200 L h^{-1} . Dissolved oxygen saturation data were collected in the upper right section of the chamber at 1 Hz in the PASCO Capstone software and at a temperature of 19.9 ± 1.36 (S.D.) °C. The chamber was completely isolated from any visual stimuli (both researchers and light) by the application of matte black stickers on the outer walls, except from above and one of the 48.5 cm sides. This side had a rectangular LED stripe ($50 \times 30 \times 50 \times 30$ cm) attached that emitted red light (wavelength peak of 630 nm) until the chamber was completely filled with water and, thereafter, white light (wavelength peaks at 460, 520 and 630 nm) with an intensity of 80 lux. These conditions allowed for the recording of movement (from above) with a Logitech C615 HD camera, and of behavior and ventilation rates (from the side) with a GoPro Hero 3⁺ Black Edition. Video recordings were made for the duration of each experiment at $1,920 \times 1,080$ pixel resolution and 24 frames per s (fps), from above, and $1,920 \times 720$ pixel and 50 fps, from the side.

Initially, 6 animals were individually exposed to a DO_2 of 100% and 6 other animals to a DO_2 of 50% oxygen saturation for 60 min. Each condition was tested pairwise, 1 cuttlefish at 100% DO_2 vs. another at 50% DO_2 . At the end of this procedure all the animals were euthanized. Samples of mantle, gill, and heart were rapidly removed and frozen in liquid nitrogen for biochemical analysis. Rates of oxygen consumption $\dot{M}\text{O}_2$ were determined for animals exposed to 50% DO_2 during 60 min. In addition, 6 further control animals (DO_2 of 100%) were used to collect $\dot{M}\text{O}_2$ during 90 min, to increase the number of available data points for regression fitting. Background $\dot{M}\text{O}_2$ (resulting from consumption of bacteria and electrode) was also assessed with no animal ($N = 3$) in the chamber for 90 min and was found to be negligible.

Hypoxia at 50% DO_2 was achieved by gassing seawater with pure N_2 using a ceramic stone in a closed 90 L reservoir. This water was then injected into the chamber, until the 41 L respirometry recirculating system was filled, and where the animal was already resting. Animals were removed from the stock tank with a black net and transported in a 10 L black bucket filled with seawater to the weighing station. They were individually weighed in a black plastic tray and quickly transported to the chamber area of the experimental room which was illuminated with 1 lux red light conditions (Philips TL-D Colored 18W Red 1SL/25). An outer room containing the data collection stations was isolated from the holding chambers. To avoid inking in the experimental chambers, animals were left resting in the bucket for 5 min before being lightly sedated (mild anesthesia) by adding

ethanol to a concentration of 2.5% (Fiorito et al., 2015). After sedation (≈ 2 min), animals were transferred to the chamber, which had 2 cm in height of seawater and thereafter filled with seawater from the 90 L reservoir (either 100 or 50% DO_2). As soon as the chamber was filled (≈ 2 min) and air purged, the system was put in recirculating mode, light switched to white and data acquisition was started on all the probes. This filling time was enough for recovery from sedation at this concentration (Gonçalves et al., 2012).

Ventilation rates were determined from video recordings, according to the following sampling scheme for each animal: (a) first and last 10 min—10 samples of 1 min each; b) from 10 to 50 min in hypoxia and normoxia (control)—1 sample every 10 min. After the procedure, the chamber was emptied, the animal removed, and euthanized with 10% ethanol in seawater (Sykes et al., 2012), the brain was then bisected downwards and forwards, followed by two lateral cuts to sever the brain from the optical lobes (Lewbart and Mosley, 2012). No relevant movements in the chamber nor extreme behavior reactions (Gonçalves et al., 2012) were recorded in any video sampling.

Biochemical Assays

Octopine, Glucose, and Glycogen

Tissues were homogenized and then sonicated (QSonica q55) in 6% HClO_4 (mantle 1 g tissue: 9 mL HClO_4 ; gill 1 g tissue: 4 mL HClO_4), and subsequently centrifuged at $14,000 \times g$ for 10 min at 4°C. The supernatant was neutralized with 2 M KHCO_3 . Octopine was assayed in buffer containing 100 mM TRIS, 8 mM NAD^+ , and excess octopine dehydrogenase (ODH) at pH 9.3. The increase in absorbance at 340 nm, following addition of enzyme, was monitored in a microplate reader. Octopine was calculated based on the NADH extinction coefficient of 6.22 and the pathlength of the plate reader assay was experimentally determined in near IR using the absorbance of water at 975 nm according to the following equation (Lampinen et al., 2012).

$$\text{pathlength} = \frac{OD_{975}^{\text{assay buffer}} - OD_{900}^{\text{assay buffer}}}{0.173} \times 10 \text{ mm}$$

ODH was isolated from frozen scallop, *Placopecten magellanicus*, adductor muscle by a modification of the method of Gäde (1985). Tissue was homogenized (25% w/v) in 100 mM TRIS (pH 7.5, 0.1 mM EDTA, 0.1 mM DTT) and centrifuged at $17,000 \times g$ for 10 min at 4°C. Ammonium sulfate was then added to the supernatant up to 65% saturation and stored at 4°C for 7 days to precipitate ODH. The sample was centrifuged, the pellet was re-suspended in 100 mM TRIS (pH 7.5, 0.1 mM EDTA, 0.1 mM DTT), the sample was desalted on a Sephadex G-50 column, and fractions containing ODH activity were retained. Contaminating dehydrogenases were removed by passing the sample through a Cibachrome Blue F3GA agarose column (Affi-Gel Blue, Bio-Rad Laboratories, Hercules, California, USA). Fractions containing ODH activity were combined and subsequently desalted on a Sephadex G-25 column to ensure removal of all residual metabolites. The resulting sample yielded 2 bands on an SDS-PAGE gel, one at ~ 43 kDa matching the molecular weight of ODH, and a second band of unknown identity at approximately

90 kDa. Lactate dehydrogenase activity was undetectable in the purified sample.

Glucose was analyzed in buffer containing 250 mM imidazole, 5 mM MgSO₄, 10 mM ATP, 0.8 mM NADP⁺, excess glucose 6-phosphate dehydrogenase and thereafter treated with excess hexokinase. A standard curve was created for glucose analysis. Assays were conducted in a microplate reader at a wavelength of 340 nm. Glycogen was assessed as glucosyl units following amyloglucosidase treatment of the neutralized supernatant (Keppler and Decker, 1974). Free glucose was subtracted from the total glucosyl equivalents to yield glycogen levels. All organic reagents were purchased from Sigma-Aldrich.

Polyubiquitin and HSP70

Polyubiquitinated proteins and HSP70 were determined by western blots. Frozen tissues were sonicated in 9 vol of 50 mM TRIS buffer (pH 7.4) and then centrifuged for 5 min at 14,000 × *g* at 4°C. The protein concentration in the supernatant was determined using the Bradford assay (Bio-Rad) and then adjusted to 3 μg/μL with the homogenization buffer. Tissue extracts were added to Laemmli sample buffer (1:1), boiled for 5 min and 4.5 μg of proteins were resolved by SDS-PAGE (TGX Stain-Free FastCast 10%, Bio-Rad). Resolved proteins were subsequently transferred to PVDF membranes, the membranes were blocked using 5% BSA before being incubated with the primary antibody. To analyze the levels of polyubiquitinated proteins, we used an antibody that only recognized proteins containing K48-linked polyubiquitin chains (Abcam ab190061) but had no cross-reactivity to monoubiquitin or polyubiquitin of other linkages. Levels of HSP70 were quantified by loading known quantities of a recombinant HSP70 (Enzo Life Sciences, #ADI-SPP-758, standard levels of 15, 30, and 60 ng per lane) along with the samples and using an HSP70 antibody (#AS05 083A, Agrisera). In all cases, secondary detection was done using an anti-rabbit HRP-linked antibody (for polyubiquitin: #7074, Cell Signaling Technology; for HSP70: SAB-300, Enzo Life Sciences). The chemiluminescence signal was detected using a CCD camera system (ImageQuant LAS 500, GE Life Sciences). Band intensity was quantified using ImageJ (imagej.nih.gov).

Oxygen Uptake by Isolated Preparations

Tissue $\dot{M}O_2$ was measured in 0.20 μm filtered seawater containing an additional 10 mM KCl, 200 mM taurine, and 1 mM glucose. The concentration of taurine used here is representative of measured tissue taurine levels in this species (Maccormack et al., 2016). Cuttlefish were euthanized as described above and 5–10 mg samples of gill, systemic heart, or mantle muscle were collected, weighed, and cut into 2–3 mg pieces using a razor blade. Tissue preparations from the same animal were then transferred to paired respirometry chambers (OX1LP, Qubit Systems Inc., Kingston, ON, Canada) containing 750 μL incubation medium maintained at 20 ± 0.1°C with a recirculating refrigerated water bath. Chambers were calibrated daily and background O₂ consumption rates were negligible without tissue. Baseline $\dot{M}O_2$ was recorded for 300 s before addition of pharmacological agents and recordings were continued for an additional 300 s before experiments were

terminated. Treatments consisted of 5 mM ouabain (e.g., Postel et al. (2000)) or 25 μM cycloheximide, a known antibiotic inhibitor of protein synthesis (Giuditta et al., 1968; Prozzo and Giuditta, 1973; Wieser and Krumschnabel, 2001) that has previously been applied in studies associated with learning in the species by blocking protein synthesis (Agin et al., 2003). Stock solutions were prepared in DMSO and delivered to the chambers in 10 μL. A tissue preparation from the same animal was run simultaneously with each drug treatment and exposed to 10 μL DMSO as a vehicle control. Chambers were thoroughly rinsed with 95% ethanol, ddH₂O, and incubation medium between runs and treatments were alternated between respirometry chambers to control for potential chamber effects. Respirometry chambers were interfaced to a LabQuest data acquisition system and DO₂ readings were collected over 10 min using LoggerPro V3.8 software (Vernier Software and Technology, Beaverton, OR, USA). Initial tissue $\dot{M}O_2$ prior to treatment was determined from the linear decrease in DO₂ between 120 and 300 s and post-treatment $\dot{M}O_2$ was determined between 400 and 600 s. $\dot{M}O_2$ was linear with tissue mass (data not shown) and consistent with previous experiments (Maccormack et al., 2016).

Data Analysis and Statistics


All values are expressed as mean ± s.e.m with the exception of water temperature, salinity, and DO₂ that are expressed as mean ± standard deviation. Statistical tests applied are stated in either the Results section or the legends to the figures. All data were tested for normal distribution with the Shapiro-Wilk test as well as for homogeneity of the variances with the Levene's test (Zar, 1999). Statistical difference was considered for *P* < 0.05.

RESULTS

Whole Animal Experiment

Figures 1A,B show examples of dissolved oxygen variation over the sampling period for individual cuttlefish either exposed to 100 or 50% DO₂ over 90 or 60 min, respectively. Oxygen consumption by hypoxic animals was 37% lower than animals held at 100% air saturation (189 ± 23 vs. 119 ± 6 nmol/g min) (Figure 1C). Cuttlefish maintained under control normoxic conditions exhibited a decrease in ventilation rate over the 60 min holding period ($y = -0.52x + 88$) (Figure 1D). Animals held under hypoxia also displayed a decrease in ventilation rate but to a lesser extent ($y = -0.1x + 102$). After 60 min ventilation rate was 85% higher in hypoxic than normoxic individuals.

Octopine was significantly higher in mantle than in gill or heart regardless of oxygenation condition. Octopine levels were significantly higher in mantle of hypoxic than normoxic animals but there was no significant difference between levels in normoxic or hypoxic animals in either gill or heart. (Figure 2A). There was no significant difference in either free glucose or glycogen levels in mantle between normoxic or hypoxic held animals (Figure 2B). HSP 70 was significantly higher in gill than in mantle or heart regardless of oxygenation condition (Figure 2C). There was no significant difference between levels in normoxic or hypoxic animals in any of the three tissues. Similarly, there



was no significant difference in levels of polyubiquitinated protein in mantle, gill or heart between treatment groups (Figure 2D).



Isolated Tissue Experiment

Pre-treatment oxygen consumption rates for gill, heart, and mantle are presented in Figure 3A. There was no significant difference in rates amongst the three tissues with values for all being approximately 500 nmol O₂/gm wet weight. min. Rates for gill and heart are from animals weighing approximately 59 g; while those for mantle are from smaller/younger animals with a mass of approximately 2 g. Preparations were treated with cycloheximide and ouabain to assess how much of the decrease in oxygen consumption noted in the whole animal study could potentially be attributed to protein synthesis and Na⁺K⁺ ATPase, respectively. Figure 3B shows the percentage decrease in oxygen consumption for each of the tissues following correction for any DMSO effect. Cycloheximide had a substantial effect upon oxygen consumption with average decreases ranging from 32% for gill to 42% for mantle. Similarly, ouabain treatment resulted in average decreases in oxygen consumption from 24% for mantle to 54% for heart.

DISCUSSION

Changes in $\dot{M}O_2$ and Ventilation Frequency under Hypoxia

Cuttlefish remained sedentary in the respirometry chambers under both normoxic and hypoxic conditions. As such, any difference in oxygen consumption between groups cannot be attributed to spontaneous swimming activity. When corrected for differences in temperature and body mass, the $\dot{M}O_2$ values reported here under control conditions are in the same range as rates previously noted (Johansen et al., 1982; Melzner et al., 2006, 2007; Grigoriou and Richardson, 2009; Lamarre et al., 2016). Here, exposure to 50% DO₂ for 60 min resulted in a 37% decrease in oxygen consumption. Although limited to 2 individuals of *S. officinalis* with a mean mass of 153 g, a decrease in $\dot{M}O_2$ of about 29% at a similar level of hypoxia was previously observed by De Wachter et al. (1988). Under comparable conditions, *S. officinalis* of body mass between 100 and 1,500 g showed a decrease in oxygen consumption of approximately 50% (Johansen et al., 1982). It is clear that a substantial reduction in $\dot{M}O_2$ is a common response to modest hypoxia



in *S. officinalis*; whether life could be sustained indefinitely at this level of hypoxia and if this will impact on the welfare (pain, suffering, distress, and lasting harm) of the animal remains unknown.

Ventilation rate under normoxic conditions decreased over the 60 min holding period in the same fashion as previously shown (Boal and Ni, 1996) and is presumably related to the stress of transferring the animals to the experimental chamber.

After 60 min the ventilation rate recorded here was in the same range as previously noted in other studies (Bone et al., 1994; Boal and Ni, 1996; Boal and Golden, 1999; Melzner et al., 2006). Ventilation rate in hypoxic animals remained relatively stable with time and was 85% higher in hypoxic than normoxic cuttlefish after 60 min. To our knowledge this is the first report of rate of ventilation in hypoxic *S. officinalis* or any other cuttlefish species. The response differs from inactive, juvenile Humbolt squid where deep hypoxia resulted in a decrease in ventilation frequency, as well as contraction strength and ventilation volume per min (Trübenbach et al., 2013a). The mantle muscle of squid and cuttlefish consists of a main mass of circular fibers that are mitochondria-poor and primarily anaerobic, as well as inner and outer layers of mitochondria-rich aerobic fibers (Bone et al., 1981; Mommsen et al., 1981). The central fibers are used in escape jetting contractions while the fibers of the inner and outer layers are responsible for rhythmical respiratory ventilation (Bone et al., 1981). It is probable that the increase in ventilation frequency observed here places additional demand to supply ATP to the contractile mechanisms under hypoxia even in the face of reduced oxygen consumption.

Anaerobic Metabolism is Minimal

Octopine dehydrogenase is the dominant opine dehydrogenase in *S. officinalis*. We were unable to detect activity of alanopine, taupine, or strombine dehydrogenase in this species' tissues (unpublished data). The level of ODH in mantle, heart and gill was 35, 20, and 2 $\mu\text{mol}/\text{min g}$ wet weight, respectively (Speers-Roesch et al., 2016). The level of octopine in mantle was substantially higher than in heart or gill, consistent with the high activity of ODH in mantle and lower activity in the other two tissues.

Sixty min of 50% hypoxia was associated with an increase in mantle octopine of 2.5 $\mu\text{mol}/\text{g}$. There was no change in octopine level in either gill or heart following the hypoxic period. The observed increase in octopine in mantle was low relative to the capacity to produce this metabolite. For instance, the increase in mantle octopine in *S. officinalis* forced to exercise and then subjected to hypoxia was 13 $\mu\text{mol}/\text{g}$ (Storey and Storey, 1979). Another study found a drastic elevation of octopine in the haemolymph of *S. officinalis* exposed to <30% O_2 hypoxia for about 60 min (Storey et al., 1979). A similar increase in mantle octopine occurred in hypoxic Humbolt squid (Seibel et al., 2014) and squid (*Lolliguncula brevis*) forced to swim to exhaustion (Finke et al., 1996). The modest increase in octopine observed in this study is in line with the lack of change in glycogen levels during the hypoxic challenge. Based on ATP equivalents that may be generated by aerobic and anaerobic metabolism, the contribution of octopine production to overall energy production is minimal.

The relative quantity of polyubiquitinated protein was similar in mantle, gill, and heart under control conditions. There was no impact of the hypoxic challenge in any of the tissues tested. This finding is in contrast to that for juvenile Humbolt squid where severe hypoxia resulted in a three-fold increase in ubiquitinated proteins with a size of approximately 100 kDa (Trübenbach

et al., 2014). These authors proposed that the increase in polyubiquitinated protein in association with a decrease in levels of at least two abundant proteins in the same size range (HSP90 and alpha-actinin) provides protection under hypoxia via a number of mechanisms, including the provision of certain amino acids for anaerobic energy production (Trübenbach et al., 2014). In *S. officinalis* food deprivation results in almost total depletion of digestive gland triglyceride, $\dot{M}\text{O}_2$ decreases by approximately 30%, and mantle polyubiquitin mRNA and polyubiquitinated proteins increase, suggesting an increase in protein degradation via the ubiquitin-proteasome pathway to provide amino acids for energetic purposes (Lamarre et al., 2012, 2016). Although in this species there appear to be mechanisms to increase protein breakdown in association with decreased $\dot{M}\text{O}_2$, we found no evidence that this occurs under modest hypoxia, at least based on levels of polyubiquitinated protein.

HSP70 is a well-recognized stress protein that plays a role in maintaining protein integrity via chaperoning newly synthesized and unfolding proteins (Balchin et al., 2016). HSP70 levels were significantly higher in gill than in mantle or heart and did not change in any tissue with exposure to hypoxia. Relatively high HSP70 levels in *S. officinalis* gill may be related to the high rate of protein synthesis in gill relative to mantle based on direct measurements (Lamarre et al., 2016), and relative to heart and mantle based on total RNA levels (Lamarre et al., 2012). Furthermore, since gill is at the interface with the aquatic environment it may experience more perturbations than other tissues and so has higher constitutive levels of HSPs as protection. HSP70 levels increased in mantle of hypoxic juvenile Humbolt squid (Trübenbach et al., 2013b) but not in adult animals (Seibel et al., 2014). Thus, the generality of a HSP70 response to hypoxia in cephalopods is unresolved but it is clear that under the conditions of the current study there was no change in HSP70 level.

The minimal increase in octopine content, along with stability in glycogen pools, polyubiquitinated protein level, and HSP70 levels in cuttlefish subject to 50% hypoxia for 1 h suggests that anaerobic metabolism is activated to only a limited extent and that the challenge is not particularly stressful to the animal. The 37% decrease in $\dot{M}\text{O}_2$ must be associated with an equivalent decrease in energy demand. Given that the animals are quiescent under both control and hypoxic conditions, the question becomes as to what metabolic processes are decreased to maintain energy balance.

Potential Sites of Energy Reduction – Insights from Isolated Tissue Experiments

There was no difference in the rate of oxygen consumption between tissue slices of gill, heart, and mantle. The *in vitro* activity of citrate synthase, a qualitative indicator of aerobic metabolism, was approximately 20-fold higher in *S. officinalis* heart than in mantle or gill which were similar to one another (Speers-Roesch et al., 2016). The relatively low rate of oxygen consumption in heart noted here is likely due to the non-contractile nature of

the preparation and reflects the basal metabolism of quiescent cells. The heart and gill samples were from juvenile animals and the mantle from hatchlings. Smaller animals have higher mass specific rates of oxygen consumption than larger animals (Johansen et al., 1982; Grigoriou and Richardson, 2009). It is likely that if oxygen consumption of mantle had been conducted on tissue removed from animals the same size as that used for gill and heart that the rate for mantle would have been relatively lower.

One of the major energy demanding processes in non-contracting cells is protein synthesis. In food deprived cuttlefish, both $\dot{M}O_2$ and protein synthesis are decreased in concert. A 36% decrease in whole animal $\dot{M}O_2$ was associated with a 63% decrease in fractional rate of protein synthesis in both mantle and gill (Lamarre et al., 2016). The importance of these findings in the context of the current study is that it demonstrates that protein synthesis is a regulatable element of the energy demand machinery in *S. officinalis*. Here we show that the inclusion of cycloheximide decreases $\dot{M}O_2$ in isolated gill, heart, and mantle preparations. Based on studies with goldfish and trout it is likely that the level of cycloheximide utilized here resulted in a total shut down of protein synthesis (Wieser and Krumschnabel, 2001), as previously suggest by the Agin et al. (2003) results on the species and because the amount used here was more than 3 times higher than the dosages used in *Octopus vulgaris* (Prozzo and Giuditta, 1973) and *Loligo pealii* (Giuditta et al., 1968), which resulted in 80% protein synthesis inhibition in the optic globe with 100 $\mu\text{g}/\text{mL}$ and 95–96% inhibition in squid axons with 200 $\mu\text{g}/\text{mL}$, respectively. If it is assumed that all tissues in the quiescent animals are consuming oxygen at similar rates, a total inhibition of all protein synthesis would be required to meet the whole animal decrease in $\dot{M}O_2$ of 37%. This is highly unlikely given that even in an anoxic resistant fish, oscar (*Astronotus crassipinnis*), protein synthesis is decreased by only 55% and 60–85% in muscle and liver, respectively, at a DO_2 of 10% (Lewis et al., 2007). Therefore, although a decrease in protein synthesis could potentially contribute to the decrease in whole animal $\dot{M}O_2$, there must be additional mechanisms to reduce energy expenditure.

Na^+/K^+ ATPase is a major driving force for many energy dependent ion transport processes and is considered one of the major cellular sites for ATP utilization.

Na^+/K^+ ATPase has been identified in numerous tissues of *S. officinalis* (Donaubauer, 1981) and is presumably ubiquitous. A ouabain sensitive Na^+/K^+ ATPase has been identified in gill of squid (*Doritheutis plei*) (Proverbio et al., 1988) and more recently, Na^+/K^+ ATPase in *Sepia* gill was shown to increase in response to increases in water CO_2 level (Hu et al., 2011). In the hypoxia tolerant intertidal clam, *Mercenaria mercenaria*, a severe hypoxic exposure resulted in a decrease in the maximal *in vitro* enzyme activity of Na^+/K^+ ATPase (Ivanina et al., 2016). As with protein synthesis, these studies illustrate that Na^+/K^+ ATPase is a site that can be controlled to alter energy

demand. Here it is shown with isolated tissues that inhibition of Na^+/K^+ ATPase with ouabain decreased $\dot{M}O_2$ by $\sim 20\%$ in mantle and 50–60% in gill and heart. Given that the Na^+/K^+ ATPase is one of the largest consumers of ATP in cells, it is likely that a curtailment of this process occurs in *S. officinalis* under hypoxia.

CONCLUSIONS

A hypoxic challenge to *S. officinalis* of 50% DO_2 results in only a minor activation of anaerobic metabolism as assessed by octopine accumulation. There were no changes in levels of HSP70 and polyubiquitinated proteins suggesting no alteration in rates of protein breakdown. The animal responds with an increase in ventilation and an overall decrease in metabolic rate, potentially due to decreases in both rates of protein synthesis and Na^+/K^+ ATPase activity. The relatively high level of HSP70 in gill compared to mantle and heart is a novel finding and may be related to the high rates of protein synthesis in this tissue.

AUTHOR CONTRIBUTIONS

SL, TM, AS, and WD participated in the conception and design of research; JC and AS performed the animal husbandry; all authors performed the experiments, acquired and analysed data of work; SL, TM, AS, and WD performed interpretation of data; all authors drafted, edited and approved the final version of the manuscript. All authors agree to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

FUNDING

TM, SL, and WD were supported by Natural Sciences and Engineering Research Council of Canada (NSERC) Discovery grants. WRD holds the Canada Research Chair in Marine Bioscience. LT held a New Brunswick Innovation Foundation (NBIF) STEM Graduate scholarship. AVS was supported by Fundação para a Ciência e a Tecnologia through Programa Investigador FCT 2014 (IF/00576/2014) and Pluriannual funding to CCMAR (UID/Multi/04326/2013). This work is a contribution to the COST (European COoperation on Science and Technology) Action FA1301 “A network for improvement of cephalopod welfare and husbandry in research, aquaculture and fisheries (CephsInAction)” that partially funded the publication of this manuscript.

ACKNOWLEDGMENTS

The authors thank Dr. Juan Fuentes, Dr. Pedro Guerreiro, Mr. João Reis and the students and staff at Ramalhete Station for valuable input and logistical assistance with the study.

REFERENCES

- Agin, V., Chichery, R., Maubert, E., and Chichery, M. P. (2003). Time-dependent effects of cycloheximide on long-term memory in the cuttlefish. *Pharmacol. Biochem. Behav.* 75, 141–146. doi: 10.1016/S0091-3057(03)00041-8
- Balchin, D., Hayer-Hartl, M., and Hartl, F. U. (2016). *In vivo* aspects of protein folding and quality control. *Science* 353, aac4354-1–aac4354-13. doi: 10.1126/science.aac4354
- Bloor, I. S. M., Attrill, M. J., and Jackson, E. L. (2013). A review of the factors influencing spawning, early life stage survival and recruitment variability in the common cuttlefish (*Sepia officinalis*). *Adv. Mar. Biol.* 65, 1–65. doi: 10.1016/B978-0-12-410498-3.00001-X
- Boal, J. G., and Golden, D. K. (1999). Distance chemoreception in the common cuttlefish, *Sepia officinalis* (Mollusca, Cephalopoda). *J. Exp. Mar. Biol. Ecol.* 235, 307–317. doi: 10.1016/S0022-0981(98)00187-7
- Boal, J. G., and Ni, J. N. (1996). Ventilation rate of cuttlefish, *Sepia officinalis*, in response to visual stimuli. *Veliger* 39, 342–347.
- Bone, Q., Brown, E. R., and Travers, G. (1994). On the respiratory plow in the cuttlefish *Sepia officinalis*. *J. Exp. Biol.* 194, 153–165.
- Bone, Q., Pulsford, A., and Chubb, A. D. (1981). Squid mantle muscle. *J. Mar. Biol. Asso. U.K.* 61, 327–342. doi: 10.1017/S0025315400046981
- Boyle, P. R. (1991). *The UFAW Handbook on the Care and Management of the Cephalopods in the Laboratory*. Herts: UFAW.
- Brett, J. R. (1979). “Environmental Factors and Growth,” in *Fish Physiology*, eds D. J. R. W. S. Hoar and J. R. Brett (New York, NY: Academic Press), 599–675.
- Brix, O., Colosimo, A., and Giardina, B. (1994). Temperature dependence of oxygen binding to cephalopod haemocyanins: ecological implications. *Mar. Freshw. Behav. Physiol.* 25, 149–162. doi: 10.1080/10236249409378914
- Burt, K., Hamoutene, D., Perez-Casanova, J., Kurt Gamperl, A., and Volkoff, H. (2013). The effect of intermittent hypoxia on growth, appetite and some aspects of the immune response of Atlantic salmon (*Salmo salar*). *Aquac. Res.* 45, 124–137. doi: 10.1111/j.1365-2109.2012.03211.x
- Carter, C. G., Lynch, K. A., and Moltschanivskiy, N. A. (2009). Protein synthesis in a solitary benthic cephalopod, the Southern dumpling squid (*Euprymna tasmanica*). *Comp. Biochem. Physiol. A Mol. Integr. Physiol.* 153, 185–190. doi: 10.1016/j.cbpa.2009.02.015
- Castro, B. G., Garrido, J. L., and Sotelo, C. G. (1992). Changes in composition of digestive gland and mantle muscle of the cuttlefish *Sepia officinalis* during starvation. *Mar. Biol.* 114, 11–20.
- Cerezo Valverde, J., and García García, B. (2005). Suitable dissolved oxygen levels for common octopus (*Octopus vulgaris* Cuvier, 1797) at different weights and temperatures: analysis of respiratory behaviour. *Aquaculture* 244, 303–314. doi: 10.1016/j.aquaculture.2004.09.036
- Chan, F., Barth, J. A., Lubchenko, J., Kirinich, A., Weeks, H., Peterson, W. T., et al. (2008). Emergence of anoxia in the California current large marine ecosystem. *Science* 319, 920–920. doi: 10.1126/science.1149016
- Cosme, N., and Hauschild, M. Z. (2016). Effect factors for marine eutrophication in LCIA based on species sensitivity to hypoxia. *Ecol. Indic.* 69, 453–462. doi: 10.1016/j.ecolind.2016.04.006
- Delaney, M. A., and Klesius, P. H. (2004). Hypoxic conditions induce HSP70 production in blood, brain and head kidney of juvenile Nile tilapia *Oreochromis niloticus* (L.). *Aquaculture* 236, 633–644. doi: 10.1016/j.aquaculture.2004.02.025
- De Wachter, B., Wolf, G., Richard, A., and Declair, W. (1988). Regulation of respiration during juvenile development of *Sepia officinalis* (Mollusca: Cephalopoda). *Mar. Biol.* 97, 365–371. doi: 10.1007/BF00397767
- Donaubauer, H. H. (1981). Sodium-activated and potassium-activated adenosine-triphosphatase in the excretory organs of *Sepia officinalis* (Cephalopoda). *Mar. Biol.* 63, 143–150. doi: 10.1007/BF00406822
- Ekau, W., Auel, H., Pörtner, H. O., and Gilbert, D. (2010). Impacts of hypoxia on the structure and processes in pelagic communities (zooplankton, macro-invertebrates and fish). *Biogeosciences* 7, 1669–1699. doi: 10.5194/bg-7-1669-2010
- Finke, E., Portner, H. O., Lee, P. G., and Webber, D. M. (1996). Squid (*Lolliguncula brevis*) life in shallow waters: oxygen limitation of metabolism and swimming performance. *J. Exp. Biol.* 199, 911–921.
- Fiorito, G., Affuso, A., Basil, J., Cole, A., De Girolamo, P., D’angelo, L., et al. (2015). Guidelines for the care and welfare of cephalopods in research – a consensus based on an initiative by CephRes, FELASA and the Boyd Group. *Lab. Anim.* 49, 1–90. doi: 10.1177/0023677215580006
- Gäde, G. (1985). “Octopine,” in *Methods of Enzymatic Analysis*, ed H. U. Bergmeyer (New York, NY: VCH Publishers Ltd.), 419–424.
- Giuditta, A., Dettbarn, W. D., and Brzin, M. (1968). Protein synthesis in the isolated giant axon of the squid. *Proc. Natl. Acad. Sci. U.S.A.* 59, 1284–1287. doi: 10.1073/pnas.59.4.1284
- Gonçalves, R. A., Aragao, C., Frias, P. A., and Sykes, A. V. (2012). The use of different anaesthetics as welfare promoters during short-term human manipulation of European cuttlefish (*Sepia officinalis*) juveniles. *Aquaculture* 370, 130–135. doi: 10.1016/j.aquaculture.2012.10.014
- Grantham, B. A., Chan, F., Nielsen, K. J., Fox, D. S., Barth, J. A., Huyer, A., et al. (2004). Upwelling-driven nearshore hypoxia signals ecosystem and oceanographic changes in the northeast Pacific. *Nature* 429, 749–754. doi: 10.1038/nature02605
- Grigoriou, P., and Richardson, C. A. (2009). Effect of body mass, temperature and food deprivation on oxygen consumption rate of common cuttlefish *Sepia officinalis*. *Mar. Biol.* 156, 2473–2481. doi: 10.1007/s00227-009-1272-4
- Houlihan, D. F., Innes, A. J., Wells, M. J., and Wells, J. (1982). Oxygen consumption and blood gases of *Octopus vulgaris* in hypoxic conditions. *J. Comp. Physiol.* 148, 35–40. doi: 10.1007/BF00688885
- Houlihan, D. F., Mcmillan, D. N., Agnisola, C., Genoino, I. T., and Foti, L. (1990). Protein synthesis and growth in *Octopus vulgaris*. *Mar. Biol.* 106, 251–259. doi: 10.1007/BF01314808
- Hu, M. Y., Tseng, Y. C., Stumpp, M., Gutowska, M. A., Kiko, R., Lucassen, M., et al. (2011). Elevated seawater PCO₂ differentially affects branchial acid-base transporters over the course of development in the cephalopod *Sepia officinalis*. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 301, R559–R559. doi: 10.1152/ajpregu.00653.2010
- Ivanina, A. V., Nesmelova, I., Leamy, L., Sokolov, E. P., and Sokolova, I. M. (2016). Intermittent hypoxia leads to functional reorganization of mitochondria and affects cellular bioenergetics in marine molluscs. *J. Exp. Biol.* 219, 1659–1674. doi: 10.1242/jeb.134700
- Johansen, K., Brix, O., Kornerup, S., and Lykkeboe, G. (1982). Factors affecting O₂-uptake in the cuttlefish, *Sepia officinalis*. *J. Mar. Biol. Asso. U.K.* 62, 187–191. doi: 10.1017/S0025315400020208
- Kepler, D., and Decker, K. (1974). “Glycogen – determination with amyloglucosidase,” in *Methods of Enzymatic Analysis*, ed H. U. Bergmeyer (New York, NY: Academic Press), 1127–1131.
- Lamarre, S. G., Ditlecadet, D., Mckenzie, D. J., Bonnaud, L., and Driedzic, W. R. (2012). Mechanisms of protein degradation in mantle muscle and proposed gill remodeling in starved *Sepia officinalis*. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 303, R427–R437. doi: 10.1152/ajpregu.00077.2012
- Lamarre, S. G., Maccormack, T. J., Sykes, A. V., Hall, J. R., Speers-Roesch, B., Callaghan, N. I., et al. (2016). Metabolic rate and rates of protein turnover in food deprived cuttlefish, *Sepia officinalis* (Linnaeus 1758). *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 310, R1160–R1168. doi: 10.1152/ajpregu.00459.2015
- Lampinen, J., Raitio, M., Peraia, A., and Oranen, H. (2012). *Microplate Based Pathlength Correction Method for Photometric DNA Quantification Assay*. Vantaa: Thermo Fisher Application Note.
- Lefevre, S., Bayley, M., and Mckenzie, D. J. (2016). Measuring oxygen uptake in fishes with bimodal respiration. *J. Fish Biol.* 88, 206–231. doi: 10.1111/jfb.12698
- Lewbart, G. A., and Mosley, C. (2012). Clinical anesthesia and analgesia in invertebrates. *J. Exot. Pet Med.* 21, 59–70. doi: 10.1053/j.jepm.2011.11.007
- Lewis, J. M., Costa, I., Val, A. L., Almeida-Val, V. M. F., Gamperl, A. K., and Driedzic, W. R. (2007). Responses to hypoxia and recovery: repayment of oxygen debt is not associated with compensatory protein synthesis in the Amazonian cichlid, *Astronotus ocellatus*. *J. Exp. Biol.* 210, 1935–1943. doi: 10.1242/jeb.005371
- Maccormack, T. J., Callaghan, N. I., Sykes, A. V., and Driedzic, W. R. (2016). Taurine depresses cardiac contractility and enhances systemic heart glucose utilization in the cuttlefish, *Sepia officinalis*. *J. Comp. Physiol. B* 186, 215–227. doi: 10.1007/s00360-015-0946-0
- Melzner, F., Bock, C., and Portner, H. O. (2006). Temperature-dependent oxygen extraction from the ventilatory current and the costs of ventilation in the cephalopod *Sepia officinalis*. *J. Comp. Physiol. B* 176, 607–621. doi: 10.1007/s00360-006-0084-9

- Melzner, F., Bock, C., and Portner, H. O. (2007). Allometry of thermal limitation in the cephalopod *Sepia officinalis*. *Comp. Biochem. Physiol. A Mol. Integr. Physiol.* 146, 149–154. doi: 10.1016/j.cbpa.2006.07.023
- Moltschaniwskij, N. A., and Carter, C. G. (2010). Protein synthesis, degradation, and retention: mechanisms of indeterminate growth in cephalopods. *Physiol. Biochem. Zool.* 83, 997–1008. doi: 10.1086/656387
- Mommsen, T. P., Ballantyne, J., Macdonald, D., Gosline, J., and Hochachka, P. W. (1981). Analogues of red and white muscle in squid mantle. *Proc. Natl. Acad. Sci. U.S.A.* 78, 3274–3278. doi: 10.1073/pnas.78.5.3274
- Pierce, G. J., Valavanis, V. D., Guerra, A., Jereb, P., Orsi-Relini, L., Bellido, J. M., et al. (2008). A review of cephalopod-environment interactions in European seas. *Hydrobiologia* 612, 49–70. doi: 10.1007/s10750-008-9489-7
- Postel, U., Becker, W., Brandt, A., Luck-Kopp, S., Riestenpatt, S., Weihrauch, D., et al. (2000). Active osmoregulatory ion uptake across the pleopods of the isopod *Idotea baltica* (Pallas): electrophysiological measurements on isolated split endo- and exopodites mounted in a micro-ussing chamber. *J. Exp. Biol.* 203, 1141–1152. Available online at: <http://jeb.biologists.org/content/203/7/1141.article-info>
- Proverbio, T., Marín, R., and Proverbio, F. (1988). Ouabain-insensitive, Na⁺-stimulated ATPase activity in squid gill microsomes. *Comp. Biochem. Physiol. B* 90, 341–345. doi: 10.1016/0305-0491(88)90084-3
- Prozzo, N., and Giuditta, A. (1973). Protein synthesis in the brain of *Octopus vulgaris*, Lam. *J. Neurochem.* 21, 1327–1336. doi: 10.1111/j.1471-4159.1973.tb07586.x
- Schmidtke, S., Stramma, L., and Visbeck, M. (2017). Decline in global oceanic oxygen content during the past five decades. *Nature* 542, 335–339. doi: 10.1038/nature21399
- Seibel, B. A., Hafker, N. S., Trübenbach, K., Zhang, J., Tessier, S. N., Portner, H. O., et al. (2014). Metabolic suppression during protracted exposure to hypoxia in the jumbo squid, *Dosidicus gigas*, living in an oxygen minimum zone. *J. Exp. Biol.* 217, 2555–2568. doi: 10.1242/jeb.100487
- Speers-Roesch, B., Callaghan, N. I., McCormack, T. J., Lamarre, S. G., Sykes, A. V., and Driedzic, W. R. (2016). Enzymatic capacities of metabolic fuel use in cuttlefish (*Sepia officinalis*) and responses to food deprivation: insight into the metabolic organization and starvation survival strategy of cephalopods. *J. Comp. Physiol. B* 186, 711–725. doi: 10.1007/s00360-016-0991-3
- Storey, K. B. (1977). Tissue specific isozymes of octopine dehydrogenase in cuttlefish, *Sepia officinalis* - Roles of octopine dehydrogenase and lactate dehydrogenase in *Sepia*. *J. Comp. Physiol.* 115, 159–169. doi: 10.1007/BF00692527
- Storey, K. B., and Storey, J. M. (1979). Octopine metabolism in the cuttlefish, *Sepia officinalis* - octopine production by muscle and its role as an aerobic substrate for non-muscular tissues. *J. Comp. Physiol.* 131, 311–319. doi: 10.1007/BF00688806
- Storey, K. B., Storey, J. M., Johansen, K., and Hochachka, P. W. (1979). Octopine metabolism in *Sepia officinalis* - Effect of hypoxia and metabolite loads on the blood-levels of octopine and related-compounds. *Can. J. Zool.* 57, 2331–2336. doi: 10.1139/z79-303
- Sykes, A. V., Baptista, F. D., Gonçalves, R. A., and Andrade, J. P. (2012). Directive 2010/63/EU on animal welfare: a review on the existing scientific knowledge and implications in cephalopod aquaculture research. *Rev. Aquaculture* 4, 142–162. doi: 10.1111/j.1753-5131.2012.01070.x
- Sykes, A. V., Domingues, P., and Andrade, J. P. (2014). “*Sepia officinalis*,” in *Cephalopod Culture*, eds J. Iglesias, L. Fuentes, and R. Villanueva (Dordrecht: Springer), 175–204. doi: 10.1007/978-94-017-8648-5_11
- Townhill, B. L., Pinnegar, J. K., Righton, D. A., and Metcalfe, J. D. (2017). Fisheries, low oxygen and climate change: how much do we really know? *J. Fish Biol.* 90, 723–750. doi: 10.1111/jfb.13203
- Trübenbach, K., Da Costa, G., Ribeiro-Silva, C., Ribeiro, R. M., Cordeiro, C., and Rosa, R. (2014). Hypoxia-driven selective degradation of cellular proteins in jumbo squids during diel migration to the oxygen minimum zones. *Mar. Biol.* 161, 575–584. doi: 10.1007/s00227-013-2360-z
- Trübenbach, K., Pegado, M. R., Seibel, B. A., and Rosa, R. (2013a). Ventilation rates and activity levels of juvenile jumbo squid under metabolic suppression in the oxygen minimum zone. *J. Exp. Biol.* 216, 359–368. doi: 10.1242/jeb.072587
- Trübenbach, K., Teixeira, T., Diniz, M., and Rosa, R. (2013b). Hypoxia tolerance and antioxidant defense system of juvenile jumbo squids in oxygen minimum zones. *Deep Sea Res. II Top. Stud. Oceanogr.* 95, 209–217. doi: 10.1016/j.dsr2.2012.10.001
- Van Den Thillart, G., Dalla Via, J., Vitali, G., and Cortesi, P. (1994). Influence of long-term hypoxia exposure on the energy metabolism of *Solea solea*. I. Critical O₂ levels for aerobic and anaerobic metabolism. *Mar. Ecol. Prog. Ser.* 104, 109–117. doi: 10.3354/meps104109
- Vaquero-Sunyer, R., and Duarte, C. M. (2008). Thresholds of hypoxia for marine biodiversity. *Proc. Natl. Acad. Sci. U.S.A.* 105, 15452–15457. doi: 10.1073/pnas.0803833105
- Wieser, W., and Krumschnabel, G. (2001). Hierarchies of ATP-consuming processes: direct compared with indirect measurements, and comparative aspects. *Biochem. J.* 355, 389–395. doi: 10.1042/bj3550389
- Zar, J. H. (1999). *Biostatistical Analysis*. Upper Saddle River, NY: Prentice-Hall Inc.

Conflict of Interest Statement: The handling Editor declared a past co-authorship with one of the authors AS and states that the process nevertheless met the standards of a fair and objective review.

The other authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2017 Capaz, Tunnah, McCormack, Lamarre, Sykes and Driedzic. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.