

ORIGINAL ARTICLE

High diversity of *Phytophthora* species in natural ecosystems and nurseries of Portugal: Detrimental side effect of plant introductions from the age of discovery to modern globalization

Marília Horta Jung^{1,2}  | Cristiana Maia³  | Beatriz Mora-Sala⁴  |
 Paloma Abad-Campos⁴  | Leonardo Schena⁵  | Saveria Mosca⁵  | Giuseppe Carella⁶  |
 Salvatore Moricca⁶  | Jan Nechwatal⁷  | Lídia Dionísio⁸  | Alfredo Cravador⁸  |
 Thomas Jung^{1,2} 

¹Department of Forest Protection and Wildlife Management, Phytophthora Research Centre, Faculty of Forestry and Wood Technology, Mendel University in Brno, Brno, Czech Republic

²Phytophthora Research and Consultancy, Nußdorf, Germany

³Centre of Marine Sciences (CCMAR), University of Algarve, Faro, Portugal

⁴Instituto Agroforestal Mediterráneo, Universitat Politècnica de València, Valencia, Spain

⁵Department of Agriculture, Mediterranean University of Reggio Calabria, Reggio Calabria, Italy

⁶Department of Agri-Food Production and Environmental Sciences, Plant Pathology and Entomology Division, University of Florence, Florence, Italy

⁷Bavarian State Research Center for Agriculture, Institute for Plant Protection, Freising, Germany

⁸MED—Mediterranean Institute for Agriculture, Environment and Development & CHANGE—Global Change and Sustainability Institute, University of Algarve, Faro, Portugal

Correspondence

Thomas Jung, Mendel University in Brno, Faculty of Forestry and Wood Technology, Department of Forest Protection and Wildlife Management, Phytophthora Research Centre, 613 00 Brno, Czech Republic.

Email: thomas.jung@mendelu.cz; dr.t.jung@gmail.com

Funding information

Fundação para a Ciência e a Tecnologia (FCT), Grant/Award Number: BIODIVERSA/0002/2012; European Regional Development Fund, Grant/Award Number: CZ.02.1.01/0.0/0.0/15_003/0000453; Italian Ministry of Education, University and Research (MIUR), Grant/Award Number: MIUR-FIRB 2010 (RBFR10PZ4N)

Abstract

Between 2010 and 2015, an extensive survey of *Phytophthora* diversity was performed across Portugal. Baiting and direct isolation tests obtained 48 *Phytophthora* taxa in 62 of 70 forest stands (89%) from 148 of 302 rhizosphere soil samples (49%) and 25 of 29 tree species (86%), and from bark cankers of *Alnus lusitanica* and *Fagus sylvatica*; from 33 of 34 rivers and forest streams (97%); and in 12 of 13 forest nurseries (92%) from 54 of 87 root ball samples (61%) and 16 of 21 tree species (76%). The 799 *Phytophthora* isolates belonged to 28 known and three recently described species; one designated and two new non-hybrid taxa; and five designated and nine new hybrid taxa. The majority, 34 *Phytophthora* taxa (71%), are introduced aliens in Europe while nine (19%) and five (10%) taxa are of cryptic and native origin, respectively. *Phytophthora cinnamomi* was the most common species in forests (56%) and forest nurseries (61%). Multiple first records for Portugal and partly Europe of both *Phytophthora* taxa and tree declines and new host–*Phytophthora* associations were found. Isolation rates from forest soil differed considerably between declining (65%)

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial-NoDerivs](https://creativecommons.org/licenses/by-nc-nd/4.0/) License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2024 The Author(s). *Plant Pathology* published by John Wiley & Sons Ltd on behalf of British Society for Plant Pathology.

and healthy trees (10%), regardless of the tree species. The widespread infestations of nurseries with aggressive wide-host range pathogens like *P. cinnamomi*, *P. pseudocryptogea*, *P. xcambivora*, *P. multivora* and *P. plurivora*, and host-specific pathogens like *P. alticola*, *P. quercetorum* and *P. quercina*, will inevitably cause their further spread to the wider environment, exacerbating pathogen threats to forest ecosystems and thus weakening their resilience to climatic extremes.

KEYWORDS

biosecurity, EU Nature Restoration Regulation, hybridization, invasive pathogens, nursery infestations, tree declines

1 | INTRODUCTION

The genus *Phytophthora* (Oomycota, Stramenopiles and Peronosporaceae) currently comprises more than 260 described species and numerous informally designated taxa organized in 14 phylogenetic clades with high biological cohesion (Abad et al., 2023; Brasier et al., 2022; Chen et al., 2022; Jung et al., 2022, 2024). Many devastating declines of horticultural crops, forest trees and natural ecosystems are driven by exotic *Phytophthora* species which, due to coevolution with their native host plants, remain unnoticed in their native habitats. However, following their introduction to other biogeographic regions, they often become invasive, threatening a non-coevolved flora containing a high number of susceptible species (Chen et al., 2022; Erwin & Ribeiro, 1996; Jung, Pérez-Sierra, et al., 2018; Lamour, 2013). Since the 1960s, the number of important *Phytophthora* diseases of forest trees and natural ecosystems has increased exponentially from five to currently 41 (Brasier et al., 2022). Several disease syndromes of forests and other natural ecosystems, namely dieback of eucalypt forests and *Banksia* heathlands in Australia, oak declines in Europe and the United States, beech decline in Europe, and decline and dieback of Mediterranean maquis vegetation in Italy and California, involve pathogens from multiple (up to nine) *Phytophthora* clades (Bourret et al., 2023; Brasier et al., 2022; Frankel et al., 2020; Jung, Pérez-Sierra, et al., 2018; Scanu et al., 2015). An array of *Phytophthora* species stand out due to their high invasiveness and their aggressiveness to either a wide range of host species (*P. cactorum*, *P. cinnamomi*, *P. palmivora*, *P. multivora*, *P. plurivora*, *P. ramorum* and *P. xcambivora*) or to specific hosts (*P. austrocedri*, *P. lateralis* and *P. xalni*) (Brasier et al., 2004, 2022; Brasier & Webber, 2010; Drenth & Guest, 2004; Erwin & Ribeiro, 1996; Greslebin et al., 2007; Grünwald et al., 2012; Hansen et al., 2000; Hardham & Blackman, 2018; Jung & Burgess, 2009; Jung, Pérez-Sierra, et al., 2018; Scott et al., 2009). Recently, worldwide surveys in natural ecosystems and molecular population analyses have unveiled the geographic origins of several notorious *Phytophthora* species: *P. cactorum* in North America; *P. cinnamomi* in Southeast Asia; *P. kernoviae* in Chile; *P. lateralis* in Taiwan; *P. multivora* in South Africa; *P. plurivora*, *P. ramorum* and *P. xcambivora* in East Asia; and *P. pseudosyringae* in Europe (Bourret et al., 2022; Brasier et al., 2012; Jung et al., 2021, 2022, 2024; Jung,

Durán, et al., 2018; Mullett et al., 2023, 2024; Shakya et al., 2021; Tsykun et al., 2022).

The intercontinental trade in living plants has been identified as the main pathway for accidental introductions of plant pathogens, including *Phytophthoras*, from other continents (Brasier, 2008; Jung et al., 2016; Liebhold et al., 2012; Santini et al., 2013). During the past two decades, several studies demonstrated almost ubiquitous infestations of nurseries with a wide range of *Phytophthora* pathogens and presented strong evidence for the planting of infested nursery stock constituting the major pathway of invasive harmful *Phytophthora* pathogens to the wider environment (Bačova et al., 2024; Bienapfl & Balci, 2014; Frankel et al., 2020; Garbelotto et al., 2018; Jung et al., 2016; Leonberger et al., 2013; Migliorini et al., 2015; Moralejo et al., 2009; Mora-Sala et al., 2022; Prigigallo et al., 2015, 2016; Rooney-Latham et al., 2019; Sims & Garbelotto, 2021).

Situated along the Atlantic in the southwesternmost corner of Europe, mainland Portugal has a long history of maritime exploration and overseas trade going back to the early 15th Century. Following the establishment of colonies and trading posts during the 16th Century in Brazil, along the West African coast (Cape Verde, Guinea Bissau, São Tomé and Príncipe, Angola) and the East African Swahili coast (e.g., Mozambique, Mombasa, Zanzibar), in India (Goa, Mumbai), the Malay archipelago (Malacca, the Moluccas, Timor), China (Macau), Taiwan and Japan (Nagasaki) (Birmingham, 2000; Disney, 2009; Page, 2013), numerous ornamental and horticultural plants were introduced to Portugal over centuries. With 667 plant taxa, 17.9% of the total Portuguese flora, originating from other biogeographic regions and more than 300 alien plant species first recorded for Europe, Portugal is a major gateway country for alien plant introductions, ranking third behind the UK and Belgium (Arianoutsou et al., 2021; Domingues de Almeida & Freitas, 2012). Consequently, it can be assumed that a rich diversity of non-native *Phytophthora* species has been introduced as hitchhikers of the intense plant trade to Portugal. Indeed, unambiguous symptoms of ink disease of sweet chestnut (*Castanea sativa*) were first reported in Portugal in 1838 (Crandall et al., 1945), suggesting the causal agents *P. cinnamomi* and *P. xcambivora* were introduced and established long before the genus *Phytophthora* had been erected in 1876 by de Bary (Chen et al., 2022). Presence of both pathogens in Portuguese

chestnut and cork oak (*Quercus suber*) stands was unveiled in the 1940s (Lopes-Pimentel, 1946, 1947). Fifty years later, Brasier et al. (1993) demonstrated the involvement of *P. cinnamomi* in the widespread oak decline in Portugal and Spain, which was later confirmed by other studies (e.g., Gallego et al., 1999; Jung et al., 2016; Moreira & Martins, 2005).

Worldwide, oomycete surveys of waterbodies and riparian ecosystems have revealed an astonishing diversity of *Phytophthora* species, including—in addition to numerous primarily aquatic species—many soilborne and airborne primary pathogens (Corcobado et al., 2023; Hüberli et al., 2013; Jung et al., 2019, 2020, 2022, 2024; Jung, Chang, Bakonyi, et al., 2017; Jung, Durán, et al., 2018; Oh et al., 2013; Reeser et al., 2011; Shrestha et al., 2013). In Portugal, *P. xalni*, the main causal agent of root and collar rot and mortality of alders, was first reported in 2016 (Jung et al., 2016; Kanoun-Boulé et al., 2016) while a small-scale survey in five riparian stands demonstrated the presence of 11 *Phytophthora* species, including *P. multivora*, *P. plurivora* and *P. rosacearum* (Bregant et al., 2023a).

This paper reports on the results of a survey of *Phytophthora* diversity performed between May 2010 and October 2015 in a diverse range of natural and planted forests, river systems and forest nurseries across Portugal, and on the association of *Phytophthora* spp. with disease symptoms of forest trees. Many nursery results were previously included in the extensive dataset from a Europe-wide survey of almost 2000 nursery stands (Jung et al., 2016) but updated molecular species identifications and DNA sequences, detailed isolation methods, infestation rates, host plants and disease symptoms are given here for the first time.

2 | MATERIALS AND METHODS

2.1 | Sampling and *Phytophthora* isolation

Seventy forest sites covering a wide range of forest types and tree species, 39 riparian sites along 34 rivers and forest streams and 13 forest nurseries across Portugal were selected for sampling. The locations of the forest and riparian sites are shown in Figure 1 and their GPS coordinates are given in Tables 1 and 2. Because anonymity was a precondition for all participating nursery owners, no precise locations and names of nurseries are provided. A selection of forest types sampled is shown in Figure 2.

With the forest sites, soil sampling and isolation tests were performed according to Jung (2009) and Pérez-Sierra et al. (2013, 2022). In total, 302 rhizosphere soil samples were taken from mature trees of 29 species in 63 natural and semi-natural forest stands, two stands with a mixture of mature cork oaks and planted younger cork oaks (F45) or *Arbutus unedo* (F67), and five agroforestry plantations of *C. sativa*, *Juglans regia*, *Pinus pinea* and *Q. suber* (F12, F18, F42, F57 and F68) (Figure 1; Table 1). After removal of the organic surface layer, three soil monoliths with a size of 20×20×30cm were taken from the rhizosphere of each tree, at a distance of 30–150cm from the stem base. Aliquots of soil together with fine and coarse roots

(diameter ≤5mm) from all monoliths were bulked, and subsamples of approximately 500–750mL were spread in a plastic container or tub (c. 18×25–27×13–14cm) and flooded with distilled water so that the waterline was around 2–3cm above the soil surface. After removal of floating debris with tweezers and tissue paper, 3- to 10-day-old leaflets of *Q. suber*, *Q. pyrenaica*, *Q. robur* and *Ceratonia siliqua* were floated as baits on the water covering the whole surface. Isolations were carried out at 18–20°C in an air-conditioned room in natural daylight. Brownish leaflets were examined at 80× under a light microscope for the presence of *Phytophthora* sporangia. Infected leaflets were blotted dry, cut into small segments and plated onto selective PARPNH agar (V8-juice agar [V8A] amended with 10µg/mL pimaricin, 200µg/mL ampicillin, 10µg/mL rifampicin, 25µg/mL pentachloronitrobenzene [PCNB], 50µg/mL nystatin and 50µg/mL hymexazol; Jung, 2009; Pérez-Sierra et al., 2022).

Bleeding bark cankers were sampled from six *Alnus lusitanica* trees growing in four riparian forests (F22, F23, F28 and F41) on the riverbanks of the Tâmega, Fervença, Alva and Côa Rivers, and from one *F. sylvatica* tree in a montane forest (F37) in the Parque Nacional da Serra da Estrela (Figures 1 and 3). Samples were taken from the upper 20cm of active lesions with orange-brown discoloration of the inner bark, placed in distilled water and transported to the laboratory where the water was replaced four times per day over 2–3 days in order to remove excess polyphenols. Small pieces (c. 8×3×3mm) were cut from the lesion, blotted on filter paper and plated onto selective PARPNH agar.

Phytophthora isolations from the 34 rivers and forest streams were performed using an in-situ baiting technique (Jung et al., 2019; Jung, Chang, Bakonyi, et al., 2017; Jung, Durán, et al., 2018). At each of the 39 sites, 15–20 non-wounded young leaves of *Q. suber*, *C. siliqua* and *Citrus sinensis* were placed as baits in a 25×30cm raft, prepared using nylon fly mesh and styrofoam, and the raft put to float at a place with calm water flow (Figure 2h). The rafts were collected after 3–4 days. Baiting leaves were washed in distilled water and blotted on filter paper. Five to 10 pieces (c. 2×2mm) were cut from each necrotic lesion of each leaf, blotted again on filter paper and plated onto PARPNH agar.

In the 13 forest nurseries, 87 bulked samples (1–13 bulked samples per nursery), each consisting of the root balls (= root systems and adhering soil) of three to six 1- to 3-year-old container plants from the same species, were harvested from 21 tree species. If present, seedlings with symptoms such as wilting, dieback, chlorosis and root and collar rot were sampled preferentially. In the laboratory, all root systems of a bulked sample were opened up and spread in a plastic container or tub (c. 18×25–27×13–14cm), flooded with distilled water and baited in the same way as the forest soil samples. In addition, in one nursery a *C. sativa* seedling with a fresh collar rot was sampled and the necrotic bark tissue plated onto PARPNH agar as described before.

All PARPNH Petri dishes with plated leaf or bark segments were incubated at 20°C in the dark and examined for *Phytophthora*-like hyphae after 24–48h. Pure cultures were obtained by transferring single hyphal tips from the edge of the colonies onto V8A. Stock




FIGURE 1 Location of 68 forest stands (F01–F68; red triangles) and 39 riparian sites (R01–R39; blue circles) along 34 rivers and streams included in the *Phytophthora* survey in mainland Portugal. Forest stands F69 (São Miguel Island) and F70 (Madeira Island) are not shown. Small maps provide detailed locations of the sites in (a) Parque Nacional da Peneda-Gerês; (b) Mata Nacional do Buçaco; (c) Parque Nacional da Serra da Estrela; (d) Reserva Natural da Serra da Malcata; and (e) Parque Natural Sintra-Cascais. For GPS coordinates see [Tables 1 and 2](#).



TABLE 1 Location, altitude, geological substrate and vegetation of 70 forest sites sampled in Portugal, sampled tree species and their health condition, and *Phytophthora* and other oomycete taxa isolated.

Site no.	GPS coordinates	Altitude (m.a.s.l.)	Location; parish, municipality, district	Geological substrate	Vegetation	Sampled tree species and health condition (no. of <i>Phytophthora</i> -positive/sampled trees) ^a	<i>Phytophthora</i> spp. (no. of positive samples) ^{b-e}
F01	41°43.110' N, 08°09.177' W	711	Parque Nacional da Peneda-Gerês (PNGP); Vilar da Veiga, Terras de Bouro, Braga	Granite	Submontane broadleaved forest	<i>Acer pseudoplatanus</i> DB (1/1) <i>A. pseudoplatanus</i> DC (0/1) <i>Arbutus unedo</i> DB (1/2) <i>Betula celtiberica</i> DB (1/2) <i>Quercus robur</i> (0/1) <i>Salix caprea</i> (1/1)	×CAM-DAPC9 (1) - SYR (1) ×CAM-DAPC9 (1) - ×CAM-DAPC9 (1)
F02	41°42.828' N, 08°08.703' W	775	PNGP; Vilar da Veiga, Terras de Bouro, Braga	Granite	Submontane birch forest	<i>B. celtiberica</i> (0/5)	-
F03	41°43.350' N, 08°07.765' W	726	PNGP; Vilar da Veiga, Terras de Bouro, Braga	Granite	Submontane broadleaved forest	<i>Quercus rubra</i> DB & bleeding bark canker (1/2)	CIN-PcG1 (1)
F04	41°41.948' N, 08°07.835' W	529	PNGP; Vilar da Veiga, Terras de Bouro, Braga	Granite	Submontane broadleaved forest	<i>Quercus suber</i> DB (1/1)	CIN-PcG2 (1)
F05	41°44.908' N, 08°09.221' W	597	PNGP; Vilar da Veiga, Terras de Bouro, Braga	Granite	Submontane broadleaved forest	<i>Castanea sativa</i> DC (1/2) <i>C. sativa</i> (0/2) <i>Q. robur</i> (0/1)	CIN-PcG1 (1) - -
F06	41°47.434' N, 08°08.235' W	699	PNGP; Campo do Gerês, Terras de Bouro, Braga	Granite	Submontane broadleaved forest	<i>Q. robur</i> DB (1/1)	QUE (1)
F07	41°48.253' N, 08°07.843' W	748	PNGP; Campo do Gerês, Terras de Bouro, Braga	Granite	Submontane mixed broadleaved–pine forest	<i>Quercus pyrenaica</i> (0/2) <i>Q. robur</i> (0/5) <i>Q. robur</i> DC & <i>Q. pyrenaica</i> DB (1/2) <i>Q. pyrenaica</i> DB & <i>Pinus pinaster</i> DB (1/1)	- - CIN (1) CIN (1), ×CAM-DAPC9 (1)
F08	41°46.083' N, 08°08.749' W	849	PNGP; Campo do Gerês, Terras de Bouro, Braga	Granite	Lower montane broadleaved forest	<i>Fagus sylvatica</i> DB (1/1)	CIN (1)
F09	41°47.208' N, 08°08.298' W	744	PNGP; Campo do Gerês, Terras de Bouro, Braga	Granite	Submontane broadleaved forest	<i>F. sylvatica</i> (0/4)	-
F10	41°42.771' N, 08°06.513' W	689	PNGP; Vilar da Veiga, Terras de Bouro, Braga	Granite	Submontane broadleaved forest	<i>Q. suber</i> DB (2/4)	CIN (2)
F11	41°46.756' N, 08°10.188' W	552	PNGP; Campo do Gerês, Terras de Bouro, Braga	Granite	Submontane broadleaved forest	<i>Q. robur</i> (0/5)	-
F12	41°53.620' N, 07°04.070' W	823	Perímetro Florestal Serra da Coroa (PFSC)/Parque Natural Montesinho (PNM); Tuizelo, Vinhais, Bragança	Phyllites, quartzites, vulcanites	Chestnut–walnut plantation	<i>C. sativa</i> DB & bleeding bark cankers (2/4) <i>Juglans regia</i> DC (0/1)	×CAM-DAPC9 (2) -

(Continues)

TABLE 1 (Continued)

Site no.	GPS coordinates	Altitude (m a.s.l.)	Location; parish, municipality, district	Geological substrate	Vegetation	Sampled tree species and health condition (no. of <i>Phytophthora</i> -positive/sampled trees) ^a	<i>Phytophthora</i> spp. (no. of positive samples) ^{b-e}
F13	41°53.783' N, 07°03.997' W	857	PFSC/PNIM; Tuizelo, Vinhais, Bragança	Phyllites, quartzites, vulcanites	Lower montane broadleaved forest	<i>Q. pyrenaica</i> DC (1/2) <i>Q. pyrenaica</i> (0/3)	xCAM-DAPC9 (1) -
F14	41°56.110' N, 07°02.070' W	826	PFSC/PNIM; União de Freguesias (UF) Moimenta e Montouto, Vinhais, Bragança	Phyllites, quartzites, vulcanites	Lower montane broadleaved forest	<i>C. sativa</i> DB (2/3) <i>C. sativa</i> (0/2) <i>Q. pyrenaica</i> DC (1/2)	xCAM-DAPC9 (2) - xCAM-DAPC9 (1)
F15	41°45.106' N, 07°36.599' W	561	UF Ardãos e Bobadela, Botiças, Vila Real	Granite	Submontane broadleaved forest	<i>Q. pyrenaica</i> DB (1/1) <i>Q. pyrenaica</i> (0/5)	xCAM-DAPC9 (1), QUE (1) -
F16	41°45.282' N, 06°52.352' W	1043	UF Castrelos e Carrazedo, Bragança, Bragança	Metagabro and schists	Montane broadleaved forest	<i>B. celtiberica</i> DB (4/4) <i>Prunus avium</i> (1/2) Streambed mud (1/1)	GON (2), PSE-DAPC1 (2) PSE-DAPC1 (1) GON (1)
F17	41°46.604' N, 06°54.116' W	890	UF Castrelos e Carrazedo, Bragança, Bragança	Metagabro and schists	Lower montane broadleaved forest	<i>Quercus ilex</i> DB (1/1) <i>Q. ilex</i> (0/2) <i>Q. ilex</i> DC & <i>Q. pyrenaica</i> DC (0/1) <i>Q. pyrenaica</i> DB (1/1)	xCAM-DAPC9 (1) - - PSE-DAPC1 (1)
F18	41°43.096' N, 06°55.949' W	871	Celas, Vinhais, Bragança	Schists, peralkaline vulcanites, phyllites, quartzites	Chestnut plantation	<i>C. sativa</i> DB & bleeding bark cankers (3/5)	CIN-PcG2 (1), xCAM-DAPC9 (2)
F19	41°41.702' N, 06°55.734' W	798	Celas, Vinhais, Bragança	Schists, peralkaline vulcanites, phyllites, quartzites	Submontane broadleaved forest and riparian gallery forest downhill of a chestnut plantation	<i>Q. pyrenaica</i> DB (1/4) <i>Alnus lusitanica</i> DC (1/1)	QUE (1) ^f CIN (1)
F20	41°31.530' N, 06°49.135' W	640	Vinhas, Macedo de Cavaleiros, Bragança	Metagabro and schists	Submontane open oak woodland	<i>Q. ilex</i> (0/4) <i>Q. suber</i> (0/3)	- -
F21	41°30.161' N, 06°48.512' W	713	Morais, Macedo de Cavaleiros, Bragança	Metagabro and schists	Submontane open oak woodland	<i>Q. suber</i> DB & bleeding bark cankers (2/5)	CIN-PcG2 (2)
F22	41°43.682' N, W07°28.620	350	Rio Tâmega; UF Madalena e Samações, Chaves, Vila Real	Quaternary alluvial sediments	Riparian forest	<i>A. lusitanica</i> bleeding bark canker (1/1)	xALN (1)
F23	41°48.201' N, 06°45.500' W	662	Rio Fervença; UF Santa Maria e Meixedo, Bragança, Bragança	Quaternary alluvial sediments	Riparian forest	<i>A. lusitanica</i> bleeding bark cankers (2/2)	xALN (2)

TABLE 1 (Continued)

Site no.	GPS coordinates	Altitude (m a.s.l.)	Location; parish, municipality, district	Geological substrate	Vegetation	Sampled tree species and health condition (no. of <i>Phytophthora</i> -positive/sampled trees) ^a	<i>Phytophthora</i> spp. (no. of positive samples) ^{b-e}
F24	40°22.596' N, 08°22.303' W	315	Mata Nacional do Buçaco (MNB); Luso, Mealhada, Aveiro	Schists, greywackes, quartzites, conglomerates	Colline broadleaved forest	<i>Phillyrea latifolia</i> DB (1/2) <i>P. latifolia</i> DC & <i>Ilex aquifolium</i> DC (1/1) <i>Q. robur</i> DB (2/4) <i>C. sativa</i> DB (1/2)	CIN (1) QUE (1) CIN (2) CIN-PcG2 (1)
F25	40°22.687' N, 08°22.055' W	339	MNB; Luso, Mealhada, Aveiro	Schists, greywackes, quartzites, conglomerates	Colline broadleaved forest	<i>Fraxinus angustifolia</i> (0/1)	- ^g
F26	40°22.240' N, 08°21.853' W	557	MNB; Luso, Mealhada, Aveiro	Schists, greywackes, quartzites, conglomerates	Colline to submontane broadleaved forest	<i>A. unedo</i> (0/1) <i>F. sylvatica</i> (0/1) <i>P. latifolia</i> DB (1/2) <i>Q. pyrenaica</i> (0/1) <i>Q. robur</i> DC (0/3) <i>Q. suber</i> (0/1)	- - CIN (1) - - -
F27	40°22.146' N, 08°21.572' W	524	MNB; Luso, Mealhada, Aveiro	Schists, greywackes, quartzites, conglomerates	Colline to submontane broadleaved forest	<i>Abies alba</i> (0/2) <i>Q. robur</i> DB (1/1)	- CIN-PcG2 (1)
F28	40°17.006' N, 07°57.554' W	192	Rio Alva; UF Vila Cova de Alva e Anseriz, Arganil, Coimbra	Quaternary alluvial sediments	Riparian gallery forest	<i>A. lusitanica</i> bleeding bark cankers (2/2)	xALN (2)
F29	40°22.914' N, 07°42.716' W	553	Parque Nacional da Serra da Estrela (PNSE); UF Seia, S. Romão e Lapa dos Dinheiros, Seia, Guarda	Biotide granitoid	Submontane broadleaved forest	<i>C. sativa</i> DC (0/1) <i>C. sativa</i> (0/2) <i>F. angustifolia</i> (0/1) <i>Q. robur</i> DC (1/2)	- - - CIN-PcG1 (1)
F30	40°21.300' N, 07°43.206' W	749	PNSE; Valezim, Seia, Guarda	Biotide granitoid	Submontane broadleaved forest	<i>C. sativa</i> DB (6/6)	CIN (5), QUE (3)
F31	40°20.514' N, 07°42.856' W	966	PNSE; Valezim, Seia, Guarda	Biotide granitoid	Lower montane broadleaved forest	<i>Q. rubra</i> (0/1)	-
F32	40°18.506' N, 07°45.206' W	464	PNSE, Rio Lima; UF Vide e Cabeça, Seia, Guarda	Biotide granitoid	Submontane broadleaved forest and riparian gallery forest	<i>A. lusitanica</i> DB (1/1) <i>Prunus lusitanica</i> DC (1/1) <i>S. caprea</i> & <i>Olea oleaster</i> DC (1/1) Streambed soil (0/1)	CIN (1) PLU (1) ^f PSC×KEL (1) ^g -

(Continues)

TABLE 1 (Continued)

Site no.	GPS coordinates	Altitude (m a.s.l.)	Location; parish, municipality, district	Geological substrate	Vegetation	Sampled tree species and health condition (no. of <i>Phytophthora</i> -positive/sampled trees) ^a	<i>Phytophthora</i> spp. (no. of positive samples) ^{b-e}
F33	40°29.619' N, 07°35.238' W	769	PNSE; UF S. Pedro e S. Julião, Gouveia, Guarda	Biotide granitoid	Submontane broadleaved forest	<i>C. sativa</i> DB (3/3) <i>C. sativa</i> DC & <i>Q. robur</i> DC (1/1) <i>Q. robur</i> DB (3/3) <i>Q. robur</i> × <i>pyrenaica</i> (0/1)	CIN (1), QUE (2) QUE (1) CIN (1), QUE (2) -
F34	40°33.005' N, 07°27.059' W	712	PNSE; Ribeira de Linhares; Linhares, Celorico da Beira, Guarda	Biotide granitoid	Submontane broadleaved forest and riparian gallery forest	<i>A. lusitanica</i> (1/1) <i>Q. pyrenaica</i> DC (4/6)	GON (1), C6PT1 (1) CIN (1), QUE (3) ^h
F35	40°24.209' N, 07°35.178' W	1428	PNSE; UF Aldeias e Mangualde da Serra, Gouveia, Guarda	Schists, greywackes, quartzites, conglomerates	Montane broadleaved forest	<i>B. celtiberica</i> (0/4) <i>Sorbus aucuparia</i> (0/1)	- -
F36	40°24.831' N, 07°32.663' W	1194	PNSE; Santa Maria, Manteigas, Guarda	Schists, greywackes, quartzites, conglomerates	Montane broadleaved forest	<i>C. sativa</i> DB (3/5)	CIN-PcG2 (3)
F37	40°24.738' N, 07°30.846' W	1083	PNSE; Santa Maria, Manteigas, Guarda	Schists, greywackes, quartzites, conglomerates	Montane broadleaved forest	<i>C. sativa</i> DB & bleeding bark cankers (2/4) <i>F. sylvatica</i> DB & bleeding bark canker (1/1) <i>F. sylvatica</i> DB (1/2) <i>Q. rubra</i> (0/1)	xCAM-DAPC9 (2) xCAM-DAPC9 (1) CIN (1), xCAM-DAPC9 (1) -
F38	40°25.303' N, 07°22.451' W	901	PNSE; Famalicão, Guarda, Guarda	Biotide granitoid	Lower montane broadleaved forest	<i>C. sativa</i> DB (2/2)	CIN-PcG2 (2)
F39	40°25.489' N, 07°22.523' W	892	PNSE; Famalicão, Guarda, Guarda	Biotide granitoid	Lower montane broadleaved forest	<i>C. sativa</i> DB & bleeding bark cankers (5/9)	CAS (4), CIN-PcG2 (3), PLU (1), xCAM-DAPC9 (1)
F40	40°28.519' N, 07°21.529' W	909	PNSE; Fernão Joanes, Guarda, Guarda	Biotide granitoid	Lower montane broadleaved forest	<i>C. sativa</i> DB (0/2)	-
F41	40°20.991' N, 07°05.656' W	745	Rio Côa; UF do Sabugal e Aldeia de Santo António, Sabugal, Guarda	Quaternary alluvial sediments	Riparian gallery forest	<i>A. lusitanica</i> bleeding bark canker (1/1)	xALN (1)
F42	40°18.596' N, 07°07.858' W	890	Reserva Natural da Serra da Malcata (RNSM); Meimão, Penamacor, Castelo Branco	Schists, greywackes, quartzites, conglomerates	Chestnut plantation	<i>C. sativa</i> DB (2/5)	CIN (2)



TABLE 1 (Continued)

Site no.	GPS coordinates	Altitude (m a.s.l.)	Location; parish, municipality, district	Geological substrate	Vegetation	Sampled tree species and health condition (no. of <i>Phytophthora</i> -positive/sampled trees) ^a	<i>Phytophthora</i> spp. (no. of positive samples) ^{b-e}
F43	40°15.846' N, 07°08.883' W	660	RNSM; Meimão, Penamacor, Castelo Branco	Schists, greywackes, quartzites, conglomerates	Submontane broadleaved forest	<i>Q. ilex</i> DC (2/2) <i>Q. ilex</i> (1/1)	QUE (2) QUE (1)
F44	40°15.166' N, 07°07.423' W	675	RNSM; Meimão, Penamacor, Castelo Branco	Schists, greywackes, quartzites, conglomerates	Submontane broadleaved forest	<i>A. unedo</i> (2/4) <i>Q. pyrenaica</i> DC (2/3)	QUE (2) QUE (2)
F45	40°11.999' N, 07°07.611' W	534	RNSM; Penamacor, Penamacor, Castelo Branco	Schists, greywackes, quartzites, conglomerates	Submontane cork oak plantation and mature forest	<i>Q. suber</i> DB & bleeding bark cankers (7/7)	CIN (6), QUE (1)
F46	40°16.669' N, 07°01.017' W	972	Reserva Natural da RNSM; Penamacor, Penamacor, Castelo Branco	Biotide granitoid	Lower montane broadleaved forest; in vicinity eucalypt plantations	<i>C. sativa</i> DB (3/5)	CIN (3)
F47	40°18.291' N, 07°01.836' W	824	RNSM; Quadrazais, Sabugal, Guarda	Biotide granitoid	Lower montane broadleaved forest	<i>Q. pyrenaica</i> DB (4/5)	QUE (3), xCAM (1)
F48	39°45.840' N, 08°59.792' W	71	Mata Nacional da Leiria (MNL); Marinha Grande, Marinha Grande, Leiria	Sand, gravels	Lowland mixed oak-pine forest and riparian gallery forest	<i>A. lusitanica</i> (0/1) <i>P. pinaster</i> (0/2) <i>Q. robur</i> (0/1) Streambed soil (0/1)	- - - -
F49	38°46.078' N, 09°25.405' W	263	Parque Natural Sintra-Cascais (PNSC); UF Santa Maria, São Miguel e São Pedro de Penaferrim, Sintra, Lisboa	Sienite	Colline, Atlantic mixed broadleaf-conifer forest	<i>Acacia dealbata</i> (1/1)	MUL (1), xPSC (1)
F50	38°46.965' N, 09°23.624' W	454	PNSC; UF S. Maria, S. Miguel e S. Pedro de Penaferrim, Sintra, Lisboa	Biotide granite	Colline, Atlantic mixed broadleaf-conifer forest	<i>A. alba</i> (0/1) <i>A. alba</i> DC & <i>C. sativa</i> DB (1/1) <i>C. sativa</i> DB (1/1) <i>F. sylvatica</i> (0/1) <i>Q. robur</i> (0/2)	- CIN (1) CIN (1) - -
F51	38°47.130' N, 09°23.574' W	425	PNSC; UF S. Maria, S. Miguel e S. Pedro de Penaferrim, Sintra, Lisboa	Biotide granite	Colline, Atlantic mixed broadleaf-conifer forest	<i>Q. robur</i> DB (2/3) <i>C. sativa</i> DB & <i>A. alba</i> DC (1/1) <i>C. sativa</i> DC & <i>F. sylvatica</i> DC (1/1)	GON (1), PLU (1) CIN-PcG1 (1) AMN (1), CHL (1), PLU (1)

(Continues)

TABLE 1 (Continued)

Site no.	GPS coordinates	Altitude (m.a.s.l.)	Location; parish, municipality, district	Geological substrate	Vegetation	Sampled tree species and health condition (no. of <i>Phytophthora</i> -positive/sampled trees) ^a	<i>Phytophthora</i> spp. (no. of positive samples) ^{b-e}
F52	38°47.063' N, 09°23.452' W	475	PNSC; UF S. Maria, S. Miguel e S. Pedro de Penaferrim, Sintra, Lisboa	Biotide granite	Colline, Atlantic mixed broadleaved forest	<i>A. pseudoplatanus</i> DC (2/2) <i>Laurus nobilis</i> (0/1) <i>Q. pyrenaica</i> DB (1/2) <i>Q. pyrenaica</i> (0/1)	CHL (2), MUL (2), PLU (1) - PLU (1) -
F53	38°47.429' N, 09°23.376' W	402	PNSC; UF S. Maria, S. Miguel e S. Pedro de Penaferrim, Sintra, Lisboa	Biotide granite	Colline, Atlantic mixed broadleaved forest	<i>A. pseudoplatanus</i> DC (2/2) <i>A. pseudoplatanus</i> DC & <i>F. angustifolia</i> DC (1/1) <i>A. pseudoplatanus</i> DC & <i>Q. robur</i> DC (0/1) <i>A. pseudoplatanus</i> DC & <i>Tilia cordata</i> DC (0/1) <i>Q. robur</i> DC (0/1)	MUL (1), PSE-DAPC5 (1) PLU (1) ^f - - -
F54	38°49.777' N, 08°20.978' W	125	Herdade das Antas; UF Cortiçadas de Lavre e Lavre; Montemor-o-Novo, Évora	Sandstones, sand, gravels	Lowland open oak woodland (Montado)	<i>Q. suber</i> DB (5/6)	CIN (5)
F55	38°47.782' N, 07°40.463' W	225	Herdade da Junceira; Évora Monte (Santa Maria), Estremoz, Évora	Biotide granitoid	Lowland open oak woodland (Montado)	<i>Q. ilex</i> DB (1/4) <i>Q. ilex</i> & <i>Q. suber</i> DB (2/2) <i>Q. suber</i> DB (1/1)	CIN (1) CIN (2) CIN (1)
F56	38°10.532' N, 08°39.951' W	200	Melides; Grandola, Setúbal	Slates, greywackes, flysch	Lowland open oak woodland (Montado)	<i>Q. suber</i> DB (2/5)	CIN-PcG2 (2)
F57	38°7.218' N, 08°33.260' W	130	UF Grândola e S. Margarida da Serra, Grândola, Setúbal	Slates, greywackes, flysch	Cork oak plantation	<i>Q. suber</i> DB (2/5) <i>Q. ilex</i> (0/1)	CIN-PcG2 (2) -
F58	38°10.935' N, 08°06.368' W	123	Alvito, Alvito, Beja	Shales, sandstones, conglomerates	Lowland open oak woodland (Montado)	<i>Q. ilex</i> DC (3/3) <i>Q. suber</i> DB (1/2)	QUE (2), PSY (2) CIN (1) ^f
F59	38°03.261' N, 08°17.110' W	75	Herdade do Outeiro; Figueira dos Cavaleiros, Ferreira do Alentejo, Beja	Shales, sandstones, conglomerates	Lowland open oak woodland (Montado)	<i>Q. suber</i> DC (2/2)	QUE (1), ULI (1)
F60	37°18.851' N, 08°32.455' W	496	Serra Algarvia (Barlavento); Monchique, Monchique, Faro	Sienite	Colline broadleaved forest	<i>C. sativa</i> (1/3) <i>C. sativa</i> DC (1/1) <i>C. sativa</i> DB (1/1)	CAS (1) CAS (1) CIN-PcG1 (1)
F61	37°23.342' N, 08°23.561' W	152	Serra Algarvia (Barlavento), Cerro do Malhão; S. Marcos da Serra, Silves, Faro	Slates, greywackes, flysch	Lowland open oak woodland (Montado)	<i>Q. suber</i> DB (1/1)	CIN (1)

TABLE 1 (Continued)

Site no.	GPS coordinates	Altitude (m a.s.l.)	Location; parish, municipality, district	Geological substrate	Vegetation	Sampled tree species and health condition (no. of <i>Phytophthora</i> -positive/sampled trees) ^a	<i>Phytophthora</i> spp. (no. of positive samples) ^{b-e}
F62	37°24.968' N, 08°24.721' W	210	Serra Algarvia (Barlavento), Foz do Grou; S. Marcos da Serra, Silves, Faro	Slates, greywackes, flysch	Lowland open oak woodland (Montado)	Q. suber DB (1/1)	CIN-PcG1 (1)
F63	37°15.088' N, 07°56.935' W	485	Serra Algarvia (Sotavento); Salir, Loulé, Faro	Slates, greywackes, flysch	Colline open oak woodland (Montado)	Q. suber DB (1/1)	CIN (1)
F64	37°13.737' N, 07°55.894' W	500	Serra Algarvia (Sotavento), Barranco do Velho; Salir, Loulé, Faro	Slates, greywackes, flysch	Colline open oak woodland (Montado)	Q. suber DB (4/6)	CIN-PcG2 (4), xPSC (1)
F65	37°12.709' N, 07°53.544' W	472	Serra Algarvia (Sotavento); S. Brás de Alportel, S.Brás de Alportel, Faro	Slates, greywackes, flysch	Colline open oak woodland (Montado)	Q. suber DB (1/1)	CIN (1)
F66	37°14.805' N, 07°51.500' W	423	Serra Algarvia (Sotavento); S. Brás de Alportel, S.Brás de Alportel, Faro	Slates, greywackes, flysch	Colline open oak woodland (Montado)	Q. suber DB (1/2)	CIN (1)
F67	37°09.548' N, 07°49.746' W	199	Santa Catarina da Fonte do Bispo, Tavira, Faro	Slates, greywackes, flysch	Arbutus plantation and open oak woodland (Montado)	A. unedo DB (1/1) Q. suber DB (1/1)	CIN (1) CIN (1), QUE (1)
F68	37°25.649' N, 07°31.695' W	223	Serra Algarvia (Sotavento); UF Alcoutim e Pereiro, Alcoutim, Faro	Slates, greywackes, flysch	Pine agroforestry plantation	Pinus pinea DC (1/1)	xPSC (1)
F69	37°51.300' N, 25°47.073' W	273	Lagoa das Sete Cidades; Sete Cidades, Ponta Delgada, São Miguel (Azores)	Basalt, trachyte	Colline broadleaved forest	Q. rubra DC (1/2)	MUL (1)
F70	32°48.569' N, 17°08.460' W	1161	Posto Forestal do Fanal; Ribeira da Janela, Porto Moniz, Funchal (Madeira)	Basalt	Atlantic montane laurisilva forest	Ocotea foetens & Persea indica (1/1)	CHL (1)

^aDB = crown dieback, DC = crown decline (increased transparency). No mentioning of disease symptoms denotes visually healthy trees.

^bAMN = *P. amnicola*, CAS = *P. castanetorum*, CHL = *P. chlamydospora*, CIN = *P. cinnamomi*, GON = *P. gonapodyides*, MUL = *P. multivora*, PLU = *P. plurivora*, PSE = *P. pseudosyringae*, PSY = *P. psychrophila*, QUE = *P. quercina*, SYR = *P. syringae*, ULI = *P. uliginosa*, C6PT1 = *P. taxon Clade06b_PT1*, PSC x KEL = *P. taxon pseudocryptogea x kelmanii*, xALN = *P. xalni*, xCAM = *P. xcambivora*, xPSC = *P. taxon xpseudocryptogea-like*.

^cDAPC groups within *P. xcambivora* according to the discriminant analysis of principal components (DAPC) of the population genomic study of Mullett et al. (2023). DAPC9 isolates belong to the A2 mating type.

^d*Phytophthora cinnamomi* panglobal clonal lineages PcG1 and PcG2, according to the population genomic study of Shakya et al. (2021). All isolates of PcG1 and PcG2 belong to the A2 mating type.

^eDAPC groups within *P. pseudosyringae* according to the DAPC of the population genomic study of Mullett et al. (2024).

^f*Elongisporangium* taxon anandrum-like also isolated.

^g*Phytophthium citrinum* also isolated.

^h*Phytophthium citrinum* also isolated.

ⁱ*Phytophthium* taxon UZ612 also isolated.

^j*Elongisporangium* sp. also isolated.

TABLE 2 Location and altitude of the 39 riparian sites sampled at 34 rivers and streams in Portugal and *Phytophthora* and other oomycete taxa isolated.

Site no.	GPS coordinates	Altitude (m.a.s.l.)	Location; parish, municipality, district	Location of catchment and vegetation	<i>Phytophthora</i> and <i>Nothophytophthora</i> spp. ^{a-d}
R01	41°44.266' N, 08°09.304' W	497	Ribeira da Quelha Verde; Vilar da Veiga, Terras de Bouro, Braga	Parque Nacional da Peneda-Gerês (PNPG); sub-/montane broadleaved forests, heath- & grasslands	CHL, GON, PLU, PSE-DAPC5
R02	41°44.637' N, 08°09.239' W	542	Ribeiro da Figueira; Vilar da Veiga, Terras de Bouro, Braga	PNPG; sub-/montane broadleaved forests, heath- & grasslands	-
R03	41°44.908' N, 08°09.221' W	597	Forest stream, Secelo; Vilar da Veiga, Terras de Bouro, Braga	PNPG; sub-/montane broadleaved forests, heath- & grasslands	GON, PLU, xCAM, xPSC
R04	41°45.421' N, 08°08.841' W	779	Ribeira da Cantina; Vilar da Veiga, Terras de Bouro, Braga	PNPG; sub-/montane broadleaved forests, heath- & grasslands	GON
R05	41°47.559' N, 08°08.285' W	673	Forest stream, Lagos (Albergaria); Campo do Gerês, Terras de Bouro, Braga	PNPG; sub-/montane broadleaved forests, heath- & grasslands	GON
R06	41°48.219' N, 08°07.925' W	725	Forest stream; Curral de S. Miguel; Campo do Gerês, Terras de Bouro, Braga	PNPG; sub-/montane broadleaved forests, heath- & grasslands	GON, xCAM-DAPC5, xCAM-DACP9
R07	41°42.516' N, 08°07.028' W	361	Rio Arado, (Ponte das Cervas); Vilar da Veiga, Terras de Bouro, Braga	PNPG; sub-/montane broadleaved forests, heath- & grasslands; horticulture	GON, xCAM-DAPC9
R08	41°16.747' N, 08°04.090' W	75	Rio Tâmega, Lufrei, Amarante, Porto	Lowland to submontane broadleaved forests; horticulture	LAC, xC9PT1, xP16855
R09	40°22.690' N, 08°21.992' W	342	Forest stream, Vale dos Fetos (Buçaco); Luso, Mealhada, Aveiro	Mata Nacional do Buçaco (MNB); colline broadleaved forest and a castle park	PLU
R10	40°22.684' N, 08°22.325' W	283	Forest stream, Fonte Fria (Buçaco); Luso, Mealhada, Aveiro	MNB; colline broadleaved forest and a castle park	CHL, PCIP, PLU, P16855 ^e
R11	40°22.622' N, 08°21.774' W	373	Forest stream, Vale dos Abetos (Buçaco); Luso, Mealhada, Aveiro	MNB; colline broadleaved forest and a castle park	CHL, PLU, xPSC
R12	40°23.896' N, 07°30.625' W	682	Rio Zêzere; Santa Maria, Manteigas, Guarda	Parque Natural da Serra da Estrela (PNSE); sub-/montane broadleaved forests, heath- & grasslands; horticulture	CHL, GON, LAC, PLU, PSC x KEL, xCAM-DAPC1
R13	40°24.476' N, 07°28.570' W	595	Rio Zêzere; Sameiro, Manteigas, Guarda	PNSE; sub-/montane broadleaved forests, heath- & grasslands; horticulture	CHL, GON, LAC, PLU ^f
R14	40°23.293' N, 07°26.841' W	545	Rio Beijames; Vale de Amoreira, Manteigas, Guarda	PNSE; sub-/montane broadleaved forests, heath- & grasslands; horticulture	AMN, CHL, LAC, PLU, PSC x KEL, xSTA ^g
R15	40°29.161' N, 07°24.953' W	816	Rio Mondego; União de Freguesias (UF) Corujeira e Trinta, Guarda, Guarda	PNSE; montane broadleaved forests, heath- & grasslands; horticulture	BIL, GON, LAC, xCAM-DAPC5
R16	40°30.478' N, 07°22.686' W	763	Rio Mondego; Videmonte, Guarda, Guarda	PNSE; sub-/montane broadleaved forests, heath- & grasslands; horticulture	PLU, xCAM-DAPC ^{h,i}
R17	40°18.802' N, 06°59.097' W	818	Rio Côa; Quadrazais, Sabugal, Guarda	Reserva Natural da Serra da Malcata (RNSM); montane broadleaved forests, heath- & grasslands; horticulture	CHL, GON, LAC, xHEN, xLAC
R18	40°18.780' N, 06°59.390' W	817	Rio Côa; Quadrazais, Sabugal, Guarda	RNSM; montane broadleaved forests, heath- & grasslands; horticulture	LAC, PSC, xHEN, xLAC
R19	40°18.085' N, 07°02.128' W	796	Ribeira da Barroca da Amieira; Quadrazais, Sabugal, Guarda	RNSM; montane broadleaved forests, heath- & grasslands; horticulture	GON
R20	40°15.378' N, 07°05.031' W	604	Ribeira da Meimosa; Penamacor, Penamacor, Castelo Branco	RNSM; sub-/montane broadleaved forests, heath- & grasslands	PLU, PSC, xPSC ^e
R21	40°15.602' N, 07°05.597' W	590	Ribeira da Meimosa; Meimão, Penamacor, Castelo Branco	RNSM; sub-/montane broadleaved forests, heath- & grasslands	BIL, LAC, PSC, THE x AMN, x AMN, x WS
R22	40°11.578' N, 07°05.549' W	442	Ribeiro da Mouca; Penamacor, Penamacor, Castelo Branco	RNSM; sub-/montane broadleaved forests, heath- & grasslands	BIL, KEL, LAC, PSC x KEL
R23	40°12.206' N, 07°06.483' W	530	Ribeira do Salgueirinho; Penamacor, Penamacor, Castelo Branco	RNSM; sub-/montane broadleaved forests, heath- & grasslands	PLU, PSC, PSC x KEL
R24	39°46.436' N, 09°00.880' W	37	Forest stream; Marinha Grande, Marinha Grande, Leiria	Mata Nacional de Leiria; Mediterranean mixed pine-oak forests and Machia vegetation	AMN, GON, MUL, AMN x CHL, THE x AMN

TABLE 2 (Continued)

Site no.	GPS coordinates	Altitude (m.a.s.l.)	Location; parish, municipality, district	Location of catchment and vegetation	Phytophthora and Notohytophthora spp. ^{a-d}
R25	38°46.843' N, 09°26.686' W	360	Fonte das 2 Bicas; Colares, Sintra, Lisboa	Parque Natural Sintra-Cascais (PNSC); Atlantic colline broadleaved and pine forests	PSE-DAPC3, THE, xAMN
R26	38°45.508' N, 09°25.179' W	113	Forest stream, Quinta do Pisão; Alcabideche, Cascais, Lisboa	PNSC; Atlantic lowland to colline broadleaved and pine forests; horticulture	AMN, AMN×CHL, MUL, THE×AMN
R27	38°46.078' N, 09°25.405' W	166	Tributary of Barragem do Rio da Mula; UF Santa Maria, São Miguel e São Pedro de Penaferrim, Sintra, Lisboa	PNSC; Atlantic colline to colline broadleaved and pine forests	AMN
R28	38°48.109' N, 09°26.763' W	18	Ribeira de Colares; Colares, Sintra, Lisboa	PNSC; Atlantic lowland to colline broadleaved and pine forests; horticulture	AMN, BIL, CHL, LAC, PCIP
R29	38°47.466' N, 09°25.479' W	193	Forest stream, Monserrate; UF S. Maria, S. Miguel e S. Pedro de Penaferrim, Sintra, Lisboa	PNSC; Atlantic colline broadleaved and pine forests & heathlands	AMN, AMP, CHL
R30	38°47.126' N, 09°23.538' W	427	Small stream, Vale dos Fetos, Pena; UF S. Maria, S. Miguel e S. Pedro de Penaferrim, Sintra, Lisboa	PNSC; Atlantic colline broadleaved and pine forest; Botanical Garden with exotic plants	AMN, AMN×CHL GON, RAM ^f
R31	37°15.530' N, 08°32.607' W	75	Ribeira de Vale do Boi; Monchique, Monchique, Faro	Mediterranean, lowland to colline broadleaved and pine forests; heathlands; acacias and eucalypts; horticulture	AMN, GON, LAC, THE, THE×AMN, THE×GON, xP16855
R32	37°13.114' N, 08°30.464' W	25	Ribeira de Odelouca; Silves, Silves, Faro	Mediterranean, lowland to colline broadleaved and pine forests; heathlands; acacias and eucalypts; horticulture	xP16855, HON, xHEN1, ZEN×PER
R33	37°12.607' N, 08°00.660' W	137	Ribeira da Menalva (Fonte Benémola); UF Querença, Tôr e Benafim, Loulé, Faro	Mediterranean, lowland to colline broadleaved and pine forests & heathlands	LAC, xP16855, ZEN×PER
R34	37°09.339' N, 07°41.966' W	33	Ribeira da Assêca, Pêgo do Inferno; UF Luz de Tavira e Santo Estevão, Tavira, Faro	Mediterranean, lowland broadleaved forests & heathlands; horticulture	HON
R35	37°09.214' N, 07°40.343' W	10	Rio Séqua; UF Santa Maria e Santiago, Tavira, Faro	Mediterranean, lowland to colline oak and pine forests; heathlands; horticulture incl. <i>Citrus</i> plantations	CIP, HON, INU, LAC, PCIP, SYL, xLUS, ZEN×PER
R36	37°28.390' N, 07°28.592' W	25	Ribeira de Cadavais; UF Alcoutim e Pereiro, Alcoutim, Faro	Mediterranean, lowland broadleaved and pine woodlands; heathlands; horticulture	HON
R37	37°20.384' N, 07°28.933' W	25	Ribeira de Odeleite; Odeleite, Castro Marim, Faro	Mediterranean, lowland broadleaved and pine woodlands; heathlands; horticulture	LAC
R38	37°24.666' N, 07°27.810' W	17	Rio Guadiana; UF Alcoutim e Pereiro, Alcoutim, Faro	Huge catchment; Mediterranean, lowland to submontane broadleaved and pine forests; heathlands; horticulture	HON, THE
R39	37°22.723' N, 07°26.423' W	5	Rio Guadiana, Álamo; UF Alcoutim e Pereiro, Alcoutim, Faro	Huge catchment; Mediterranean, lowland to submontane broadleaved and pine forests; heathlands; horticulture	HON

^aNon-hybrid taxa: AMN = *P. amnicola*, AMP = *N. amphigynosa*, BIL = *P. bilobang*, CHL = *P. chlamydospora*, CIP = *P. citrophthora*, GON = *P. gonapodyides*, HON = *P. honggalleglyana*, INU = *P. inundata*, KEL = *P. kelmanii*, LAC = *P. lacustris*, MUL = *P. multivora*, PLU = *P. plurivora*, PCIP = *P. pseudocitrophthora*, PSC = *P. pseudocryptogea*, PSE = *P. pseudosyringae*, RAM = *P. ramorum*, SYL = *P. taxon sylvatica*-like 1, P16855 = *P. taxon P16855*, THE = *P. thermophila*.

^bHybrid taxa: AMN×CHL = *P. taxon amnicola* × *chlamydospora*, PSC×KEL = *P. taxon pseudocryptogea* × *kelmanii*, THE×AMN = *P. taxon thermophila* × *amnicola*, THE×GON = *P. taxon thermophila* × *gonapodyides*, xAMN = *P. taxon amnicola*-like, xCAM = *P. xambivora*, xC9PT1 = *P. taxon xClade09a1_PT1*, xHEN = *P. taxon xHennops*, xHEN1 = *P. taxon xHennops*-like 1, xLAC = *P. taxon xlacustris*-like, xLUS = *P. xlusitanica*, xPSC = *P. taxon xpseudocryptogea*-like, xP16855 = *P. taxon xP16855*-like, xSTA = *P. xstagnum*, xWS = *P. taxon xWS*, ZEN×PER = *P. taxon zentmyerii* × *Peru4*-like.

^cDAPC groups within *P. pseudosyringae* according to the discriminant analysis of principal components (DAPC) of the population genomic study of Mullett et al. (2024).

^dDAPC groups within *P. xambivora* according to the DAPC of the population genomic study of Mullett et al. (2023). DAPC1 isolates belong to the A1 mating type, DAPC5 and DAPC9 isolates to the A2 mating type.

^e*Phytophthora litorale* also isolated.

^f*Phytophthora* taxon PV So7 also isolated.

^g*Pythium* taxon strain 1-9 also isolated.

^h*Phytophthora* taxon PB-2013 also isolated.

ⁱ*Phytophthora* taxon KC-2014 also isolated.

^jThe isolate of *P. ramorum* belongs to the EU1 lineage according to the genomic study of Van Poucke et al. (2021).



FIGURE 2 Representative forest stands and rivers sampled in Portugal; (a) submontane mixed forest F01 in the Parque Nacional da Peneda-Gerês (PNPG); (b) submontane *Quercus suber* forest F10 in the PNPG; (c) colline to submontane, mixed forest F25 in the Mata Nacional do Buçaco; (d) submontane *Quercus pyrenaica* forest F34 in the Parque Nacional da Serra da Estrela (PNSE); (e) montane forest (F37) of *Castanea sativa*, *Fagus sylvatica* and *Quercus rubra* in the PNSE; (f) colline mixed forest F50 in the Parque Natural Sintra-Cascais; (g) open lowland woodland (Montados) F54 of *Quercus ilex* and *Q. suber* in the Alentejo region; (h) baiting raft (arrow) floating in the Rio Arado (R07) in the PNPG. For GPS coordinates see [Tables 1](#) and [2](#); for location of sites see [Figure 1](#).

cultures were maintained on both V8A and carrot agar (CA; Pérez-Sierra et al., 2022) at 10°C in the dark.

2.2 | Molecular identification of isolates

For all *Phytophthora* isolates obtained in this study, mycelial DNA was extracted from pure 1-week-old V8A cultures. Total DNA was extracted using the DNeasy Plant Mini kit (QIAGEN) or the Phire Plant Direct PCR Kit (Thermo Fisher Scientific) following the manufacturer's instructions. DNA was stored at -20°C until further use. For all isolates, the region spanning the internal transcribed spacer (ITS1-5.8S-ITS2) region of the ribosomal DNA was amplified using

primer pairs ITS1/ITS4 or ITS6/ITS4 (Cooke et al., 2000; White et al., 1990). For all isolates from new taxa and a representative set of isolates from known species of *Phytophthora*, the mitochondrial *cox1* gene was amplified with both primer pairs OomCox1-Levup/OomCox1-Levlo and COXF4N/COXR4N (Kroon et al., 2004; Robideau et al., 2011) or FM84/FM83 (Martin & Tooley, 2003). The PCR mixture and the amplification conditions for ITS and *cox1* were as described by Cooke et al. (2000), Kroon et al. (2004), Robideau et al. (2011) and Martin and Tooley (2003). PCR consumables were purchased from Thermo Fisher Scientific. PCR was carried out in an Eppendorf Mastercycler Ep Gradient thermocycler. PCR products were purified and sequenced by MacroGen Europe (Amsterdam, Netherlands) in both directions using the amplification primers.

Electropherograms were quality-checked and forward and reverse reads were compiled using Geneious Prime v. 2024.0.2 (Biomatters Ltd). Pronounced double peaks were considered as heterozygous positions and labelled according to the International Union of Pure and Applied Chemistry (IUPAC; <https://iupac.org>) coding system. The consensus sequences were subjected to an NCBI BLAST search (<http://www.ncbi.nlm.nih.gov/BLAST/>) and to a BLAST search in a local database containing sequences of ex-type isolates or key isolates from published studies to identify the closest related species. Isolates were assigned to a species when sequence identities were above a 99% cut-off in respect to those of ex-type isolates or key isolates. ITS sequences from representative isolates of all oomycete species and *cox1* sequences of representative isolates of several known species and putative new species obtained in this study were deposited at GenBank.

2.3 | Classical identification of isolates

Colony growth patterns of 7-day-old cultures grown at 20°C in the dark on V8A, CA and potato dextrose agar (PDA; Jung et al., 2011) and morphological features of sporangia, oogonia, antheridia, chlamydospores, hyphal swellings and aggregations were compared with known isolates and with species descriptions in the literature.

Sporangia were produced in non-sterile soil filtrate (Jung, 2009). Microscopic examinations and measurements of morphological structures at 200× and 400× were according to Jung, Horta Jung, Cacciola, et al. (2017) and Jung et al. (2022) using a compound microscope (Zeiss Imager.Z2), a digital camera (Zeiss Axiocam ICc3) and biometric software (Zeiss ZEN). Self-sterile isolates were paired on V8A and CA with known A1 and A2 mating type tester strains of *P. cinnamomi* (A1: TW12; A2: MP74; Jung, Horta Jung, Scanu, et al., 2017) and *P. xambivora* (A1: Resi75; A2: SK9; Jung, Horta Jung, Scanu, et al., 2017) (isolates with non-papillate sporangia) or *P. botryosa* (A2: MYA-4059; Jung et al., 2024), *P. colocasiae* (A1: MYA-4159; A2: 35D3; Yang et al., 2017; Jung et al., 2024) and *P. meadii* (A1: MYA-4042; A2: MYA-4043; Jung et al., 2024) (isolates with papillate sporangia), and examined after 4–6 weeks incubation at 20°C in order to determine whether they were heterothallic or sterile (Jung et al., 2022, 2024). All isolates are preserved in the culture collections maintained at the University of Algarve and Mendel University in Brno.

3 | RESULTS

Overall, 799 isolates of 48 *Phytophthora* taxa were obtained from forest stands (Table 1), river systems (Table 2) and forest nurseries (Table 3) in Portugal. Although isolates from other oomycete genera were discarded regularly during the isolation process, 25 isolates of *Elongisporangium*, *Nothophytophthora*, *Phytophthium* and *Pythium* spp. were also taken into culture. GenBank accession numbers of ITS and *cox1* sequences of representative isolates of all oomycete taxa are

given in Table S1. Official species descriptions of new *Phytophthora* and other oomycete taxa including detailed descriptions of morphological characteristics, morphometric data, temperature-growth data and multigenic phylogenies have been published recently (Jung et al., 2024; Jung, Horta Jung, Cacciola, et al., 2017; Jung, Scanu, Bakonyi, et al., 2017) and more will be presented in future publications. Comprehensive lists for associating sample sources (tree species, river/stream) with oomycete taxa, and oomycete taxa with sample sources and site categories are given in Tables S2–S4.

3.1 | *Phytophthora* diversity and disease symptoms in natural and semi-natural forest stands of Portugal

Most tree species in the *Phytophthora*-infested forest stands showed a range of symptoms typical of *Phytophthora* infections, such as decline (increased transparency) and dieback of crowns and losses of fine and lateral roots. Moreover, in several stands *A. lusitanica*, *C. sativa*, *F. sylvatica*, *Q. rubra* and *Q. suber* also suffered from bleeding bark cankers (Figures 3 and 4; Table 1). Declines and diebacks of tree species reported for the first time in Europe and Portugal, respectively, are listed in Table S5.

In 62 of the 70 forest stands (88.6%), 17 *Phytophthora* taxa were isolated from 148 of the 302 rhizosphere soil samples (49%) of 25 of the 29 tree species sampled (86.2%) and from all seven bark cankers of *A. lusitanica* and *F. sylvatica* sampled. The 254 *Phytophthora* isolates belonged to 13 known species from six phylogenetic clades, namely *P. multivora* and *P. plurivora* from Clade 2; *P. pseudosyringae* and *P. psychrophila* from Clade 3; *P. amnicola*, *P. chlamydospora* and *P. gonapodyides* from Clade 6; *P. cinnamomi*, *P. uliginosa*, *P. xalni* and *P. xambivora* from Clade 7; and *P. syringae* and *P. quercina* from Clades 8 and 12, respectively; one new Clade 12 species, which due to its specificity to chestnut was named *P. castanetorum* (Jung, Horta Jung, Cacciola, et al., 2017); and three undescribed taxa, one new non-hybrid taxon provisionally named *P. taxon Clade06b_PT* and two Clade 8 hybrid taxa, the designated *P. taxon pseudocryptogea* × *kelmanii* and the new provisionally named *P. taxon xpseudocryptogea-like* (Figure 5; Table 1). In addition, *Phytophthium citrinum*, the undescribed *Phytophthium* taxa REB326-70 and UZ612, and the undescribed *Elongisporangium* taxon anandrum-like were recovered from five soil samples (Table 1). With 39 forest sites (55.7%), an altitudinal range between 123 and 1194 m a.s.l. and 12 tree species (41.4%), *P. cinnamomi* was by far the most common species (Figure 5; Table 1). All 82 isolates belonged to the A2 mating type. A subgroup of these isolates was included in the population genomic studies of Shakya et al. (2021) that revealed the presence of both globally distributed clonal A2 mating type lineages, PcG1-A2 and PcG2-A2. *P. quercina* was isolated in 15 forest stands from the rhizosphere of four oak species, *Q. ilex*, *Q. pyrenaica*, *Q. robur* and *Q. suber*, and four tree species belonging to other genera, namely *A. unedo*, *C. sativa*, *Ilex aquifolium* and *Phyllirea latifolia* (Figure 5; Table 1). With 15 forest sites (21.4%) and eight tree species (38.1%), namely *Acer pseudoplatanus*, *Betula*




FIGURE 3 Disease symptoms of mature trees in natural and semi-natural forest stands in Portugal caused by *Phytophthora* pathogens; (a) crown dieback and mortality of *Castanea sativa* in the montane forest F37 in the Parque Nacional da Serra da Estrela (PNSE) (*P. cinnamomi* and *P. ×cambivora*); (b) severe dieback and mortality of *Quercus pyrenaica* in the submontane forest F19 in the Serra da Nogueira (*P. quercina*); (c) severe dieback of *Quercus robur* in submontane forest F06 in the Parque Nacional da Peneda-Gerês (PNPG) (*P. quercina*); (d) high mortality of *Alnus lusitanica* (arrows) along the Rio Alva (*P. xalni*); (e) bleeding bark canker on *A. lusitanica* at the Rio Côa (*P. xalni*); (f) bleeding bark canker on *Quercus rubra* in the submontane forest F03 in the PNPG (*P. cinnamomi*); (g) bleeding bark canker on *Fagus sylvatica* in the montane forest F37 in the PNSE (*P. ×cambivora*).

celtiberica, *C. sativa*, *F. sylvatica*, *Pinus pinaster*, *Q. ilex*, *Q. pyrenaica* and *Salix caprea*, *P. ×cambivora* was the third most frequent species (Figure 5; Table 1). This pathogen was also isolated from the bleeding collar rot lesion of a mature *F. sylvatica* tree at the montane stand F37 in the Parque Nacional da Serra da Estrela (Figure 3; Table 1). All 37 *P. ×cambivora* isolates from the forest sites were included in the population genomic study of Mullett et al. (2023) and belonged to the discriminant analysis of principal components (DAPC) group 9. Mating tests revealed all isolates were of the A2 mating type. *P. plurivora* and *P. multivora* were obtained from five (7.1%) and four (5.7%) forest stands, respectively (Figure 5; Table 1). *C. sativa* hosted the highest *Phytophthora* diversity in the rhizosphere, with seven species (*P. amnicola*, *P. castanetorum*, *P. chlamydospora*, *P. cinnamomi*, *P. plurivora*, *P. quercina* and *P. ×cambivora*, *C. sativa*); followed with five species by *Q. pyrenaica* (*P. cinnamomi*, *P. plurivora*, *P. pseudosyringae*, *P. quercina* and *P. ×cambivora*) and *Acer pseudoplatanus* (*P. chlamydospora*, *P. multivora*, *P. plurivora*, *P. pseudosyringae* and *P. ×cambivora*); and with four species by *Q. robur* and *Q. suber* (Table 1, Table S2). The *P. pseudosyringae* isolates from the forest sites included in the population genomic study of Mullett et al. (2024) belonged to two different DAPC groups, 1 and 5.

Regardless of the tree species, there was a considerable difference in the isolation rates from rhizosphere soil between visually healthy trees (nine of the 89 trees infested = 10.1%) and declining trees (139 of the 213 trees infested = 65.3%). Decline and dieback of *C. sativa* was associated with *P. cinnamomi* and *P. ×cambivora* in 12 and five stands, respectively (Table 1). *P. cinnamomi* was also associated with the decline and dieback of *Q. suber* in 16 stands (Table 1). Decline and dieback of *Q. pyrenaica* were associated with *P. ×cambivora* in four stands, with *P. cinnamomi* and *P. quercina* in each of three stands and with *P. plurivora* in one stand (Table 1). With *Q. robur*, *P. cinnamomi* and *P. quercina* were recovered from declining trees in four and two stands, respectively, and *P. plurivora* in one stand. In contrast, healthy *Q. robur* trees in six stands did not yield any *Phytophthora* species (Table 1). Decline and dieback of *Q. ilex* was found associated with *P. quercina* in two stands and with *P. cinnamomi* and *P. ×cambivora* in one stand each (Table 1). The bark cankers of riparian *A. lusitanica* trees at the Rio Alva, Rio Côa, Rio Tâmega and Rio Fervença (Figure 3) were caused by *P. xalni* (Table 1). At three other riparian sites, *P. cinnamomi* (F19 along a forest stream, F32 along the Rio Lima) and *P. gonapodyides* (F34 along the Ribeira de Linhares) were isolated from the rhizosphere

TABLE 3 Forest nursery stock sampled in Portugal, health condition and *Phytophthora* and other oomycete taxa isolated.

Nursery no.	Sampled tree species (no. of <i>Phytophthora</i> -positive/sampled trees)	Symptoms of sampled plants ^a	<i>Phytophthora</i> spp. (no. of positive samples) ^{b-d}
N01	Soil of transplanting bed (0/2)	-	- ^e
N02	<i>Abies alba</i> (1/1)	Stunted, chlorotic, small needles	CIN-PcG2 (1)
	<i>Arbutus unedo</i> (1/1)	Wilting	CIN-PcG1 (1)
	<i>Chamaecyparis lawsoniana</i> (0/1)	Healthy	-
	<i>Cupressus sempervirens</i> (1/1)	Some stunted & chlorotic or dead	CIN (1)
	<i>Pinus pinea</i> (1/2)	Some stunted & chlorotic or dead	×PSC (1) ^f
	<i>Pinus radiata</i> (0/1)	Some stunted & chlorotic or dead	- ^{g,h}
	<i>Pinus sylvestris</i> (1/1)	Some stunted & chlorotic or dead	PSC (1), ×PSC (1)
	<i>Quercus pyrenaica</i> (1/2)	Some wilting or dead	×CAM DAPC11 (1) ^g
	<i>Quercus robur</i> (1/1)	Some wilting or dead	×PSC (1) ⁱ
N03	<i>Quercus suber</i> (0/2)	Healthy	-
N03	<i>Castanea sativa</i> (1/1)	Some stunted & chlorotic or dead	CIN-PcG2 (1)
	<i>Eucalyptus nitens</i> (1/1)	Some stunted & chlorotic or dead	ALT (1), CIN (1)
	<i>P. pinea</i> (0/1)	Healthy	-
	<i>Quercus ilex</i> (1/1)	Some stunted & dead	CIN (1), QUT (1)
	<i>Q. pyrenaica</i> (0/1)	Healthy	-
	<i>Q. robur</i> (0/1)	Healthy	-
	<i>Q. suber</i> (0/1)	Healthy	-
	<i>Quercus rubra</i> (1/1)	Healthy	QUT (1)
N04	<i>C. sativa</i> (1/1)	Some stunted & chlorotic or dead	CIN-PcG2 (1)
	<i>P. pinea</i> (0/1)	Healthy	-
	<i>Q. pyrenaica</i> (0/1)	Healthy	-
	<i>Q. robur</i> (0/1)	Healthy	-
	<i>Q. rubra</i> (0/1)	Healthy	-
	<i>Q. suber</i> (1/1)	Some stunted & chlorotic	PLU (1), ×CAM DAPC11 (1) ^j
N05	<i>C. sativa</i> (1/1)	Some stunted & chlorotic or dead	CIN-PcG2 (1)
	<i>C. sativa</i> (1/1)	Chlorotic & collar rot, direct isolation	CIN (1)
	<i>P. pinea</i> (0/1)	Some stunted & chlorotic	- ^k
	<i>Pinus pinaster</i> (0/1)	Healthy	- ^h
	<i>Quercus coccinea</i> (0/1)	Healthy	-
	<i>Q. pyrenaica</i> (1/1)	Some stunted & chlorotic	QUT (1)
	<i>Q. rubra</i> (1/1)	Some stunted & chlorotic	QUT (1)
	<i>Q. suber</i> (0/1)	Healthy	-
N06	<i>C. sativa</i> (0/1)	Healthy	- ^g
	<i>P. pinea</i> (1/1)	Some stunted & chlorotic	PSC (1)
	<i>Quercus faginea</i> (1/1)	Healthy	QUT (1)
	<i>Q. ilex</i> (0/1)	Healthy	-
	<i>Q. pyrenaica</i> (2/2)	Some stunted & chlorotic	×CAM DAPC9 (1), QUT (1)
	<i>Q. robur</i> (1/1)	Healthy	×CAM DAPC9 (1) ^j
	<i>Q. rubra</i> (2/2)	Some stunted & chlorotic or dead	CIN (1), QUT (1) ^j
	<i>Q. suber</i> (0/2)	Healthy	-
N07	<i>P. pinea</i> (1/1)	Stunted & chlorotic or dead	PAR (1), PSC (1)
	<i>P. pinaster</i> (1/1)	Some stunted & chlorotic or dead	CIN (1)
	<i>Q. faginea</i> (1/1)	Some stunted & chlorotic	MUL (1), PSC (1)
	<i>Q. ilex</i> (1/1)	Stunted & chlorotic or dead	CIN (1)
	<i>Q. suber</i> (1/1)	Stunted & chlorotic or dead	CIN (1)

(Continues)

TABLE 3 (Continued)

Nursery no.	Sampled tree species (no. of <i>Phytophthora</i> -positive/sampled trees)	Symptoms of sampled plants ^a	<i>Phytophthora</i> spp. (no. of positive samples) ^{b-d}
N08	<i>P. pinea</i> (1/1)	Some stunted & chlorotic	PSC×KEL
	<i>Q. ilex</i> (1/1)	Dead	CAC (1), ×CAM DAPC4 (1)
	<i>Q. ilex</i> (1/1)	Healthy	×CAM DAPC4 (1)
	<i>Q. suber</i> (1/1)	Some stunted & chlorotic or dead	CIN-PcG2 (1), MUL (1)
N09	<i>P. pinea</i> (1/1)	Some stunted & chlorotic	CAC (1)
N10	<i>Q. ilex</i> (1/1)	Dead	CIN (1)
	<i>Q. ilex</i> (1/1)	Healthy	×CAM DAPC1 (1)
N11	<i>Alnus lusitanica</i> (1/1)	Healthy	CRA (1)
	<i>A. unedo</i> (1/1)	Some chlorotic or wilting	×CRA (1)
	<i>C. sativa</i> (2/2)	Chlorotic or dead, collar rot	CIN (1), CRA (1), ×CAM (1)
	<i>Cupressus lusitanica</i> (0/1)	Some stunted & chlorotic or dead	-
	<i>Q. faginea</i> (1/1)	Some stunted & chlorotic or dead	PLU (1)
	<i>Q. ilex</i> (1/1)	Stunted & chlorotic or dead	QUE (1), PSC (1)
	<i>Q. pyrenaica</i> (1/1)	Some stunted & chlorotic or dead	PSC (1)
	<i>Q. robur</i> (1/1)	Some stunted & chlorotic or dead	CRA (1), QUE (1) ^g
	<i>Q. suber</i> (1/1)	Stunted & chlorotic or dead	CRA (1), PSC (1)
	<i>Taxus baccata</i> (1/1)	Some stunted & chlorotic or dead	CRA (1)
N12	<i>A. lusitanica</i> (0/1)	Healthy	-
	<i>A. unedo</i> (0/1)	Mostly healthy, some wilting	-
	<i>C. sativa</i> (0/1)	Healthy	-
	<i>Juglans regia</i> (0/1)	Healthy	-
	<i>P. pinea</i> (0/1)	Healthy	-
	<i>P. pinea</i> (1/1)	Some stunted & chlorotic or dead	CAC (1)
	<i>Q. faginea</i> (1/1)	Some stunted & chlorotic or dead	PSC (1), QUE (1)
	<i>Q. ilex</i> (1/1)	Healthy	PSY (1)
	<i>Q. pyrenaica</i> (1/1)	Some stunted & chlorotic	QUE (1)
	<i>Q. robur</i> (1/1)	Healthy	GON (1)
	<i>Q. suber</i> (0/1)	Healthy	-
N13	<i>A. lusitanica</i> (1/1)	Healthy	PSY (1), QUE (1)
	<i>A. unedo</i> (0/1)	Healthy	-
	<i>P. pinaster</i> (1/1)	Some stunted & chlorotic or dead	MUL (1), PSC (1)
	<i>P. pinea</i> (0/1)	Healthy	-
	<i>Q. faginea</i> (0/1)	Healthy	-
	<i>Q. ilex</i> (0/1)	Mostly healthy, some stunted & chlorotic	-
	<i>Q. suber</i> (0/1)	Mostly healthy, some stunted & chlorotic	-

^aAlmost all plants with above-ground symptoms had extensive fine root losses and/or root rot.

^bALT=*P. alticola*, CAC=*P. cactorum*, CIN=*P. cinnamomi*, CRA=*P. crassamura*, GON=*P. gonapodyides*, MUL=*P. multivora*, PAR=*P. parvispora*, PLU=*P. plurivora*, PSC=*P. pseudocryptogea*, PSY=*P. psychrophila*, QUE=*P. quercina*, QUT=*P. quercetorum*, PSC×KEL=*P. taxon pseudocryptogea* × *kelmanii*, ×CAM=*P. xcambivora*, ×CRA=*P. taxon xcrassamura*-like, ×PSC=*P. taxon xpseudocryptogea*-like.

^c*Phytophthora cinnamomi* panglobal clonal lineages PcG1 and PcG2, according to the population genomic study of Shakya et al. (2021). All isolates PcG1 and PcG2 belong to the A2 mating type.

^dDAPC groups within *P. xcambivora* according to the discriminant analysis of principal components (DAPC) of the population genomic study of Mullett et al. (2023). DAPC4 and DAPC11 isolates belong to the A1 mating type, DAPC9 isolates to the A2 mating type.

^e*Phytophthora vexans* isolated.

^f*Pythium diclinum* also isolated.

^g*Phytophthora mercuriale* also isolated.

^h*Phytophthora* taxon *mercuriale*-like also isolated.


ⁱ*Phytophthora* taxon IRAN4019C also isolated.

^j*Phytophthora citrinum* also isolated.

^k*Elongisporangium undulatum* also isolated.

^l*Phytophthora litorale* also isolated.

FIGURE 4 Symptoms of cork oak (*Quercus suber*) decline and dieback in Portugal caused by *Phytophthora cinnamomi*; (a) acute wilting and mortality of a cork oak in stand F61 in the Serra Algarvia; (b) severe mortality and dieback of cork oak stand F64 in the Serra Algarvia; (c) small woody roots with extensive loss of lateral and fine roots of a declining cork oak in stand F64 in the Serra Algarvia; (d) cork oak in stand F21 at Monte Morais with aerial bleeding cankers (arrows); (e) bleeding collar rot canker of a cork oak in stand F45 in the Reserva Natural da Serra da Malcata.



of *A. lusitanica* trees with decline and dieback symptoms (Table 1). The dieback of *B. celtiberica* in a natural montane forest (F17) of the Serra da Nogueira at 1043 m.a.s.l. was associated with *P. gonapodyides* and *P. pseudosyringae* (Table 1). Conversely, no *Phytophthora* species was found in the healthy natural *B. celtiberica*-*Sorbus aucuparia* stand F35 at 1428 m.a.s.l. in the Parque Nacional da Serra da Estrela (Table 1). On the Azores island of São Miguel, *P. multivora* was obtained from the soil sample of a declining non-native *Q. rubra* tree. On Madeira Island, the rhizosphere of *Ocotea foetens* and *Persea indica* in a natural montane laurisilva forest at 1161 m.a.s.l. harboured *P. chlamydospora* (Table 1). Although conifers were sampled only occasionally, several *Phytophthora*-conifer associations were found. In two Atlantic forest stands in the Parque Natural Sintra-Cascais (F50, F51), the decline of *Abies alba* trees was associated with *P. cinnamomi* (Table 1). Both *P. cinnamomi* and *P. xambivora* were recovered from the rhizosphere of a declining *P. pinaster* tree in the Parque Nacional da Peneda-Gerês (F07) in the north of Portugal whereas a declining *P. pinea* tree in an agroforestry plantation (F68) in the southernmost Serra Algarvia yielded the undescribed hybrid *P. taxon pseudocryptogea* × *kelmanii*. The only tree species without *Phytophthora* records were *J. regia*, *Laurus nobilis*, *S. aucuparia* and *Tilia cordata*. However, with only one tree each, their sample size was not representative.

3.2 | *Phytophthora* diversity in rivers and forest streams of Portugal

Using an in-situ baiting approach with rafts, 34 *Phytophthora* taxa were isolated from 33 of the 34 rivers and streams tested (97.1%).

The 425 *Phytophthora* isolates belonged to 34 taxa from six phylogenetic clades: (a) 18 non-hybrid taxa including the known species *P. citrophthora*, *P. multivora* and *P. plurivora* from Clade 2; *P. pseudosyringae* from Clade 3 (DAPC groups 3 and 5; Mullett et al., 2024); *P. amnicola*, *P. bilorbang*, *P. chlamydospora*, *P. gonapodyides*, *P. inundata*, *P. lacustris* and *P. thermophila* from Clade 6; *P. kelmanii*, *P. pseudocryptogea* and *P. ramorum* from Clade 8; and *P. honggalleglyana* (previously *P. hydropathica*; Abad et al., 2023; Hong et al., 2010) from Clade 9; a new Clade 2 species recently described as *P. pseudocitrophthora* (Jung et al., 2024); and two undescribed taxa, the designated *P. taxon* P16855 from Clade 9 and the provisionally named *P. taxon* sylvatica-like 1 from Clade 6 (Figure 6; Table 2); and (b) 16 hybrid taxa including the known species *P. xambivora* (DAPC groups 1, 5 and 9; Mullett et al., 2023) from Clade 7 and *P. xstagnum*, a Clade 6 hybrid between *P. chlamydospora* as its paternal parent and an unknown species closely related to *P. mississippiiae* as its maternal parent (Yang et al., 2014); a new Clade 2 hybrid species with *P. citrophthora* as maternal and an unknown Clade 2 taxon as paternal parent that was recently described as *P. xlusitanica* (Jung et al., 2024); the informally designated taxa *P. taxon* xWS (Oh et al., 2013), *P. taxon* *amnicola* × *chlamydospora* and *P. taxon* *thermophila* × *amnicola* (Burgess, 2015; Nagel et al., 2013) from Clade 6, *P. taxon* *pseudocryptogea* × *kelmanii* from Clade 8 and *P. taxon* xHennops from Clade 9; and eight new hybrid taxa provisionally named here as *P. taxon* xamnicola-like, *P. taxon* xlacustris-like and *P. taxon* *thermophila* × *gonapodyides* from Clade 6; *P. taxon* xpseudocryptogea from Clade 8; and *P. taxon* xClade09a1_PT1, *P. taxon* xHennops1, *P. taxon* zentmyerii × Peru4-like and *P. taxon* xP16855-like from Clade 9 (Figure 7; Table 2). In addition, *Phytophthora litorale*, a new species from the




FIGURE 5 Diversity and frequency of *Phytophthora* taxa in 70 forest stands across Portugal; P. t. = *Phytophthora* taxon.

previously unknown oomycete genus *Nothophytophthora* recently described as *N. amphigynosa* (Jung, Scanu, Bakonyi, et al., 2017), the undescribed *Phytophythium* taxa PV So7, KC-2014 and PB-2013 and the undescribed *Pythium* taxon strain 1-9 were recovered from five waterways (Table 2).

Most common were *P. gonapodyides* (13 streams=38.2%), *P. lacustris* (12 streams=35.3%), *P. plurivora* (10 streams=29.4%), *P. chlamydospora* (nine streams=26.5%) and *P. amnicola* (eight streams) followed by *P. honggalleglyana* and *P. xambivora* (five streams each=14.7%), and *P. taxon thermophila* × *amnicola* and *P. taxon* ×P16855-like (four streams each) (Figure 7; Table 2). Comprising isolates from groups DAPC9 (A2 mating type), DAPC1 (A1 mating type) and DAPC5, an A1 hybrid between DAPC1 and DAPC9 (Mullett et al., 2023), the aquatic population of *P. xambivora* was genetically more diverse than the isolates recovered from forest soils and bark cankers. *P. plurivora* was widespread in upper colline and submontane rivers and streams of the Mata Nacional do Buçaco, the National Parks Peneda-Gerês and Serra da Estrela, and the Reserva Natural da Serra da Malcata between 273 and 682 m a.s.l. (Table 2). In contrast, *P. multivora* was found only in two lowland

streams (R24, R26) near the Atlantic coast at Leiria and in the Parque Natural Sintra-Cascais (Table 2). *P. ramorum* (belonging to the EU1 lineage, according to Van Poucke et al., 2021) was detected exclusively in a small stream (R30) running through a Botanical Garden with *Rhododendron* spp. and other exotic plants established within a mixed broadleaved and pine forest in the Parque Natural Sintra-Cascais (Table 2). In Portugal, this pathogen was previously found on ornamental *Viburnum* spp. (Gomes & Amaro, 2008).

With eight *Phytophthora* taxa, the highest diversity was found in the Rio Séqua (R35), which runs through lowland to colline oak and pine forests, heathlands and *Citrus* plantations in the Algarve near Tavira, and in the Ribeira da Meimoa (sites R20 and R21) with its catchment in the Reserva Natural da Serra da Malcata covered by submontane and montane broadleaved forests, heath- and grasslands (Table 2). In the Rio Séqua, all three species from the *P. citrophthora* complex—*P. citrophthora*, *P. pseudocitrophthora* and *P. xlusitanica*—were present (Table 2). Seven *Phytophthora* taxa were detected in the Ribeira de Vale do Boi (R31) running through the foothills of the Algarvian mountain range of Monchique covered with broadleaved and pine forests, heathlands, introduced




FIGURE 6 Diversity and frequency of *Nothophytophthora* and *Phytophthora* non-hybrid taxa in 34 rivers and forest streams across Portugal; P. t. = *Phytophthora* taxon.

acacias, eucalypt plantations and horticulture (Table 2). Six *Phytophthora* species occurred in the Rio Zêzere (R12) and the Rio Beijames (R14) in the Parque Natural da Serra da Estrela and in the Rio Côa (R17 and R18) in the Reserva Natural da Serra da Malcata, all having catchment areas with diverse submontane and montane broadleaved forests, heath- and grasslands and horticulture (Table 2). Six waterways—the Rio Mondego (R15) in the Parque Natural da Serra da Estrela, the Ribeiro da Mouca (R22) in the Reserva Natural da Serra da Malcata, the forest stream R24 in the Mata Nacional de Leiria, the Ribeira de Colares (R28) and the forest stream R26 in the Parque Natural Sintra-Cascais, and the Ribeira de Odelouca near Silves in the Algarve—hosted five *Phytophthora* taxa each (Table 2).

3.3 | *Phytophthora* diversity in forest nurseries of Portugal

In 12 of the 13 forest nurseries (92.3%), a total of 135 oomycete isolates were obtained from the collar rot of one chestnut seedling and from 54 of the 87 bulked root ball samples (61.4%) from 16 of the 21 tree species (76.2%). The only nursery without a *Phytophthora* record was nursery NO1 where only two mixed soil samples from a

bare transplanting bed were tested (Table 3). The tree species without *Phytophthora* isolations were *J. regia* and the four exotic species *Chamaecyparis lawsoniana*, *Cupressus lusitanica*, *Pinus radiata* and *Quercus coccinea* (Table 3). The 122 *Phytophthora* isolates belonged to 16 taxa from eight phylogenetic clades including (a) 13 known species, namely *P. cactorum* (Clade 1), *P. multivora* and *P. plurivora* (Clade 2), *P. psychrophila* (Clade 3), *P. alticola* and *P. quercetorum* (Clade 4), *P. crassamura* and *P. gonapodyides* (Clade 6), *P. cinnamomi*, *P. parvispora* and *P. xcambivora* (Clade 7), *P. pseudocryptogea* (Clade 8) and *P. quercina* (Clade 12); (b) the informally designated *P. taxon pseudocryptogea* × *kelmanii* (Clade 8); and (c) two new hybrid taxa provisionally named here *P. taxon xcassamura-like* (Clade 6) and *P. taxon xpseudocryptogea-like* (Clade 8) (Figure 8; Table 3). In addition, 13 isolates belonging to nine taxa from three other oomycete genera were also isolated: *Elongisporangium undulatum*, *Elongisporangium* sp., *Pythium diclinum*, *Phytophythium citrinum*, *Phytophythium litorale*, *Phytophythium mercuriale*, *Phytophythium vexans*, the informally designated *Phytophythium* taxon IRAN4019C, and a previously unknown taxon provisionally named here as *Phytophythium* taxon mercuriale-like (Table 3).

Most common was *P. cinnamomi*, which was isolated in eight nurseries (61.5%) from 15 bulked root ball samples (17.2%) of nine host plants (42.9%), including *A. alba*, *A. unedo*, *C. sativa*, *Cupressus*




FIGURE 7 Diversity and frequency of *Phytophthora* hybrid taxa in 34 rivers and forest streams across Portugal; P. t. = *Phytophthora* taxon.

sempervirens, *Eucalyptus nitens*, *P. pinaster*, *Q. ilex*, *Q. rubra* and *Q. suber*, and from the collar rot of a *C. sativa* seedling (Figures 8 and 9; Table 3). *Phytophthora pseudocryptogea* was found in six nurseries (46.2%) in eight bulked root ball samples (9.2%) from seven host species (33.3%), all belonging to the genera *Pinus* and *Quercus* (*P. pinea*, *P. pinaster*, *P. sylvestris*, *Q. faginea*, *Q. ilex*, *Q. pyrenaica* and *Q. suber*; Table 3). Interestingly, the hybrids *P. taxon xpseudocryptogea*-like and *P. taxon pseudocryptogea x kelmanii* were also obtained exclusively from oak and pine plants (Figure 8; Table 3). *Phytophthora xcambivora* also occurred in six nurseries (40%) where it was isolated from eight bulked root ball samples (9.2%) of five host species (23.8%), all belonging to Fagaceae genera (*C. sativa*, *Q. ilex*, *Q. pyrenaica*, *Q. robur* and *Q. suber*; Figure 8; Table 3). The isolates belonged to groups DAPC4 and DAPC11 (both A1 mating type) and DAPC9 (A2 mating type) of the population genomic study by Mullett et al. (2023). *P. cactorum*, *P. multivora*, *P. quercetorum* and *P. quercina* were each found in three nurseries (Figure 8): *P. cactorum* in two nurseries on *P. pinea* and in one nursery on *Q. ilex*; *P. multivora* in separate nurseries on *P. pinea*, *Q. faginea* and *Q. suber*; *P.*

quercetorum on *Q. rubra* in three nurseries, *Q. pyrenaica* in two nurseries and *Q. faginea* and *Q. ilex* in one nursery each; and *P. quercina* on *A. lusitanica*, *Q. faginea*, *Q. ilex*, *Q. pyrenaica* and *Q. robur* in one nursery each (Table 3). *P. plurivora* and *P. psychrophila* were both isolated in two nurseries (Figure 8): *P. plurivora* from *Q. faginea* and *Q. suber*; and *P. psychrophila* from *A. lusitanica* and *Q. ilex* (Table 3). All remaining *Phytophthora* taxa, including *P. alticola*, which was obtained from *E. nitens*, were found in only one nursery each (Figure 8; Table 3). From five bulked root ball samples without *Phytophthora* record, *P. vexans*, *P. mercuriale*, *Phytophythium taxon mercuriale*-like and *E. undulatum* were isolated (Table 3). Nursery N11 showed the highest *Phytophthora* diversity with seven taxa detected, including *P. cinnamomi*, *P. crassamura*, *P. plurivora*, *P. pseudocryptogea*, *P. quercina*, *P. xcambivora* and *P. taxon xcrassamura*-like (Table 3). Five *Phytophthora* taxa occurred in nursery N08 whereas each of four *Phytophthora* taxa were found in five nurseries (N02, N06, N07, N12 and N13; Table 3).

Although in most nurseries, the majority of plants appeared visually healthy, plants with symptoms indicative of *Phytophthora*

infections were present in all nurseries. Symptoms comprised fine root losses and root rot (found in almost all plants with above-ground symptoms), collar rot lesions (only found in *C. sativa*), stunted growth, small sized and/or chlorotic leaves/needles, wilting and mortality (Figure 9; Table 3). Overall, 44 of the 50 bulked samples taken from symptomatic or partly symptomatic plants (88%) yielded 16 *Phytophthora* taxa. In contrast, only nine of the 35 bulked samples taken from visually healthy plants (25.7%) were found infested with five *Phytophthora* species (Table 3). All bulked root ball samples from which *P. cactorum*, *P. cinnamomi*, *P. multivora*, *P. plurivora*, *P. pseudocryptogea* and its hybrids have been isolated came, at least partially, from symptomatic plants (Table 3). From two of the six symptomatic samples without a *Phytophthora* record, *E. undulatum*, *P. mercuriale* and *Phytophthium* taxon *mercuriale*-like were recovered. The latter taxon was also obtained from two samples of healthy plants (Table 3).

4 | DISCUSSION

This paper reports on the most extensive and diverse *Phytophthora* survey conducted in natural ecosystems and forest nurseries of a country in Europe. Across Portugal, 48 *Phytophthora* taxa from eight phylogenetic clades were obtained from 62 of 70 forest stands (88.6%), 33 of 34 rivers and forest streams (97.1%) and 12 of 13 nurseries (92.3%) tested. These comprised 28 known *Phytophthora* species; three new recently described species (*P. castanetorum*, *P. pseudocitrophthora* and *P. xlusitanica*; Jung et al., 2024; Jung, Horta Jung, Cacciola, et al., 2017); one informally designated and two new non-hybrid taxa; five informally designated and nine new hybrid taxa. Only five *Phytophthora* species (10.4%)—*P. pseudosyringae* and *P. psychrophila* from Clade 3, *P. uliginosa* from Clade 7, and *P. castanetorum* and *P. quercina* from Clade 12—are considered native to Europe (Jung et al., 2002, 2003, 2016; Jung, Horta Jung, Cacciola, et al., 2017; Mullett et al., 2024). Another nine taxa (18.8%)—*P. bilorbang*, *P. chlamydospora*, *P. crassamura*, *P. gonapodyides*, *P. inundata*, *P. lacustris*, the new *P.* taxon Clade06b_PT1, *P.* taxon *xcrassamura*-like and *P.* taxon *xlacustris*-like (all from Clade 6)—are of cryptic origin (Aghighi et al., 2012; Brasier, Cooke, et al., 2003; Brasier, Sanchez-Hernandez, & Kirk, 2003; Jung et al., 2016; this study). However, the remaining majority of 34 *Phytophthora* taxa (70.8%) are considered in Europe as introduced aliens. Because the international trade in living plants is considered a major pathway for the introduction of exotic *Phytophthora* pathogens (Brasier et al., 2022; Jung et al., 2016, 2024; Liebhold et al., 2012; Santini et al., 2013), the unique diversity of exotic *Phytophthora* taxa in Portugal is most probably related to the long-term import of living plants from overseas that started more than 500 years ago with the establishment of trading posts and colonies in Brazil and across Africa, Southeast and East Asia, and has intensified during the past decades with modern globalization.

A phylogeographic study of a global population of *P. cinnamomi* isolates demonstrated recently that this wide-host range pathogen

originates from Taiwan, the previous Portuguese colony of Formosa, and possibly other regions in Southeast Asia (Shakya et al., 2021). Another recent phylogeographic study indicates for *P. xambivora* an origin in Japan and most likely also in neighbouring regions of East Asia (Mullett et al., 2023). Early reports of typical symptoms of ink disease of *C. sativa* suggest that the causal agents *P. cinnamomi* and *P. xambivora* were already established in Portugal in the 1830s (Crandall et al., 1945). Eventually, the association of both pathogens with declining chestnuts in Portugal was confirmed in the 1940s (Lopes-Pimentel, 1946, 1947). Numerous isolates of *P. cinnamomi* and *P. xambivora* were included in the population genomic studies of Shakya et al. (2021) and Mullett et al. (2023), respectively. These demonstrated comparatively high genetic variability of the Portuguese populations of *P. cinnamomi*, comprising both clonal A2 mating type lineages PcG1-A2 and PcG2-A2 that are driving the global pandemic (Shakya et al., 2021), and *P. xambivora*, comprising five of the 11 global DAPC groups and both mating types (Mullett et al., 2023; this study), indicating multiple introductions of both pathogens. Also for *P. parvispora*, the closest relative of *P. cinnamomi* in Clade 7c; the Clade 2 species *P. citrophthora*, *P. pseudocitrophthora*, *P. plurivora* and *P. xlusitanica*; and the Clade 8 species *P. kelmanii* and *P. pseudocryptogea* including their interspecific hybrids, and *P. ramorum*, an origin in Southeast and/or East Asia is likely due to their occurrence in natural healthy ecosystems and/or the origin of their closest relatives in this region (Jung et al., 2020, 2021, 2024; Jung, Chang, Bakonyi, et al., 2017; Scanu, Hunter, et al., 2014). Several Clade 6 isolates from the Rio Séqua in the south of Portugal have ITS and *cox1* sequences identical to those of *P.* taxon *sylvatica*-like 1, a natural inhabitant of forest streams in Vietnam (Jung et al., 2020). *P. alticola* from Clade 4 was described from exotic eucalypt plantations in South Africa (Maseko et al., 2007) and was later also found widespread in *Acacia* plantations and natural forests (Bose et al., 2018, 2021). It is closely related to the eucalypt pathogens *P. arenaria* and *P. boodjera* from Western Australia (Bose et al., 2017; Rea et al., 2011; Simamora et al., 2015). From Australia are the mostly aquatic Clade 6 species *P. amnicola*, *P. thermophila* and their hybrid *P.* taxon *thermophila* × *amnicola* (Burgess, 2015; Crous et al., 2012; Jung et al., 2011), which were all found widespread in Portuguese rivers and streams during this study. Another aquatic Clade 6 hybrid found in Portuguese streams, *P.* taxon *xWS* (= *P. amnicola* × *P. thermophila*), most likely originates from South Africa (Burgess, 2015; Nagel et al., 2013). However, it cannot be ruled out that the hybridizations that created *P.* taxon *thermophila* × *amnicola* and *P.* taxon *xWS* occurred spontaneously in Portuguese waterbodies where their introduced parental species were found widespread in this survey. The emerging wide-host range pathogen *P. multivora* from Clade 2 is also native to South Africa as has been unveiled recently by a population genomic study (Tsykun et al., 2022). Another recent population genetic study presented evidence that the global wide-host range pathogen *P. cactorum* from Clade 1 most likely originates from North America (Bourret et al., 2022). Likewise, the oak-specific Clade 4 species *P. quercetorum* (Balci et al., 2008), the aquatic Clade 6 hybrid *P. xstagnum* (Yang et al., 2014) and *P. honggalleglyana* from




FIGURE 8 Diversity and frequency of *Phytophthora* taxa in 13 forest nurseries across Portugal; *P. t.* = *Phytophthora* taxon.

a cluster of aquatic high-temperature species within Clade 9 (Abad et al., 2023; Hong et al., 2010) are native to North America. From this Clade 9 cluster, an array of six hybrid and non-hybrid taxa were obtained from Portuguese rivers and streams which, due to their own or their closest relatives' occurrence in natural undisturbed rivers and forests, most likely have their origins in South Africa (*P. taxon* ×Hennops, *P. taxon* ×Hennops-like 1; Oh et al., 2013) and South America like *P. taxon* P16855, first found in an Amazonian rainforest in Ecuador (*cox1* GenBank accession GU594797), and three related new hybrid taxa *P. taxon* ×P16855-like, *P. taxon* ×Clade09a1_PT1 and *P. taxon* zentmyerii × Peru4-like.

Figure 10 summarizes the distribution of *Phytophthora* taxa across the three site categories surveyed. While 35 *Phytophthora* taxa were exclusively found in forests (five taxa), forest nurseries (six taxa) and river systems (24 taxa), 13 *Phytophthora* taxa occurred in more than one site category. The primarily aquatic *P. gonapodyides* with cryptic geographic origin was found at all site categories. Five introduced taxa—*P. multivora*, *P. plurivora*, *P. xambivora*, *P. taxon pseudocryptogea* × *kelmanii* and *P. taxon xpseudocryptogea-like*—also occurred at all site categories, indicating their spread from

nurseries to both forests (via planting of infested nursery stock) and river systems (via nursery effluents or indirectly via infested plantings). For *P. xambivora* this is supported by the higher genetic variability of nursery populations (four DAPC groups) compared with river (three DAPC groups) and forest (one DAPC group) populations. However, due to the use of unfiltered non-sterile river water for irrigation in several nurseries included in this study, an introduction of *Phytophthora* taxa to the nurseries via infested river water cannot be ruled out. Besides the six *Phytophthora* taxa recorded from all three site categories, nurseries shared *P. pseudocryptogea* with rivers/streams and another three *Phytophthora* species, the invasive *P. cinnamomi* and the native *P. psychrophila* and *P. quercina*, with forest sites. Forest sites and river systems also had two primarily aquatic species, *P. amnicola* and *P. chlamydospora*, and the forest pathogen *P. pseudosyringae* in common.

Over all 29 tree species sampled in forest stands across Portugal, isolation rates from rhizosphere soil of declining trees exceeded those from visually healthy trees considerably (65.3% versus 10.1%), demonstrating a clear association between *Phytophthora* presence and decline and dieback of trees. Our survey showed that *P.*

FIGURE 9 Container plants in forest nurseries of Portugal; (a) *Castanea sativa* in nursery N11; healthy seedlings with yellow leaves due to autumnal senescence and dead seedlings with brown leaves due to root and collar rot caused by *Phytophthora cinnamomi* and *P. x cambivora*; (b) *C. sativa* seedling in nursery N11 with a girdling collar rot lesion (arrow) caused by *P. cinnamomi*; (c) *C. sativa* seedling in nursery N05 with a girdling collar rot lesion caused by *P. cinnamomi*; note the tongue-shaped inner bark lesion (arrow); (d) high mortality of *Quercus ilex* seedlings caused by *P. cinnamomi* in nursery N08; note that the containers are standing directly on the wet ground; (e) wilting of *Arbutus unedo* seedlings in nursery N02 due to root rot caused by *P. cinnamomi*; (f) chlorosis and mortality of *Pinus sylvestris* seedlings in nursery N02 due to root rot caused by *P. pseudocryptogea* and *P. taxon* x *pseudocryptogea*-like; (g) *Quercus robur* container stand in nursery N12 infested by *P. gonapodyides*; note the close distance to the ground and the debris splashed onto the containers (arrows).



cinnamomi is by far the most common *Phytophthora* species in forests and forest nurseries across Portugal, with infestation rates of 55.7% and 61.4%, respectively, and an altitudinal range of 123–1194 m.a.s.l. The high *P. cinnamomi* infestation rates of forest sites and forest nurseries are most probably correlated via the planting of infested nursery stock. The invasive success of this self-sterile species in Portugal and numerous other countries with hot and dry summers like Australia, California, Italy, Spain and South Africa (Bose et al., 2018; Burgess et al., 2017; Erwin & Ribeiro, 1996; Hardham & Blackman, 2018; Jung, Pérez-Sierra, et al., 2018; Oh et al., 2013; Sanchez et al., 2002; Vettraino et al., 2005) is certainly facilitated by the ability of the A2 mating type to form oospores, highly enduring thick-walled survival structures, by selfing, in particular under environmental stress or in the presence of antagonistic fungi and plants (Brasier, 1978; Jayasekera et al., 2007; Jung et al., 2013). *P. cinnamomi* is arguably the most notorious invasive plant pathogen globally, causing root and collar rot, epidemic dieback and mortality of more than 5000 woody species in natural ecosystems, nurseries and planted stands (Brasier et al., 2022; Erwin & Ribeiro, 1996; Hardham & Blackman, 2018). In all 12 forest tree species from which it was recovered in forest stands, *P. cinnamomi* was associated with severe symptoms like fine root losses, root rot, decline, dieback and mortality, and in some stands of *C. sativa* and *Q. suber* also with bleeding

bark cankers. With eight tree species in 15.7% of forest sites and 46.2% of forest nurseries, the other wide-host range species from Clade 7, *P. x cambivora*, was also common in Portugal. Like in other European countries (Day, 1938; Jung, Pérez-Sierra, et al., 2018; Vettraino et al., 2005) and Chile (Jung, Durán, et al., 2018), *P. cinnamomi* and *P. x cambivora*, either separately or jointly, were responsible for ink disease and mortality of *C. sativa* in 16 of the 17 affected stands (94.1%) in this study. The severity of this epidemic in Portugal was demonstrated by Martins et al. (2007) who reported that both pathogens caused an average mortality of 58.6% in around 90 chestnut stands between 1995 and 2004. Considering their aggressiveness to *C. sativa*, the presence of *P. cinnamomi* and/or *P. x cambivora* in 67% of the chestnut container stands tested in forest nurseries in this survey is concerning. *P. cinnamomi* was also recovered from *Q. suber* and *Q. ilex* trees with decline and dieback in 15 of the 19 affected stands (78.9%), reinforcing previous studies that demonstrated the involvement of *P. cinnamomi* in the devastating decline of both oak species in Portugal (Brasier et al., 1993; Moreira & Martins, 2005) and other countries (Gallego et al., 1999; Jung, Pérez-Sierra, et al., 2018; Robin et al., 1998; Sanchez et al., 2002; Seddaiu et al., 2020; Vettraino et al., 2002). In addition, *P. quercina* and in some stands also *P. pseudocryptogea*, *P. psychrophila* and *P. uliginosa* were isolated for the first time from the rhizosphere of declining




FIGURE 10 Venn diagram showing the exclusive and overlapping *Phytophthora* diversity in 70 forest stands, 34 rivers and forest streams and 13 forest nurseries across Portugal. Black and yellow taxa names indicating the first reports for Europe and Portugal, respectively. First reports from this survey previously reported: *Jung et al. (2016), **Jung, Horta Jung, Cacciola, et al. (2017), ***Jung et al. (2024).

cork and holm oaks in Portugal. Their involvement in cork and holm oak decline was previously demonstrated in eastern Spain (Pérez-Sierra et al., 2013) and Italy (Jung et al., 1999; Seddaiu et al., 2020). Although commonly considered as specific to oaks, *P. quercina* was also recovered from the rhizosphere of tree species from other genera, namely *A. unedo*, *C. sativa*, *P. latifolia* and *I. aquifolium*, in several forest stands. Pathogenicity trials using a standardized soil infestation method are needed to clarify whether this fine root pathogen is pathogenic to these four tree species. The *Phytophthora*-mediated declines of *F. sylvatica* and *Q. robur*, found in several stands during this survey, are widespread in temperate regions of Europe and parts of Italy and the Balkans (Balci & Halmschlager, 2003; Cacciola et al., 2005; Corcobado et al., 2020; Day, 1938; Hansen & Delatour, 1999; Jönsson et al., 2003; Jung, 2009; Jung et al., 1996, 2000, 2019; Jung, Pérez-Sierra, et al., 2018; Milenković et al., 2012; Stępniewska & Dłuszyński, 2010), but were previously not known to occur in Portugal. Other first records from non-riparian forests in Portugal included (a) the decline and dieback of the Iberian *Q. pyrenaica*, which was mainly associated with *P. cinnamomi*, *P. quercina* and *P. xcambivora*; the findings of (b) the emerging wide-host range pathogen *P. multivora* from a declining *Q. rubra* tree on the Azores island of São Miguel and from the invasive *Acacia dealbata* and declining *A. pseudoplatanus* trees in several stands of the Parque Natural Sintra-Cascais; (c) *P. cinnamomi* from declining mature silver fir (*A. alba*) trees in the same Natural Park; (d) *P. plurivora* in various forest stands of Portugal; (e) *P. xcambivora*, *P. gonapodyides* and *P.*

pseudosyringae from declining *B. celtiberica* stands; (f) *P. xcambivora* from *S. caprea* and *P. pinaster* with dieback; and (g) *P. chlamydospora* from *O. foetens* and *P. indica* in an Atlantic laurisilva forest on the island of Madeira. Overall, 15 and 21 new *Phytophthora*-associated decline and dieback diseases of trees were recorded for the first time in Europe and Portugal, respectively (Table S5).

Following the first reports of *P. xalni* from bark cankers of declining planted *Alnus glutinosa* trees on the banks of the Rio Tâmega in Chaves (Jung et al., 2016) and declining mature *A. lusitanica* trees along a small stream in central Portugal (Kanoun-Boulé et al., 2016), bleeding bark cankers, decline and dieback of riparian *A. lusitanica* trees were commonly observed during this study along several rivers and streams in northern and central Portugal. At the Alva, Côa, Fervença and Tâmega Rivers where mortality rates along some river sections exceeded 50%, the disease was caused by the aggressive alder-specific hybrid pathogen *P. xalni* in accordance with the situation in other European countries (Brasier et al., 2004; Černý & Strnadová, 2010; Corcobado et al., 2023; Gomes Marques et al., 2024; Haque et al., 2015; Jung & Blaschke, 2004; Jung, Pérez-Sierra, et al., 2018; Nagy et al., 2003; Schumacher et al., 2006; Solla et al., 2010; Thoirain et al., 2007; Trzewik et al., 2015). Our findings of *P. cinnamomi* and *P. gonapodyides* in the rhizosphere of declining *A. lusitanica* trees along several streams with scattered alder dieback confirm earlier reports of these pathogens from alders in northern Spain, Germany and Austria (Corcobado et al., 2023; Jung et al., 2016; Jung & Blaschke, 2004). In southern Portugal, *Phytophthora*-related

dieback of riparian alder forests was not observed, which is in accordance with a recent model that predicts absence of *P. xalni* in this region due to summer temperatures exceeding the maximum temperature of the pathogen (Brasier et al., 2004; Gomes Marques et al., 2024; Jung, Horta Jung, Scanu, et al., 2017). In a recent survey in five riparian alder stands in northern Portugal, *P. cinnamomi*, *P. gonapodyides* and nine other *Phytophthora* species, including three species not found in our survey (*P. asparagi*, *P. polonica* and *P. rosacearum*) were found in the rhizosphere of declining alder trees whereas *P. multivora* and *P. plurivora* were isolated from bark cankers at two sites (Bregant et al., 2023a). *P. plurivora* was isolated earlier from bark cankers of *A. glutinosa* in Germany (Jung & Blaschke, 2004; Jung & Burgess, 2009) and was also found widespread in the rhizosphere of riparian alder stands in Austria, Germany, Poland, Slovakia and Turkey (Aday Kaya et al., 2018; Corcobado et al., 2023; Jung & Blaschke, 2004; Tkaczyk et al., 2023; Trzewik et al., 2015). Noteworthy, *P. uniformis*, paternal parent of *P. xalni* and contributing causal agent of the Europe-wide alder epidemic (Brasier et al., 2004; Corcobado et al., 2023; Husson et al., 2015; Jung & Blaschke, 2004; Jung, Pérez-Sierra, et al., 2018; Macháčová et al., 2024), was not detected in this or any previous survey in Portugal. Its absence is most probably related to the low cardinal temperatures of *P. uniformis* as an adaptation to the temperate and sub-boreal climates in its native habitats in the Pacific Northwest and Alaska (Aguayo et al., 2013; Černý et al., 2012; Jung, Horta Jung, Scanu, et al., 2017; Jung, Pérez-Sierra, et al., 2018; Redondo et al., 2015; Schumacher et al., 2006; Sims et al., 2015), rendering the pathogen unsuited for the high summer temperatures in Portugal.

Overall, a diverse community of 20 *Phytophthora* taxa has been unveiled in rhizosphere soil and bark cankers of forest stands in Portugal by this study (17 taxa) and Bregant et al. (2023a, 2023b, nine species). Using similar or identical isolation methods, slightly lower *Phytophthora* diversities were found in various surveys in Spain (13 species; Brasier et al., 1993; Corcobado et al., 2010; Pérez-Sierra et al., 2013; Sanchez et al., 2002; Solla et al., 2010; Štraus et al., 2023; Vettraino et al., 2005), Austria (14 species; Balci & Halmschlager, 2003; Corcobado et al., 2020, 2023), France (17 taxa; Hansen & Delatour, 1999; Husson et al., 2015; Jung et al., 2002, 2003; Robin et al., 1998, 2011; Schenck et al., 2018; Vettraino et al., 2005) and Germany (17 taxa; Jung, 2009; Jung & Blaschke, 2004; Jung & Nechwatal, 2008; Jung et al., 2000, 2002, 2003; Peters et al., 2019; Schumacher et al., 2006). The European country with the highest *Phytophthora* diversity in forest stands, however, is Italy, including the islands of Sardinia and Sicily, where 38 *Phytophthora* species were isolated from rhizosphere soil, bark cankers, leaf and twig blights in numerous surveys (Bregant et al., 2020; Cacciola et al., 2005; Ginetti et al., 2014; Jung et al., 2019, 2024; Jung, Horta Jung, Cacciola, et al., 2017; Jung, Pérez-Sierra, et al., 2018; Riolo et al., 2020; Scanu et al., 2015; Scanu, Hunter, et al., 2014; Scanu, Linaldeddu, et al., 2014; Seddaiu et al., 2020; Vettraino et al., 2002, 2005).

Rivers and streams in Asia, Australia, Europe, the Americas and South Africa were shown to be inhabited by a wide range of

Phytophthora species (Caballot et al., 2024; Català et al., 2015; Corcobado et al., 2023; Dunstan et al., 2016; Huai et al., 2013; Hüberli et al., 2013; Jung et al., 2019, 2020, 2022, 2024; Jung, Chang, Bakonyi, et al., 2017; Jung, Durán, et al., 2018; Oh et al., 2013; Reeser et al., 2011; Shrestha et al., 2013). In our survey in Portugal, 34 *Phytophthora* taxa were detected in 33 of the 34 rivers and forest streams (97.1%) tested. These comprised 18 non-hybrid and 16 hybrid taxa, among them two and eight taxa, respectively, previously unknown to science. As in other river surveys listed before, *P. cinnamomi* was not recovered from any Portuguese stream, emphasizing its strictly soilborne lifestyle. Because surface water is commonly used for irrigation of nursery beds and horticultural plantations, the presence of the notorious pathogens *P. plurivora* and *P. x cambivora* in 29.4% and 14.7% of the streams, the detection of *P. multivora* and *P. ramorum* in two and one stream, respectively, and the widespread occurrence of 10 new taxa with unknown host ranges are of particular concern. Bregant et al. (2023a) reported in total six *Phytophthora* species from five streams in central Portugal including *P. multivora*, four Clade 6 species and the Clade 9 species *P. polonica*, which was not found in the present study. The widespread occurrence of a diverse community of 16 hybrid taxa from Clades 2, 6, 8 and 9 in our stream survey confirms earlier studies showing that aquatic ecosystems provide ideal conditions for interspecific *Phytophthora* hybridizations and the subsequent survival of the hybrids (Burgess, 2015; Jung et al., 2020, 2024; Jung, Chang, Bakonyi, et al., 2017; Jung, Durán, et al., 2018; Nagel et al., 2013; Oh et al., 2013). It is becoming increasingly clear that in the genus *Phytophthora*, interspecific hybridizations are a common evolutionary feature facilitating the adaptation to new environmental conditions and new host species (Brasier et al., 2004; Burgess, 2015; Jung et al., 2020, 2024; Jung, Chang, Bakonyi, et al., 2017; Jung, Horta Jung, Scanu, et al., 2017; Van Poucke et al., 2021). The use of infested surface water for irrigation of nurseries and agricultural, horticultural and forest plantings poses, therefore, an inherent risk of creating new *Phytophthora*-host combinations and potentially new epidemics. The occurrence of the Clade 8 hybrids *P. taxon pseudocryptogea* × *kelmanii* and *P. taxon* × *pseudocryptogea*-like in several streams and in one nursery each might suggest an introduction to the nurseries with irrigation water taken from infested rivers.

An extensive study in 23 European countries revealed the presence of (a) 49 *Phytophthora* taxa in 81% of the almost 2000 beds and container stands in 92% of the 732 nurseries tested and (b) 56 *Phytophthora* taxa in 66% of the more than 2500 planting sites tested, clearly demonstrating that the planting of infested nursery stock is the primary pathway of exotic *Phytophthora* pathogens to the wider environment (Jung et al., 2016). The efficacy and detrimental effects of *Phytophthora* spread to natural ecosystems via the nursery pathway were also demonstrated for restoration outplantings in heathlands and woodlands in California (Frankel et al., 2020; Garbelotto et al., 2018; Rooney-Latham et al., 2019; Sims & Garbelotto, 2021). Recently, using both baiting and metabarcoding approaches, 12 and 45 *Phytophthora* taxa, respectively, were reported from soil and water samples in eight Czech nurseries, among

them 15 taxa previously unknown to science (Bačová et al., 2024). In the present study, 16 *Phytophthora* taxa were detected in 12 of the 13 nurseries, which is comparable to the diversity reported for Spanish nurseries by Moralejo et al. (2009; 17 *Phytophthora* taxa) and Mora-Sala et al. (2022; 14 *Phytophthora* taxa). Besides the high infestation rates with the aggressive wide-host range pathogens *P. cinnamomi*, *P. pseudocryptogea* and *P. xambivora*, the findings of two other aggressive pathogens with wide-host ranges, namely *P. multivora* and *P. plurivora*, and the host-specific pathogens *P. alticola*, *P. quercetorum* and *P. quercina*, are particularly concerning. The oak-specific fine root pathogen *P. quercetorum* is involved in temperate oak decline in the eastern United States (Balci et al., 2008) and, hence, its spread with infested nursery plants to forests could exacerbate the decline of oak forests in Portugal. *P. alticola* was described from eucalypt plantations in South Africa (Maseko et al., 2007) but its status has been doubtful due to the loss of all isolates associated with the original description except one ex-paratype; discrepancies between sequence data and morphology of the single remaining isolate and the original species description; and inconclusiveness of the holotype material of *P. alticola* (Simamora et al., 2015). Based on a thorough morphological and phylogenetic study of the only existing ex-paratype isolate and a collection of new isolates from plantations and forests across South Africa, Bose et al. (2017) demonstrated that *P. alticola* and the Western Australian eucalypt pathogen *P. boodjera* (Simamora et al., 2015) are distinct, though very closely related, sister species and provided an emended description for *P. alticola*. During a survey conducted between 2012 and 2020 in numerous *Eucalyptus globulus* plantations across Portugal, crown symptoms indicative of root and collar rot were observed and, besides *P. cinnamomi*, *P. alticola* was obtained from 10 plantations (Diogo et al., 2023). Our results indicate that *P. alticola* might have been introduced to the eucalypt plantations with infested nursery plants. In another study of eucalypt plantations in Portugal, Bregant et al. (2023b) reported *P. hibernalis*, *P. multivora* and *P. niederhauserii* causing root rot and bleeding cankers on *E. globulus*. The aggressive eucalypt pathogens *P. alticola*, *P. cinnamomi*, *P. hibernalis*, *P. multivora* and *P. niederhauserii* could pose a serious threat to the Portuguese paper and pulp industry, which is primarily based on timber produced on 845,000 ha of domestic eucalypt plantations (ICNF, 2019).

In Portuguese forest nurseries, both infestation rate (88%) and *Phytophthora* diversity (16 taxa) were higher in symptomatic plants, but 26% of samples from visually healthy plants also yielded a total of five *Phytophthora* species. Furthermore, in container stands with *Phytophthora* isolations from symptomatic plants, many healthy-looking plants were almost certainly also infested with *Phytophthora*, as has been demonstrated by Jung et al. (2016) and Migliorini et al. (2015). The almost ubiquitous infestation of forest and other nurseries in Portugal and elsewhere in Europe with almost 100 known and unknown *Phytophthora* taxa will inevitably cause further spread of these pathogens to the wider environment, increasing the pathogen pressure on forest ecosystems and thus weakening their resilience to climatic extremes and decreasing the sustainability of their multiple functions to society.

This is particularly concerning as the newly approved EU regulation 2024/1991 (Nature Restoration Law; Anonymous, 2024), aiming to ensure the recovery of biodiverse and resilient nature across the EU territory, obliges the member states to perform extensive measures on nature restoration including re-establishing numerous listed habitat types in areas where those habitat types do not occur and planting of at least three billion additional trees by 2030 at Union level. Consequently, to prevent rapid spread of aggressive invasive *Phytophthora* and other plant pathogens to highly sensitive natural areas across Europe, it is paramount at the EU level to avoid *Phytophthora* infestations of nursery stock by immediately implementing an array of mandatory management measures encompassing: (a) the use of thermosterilized substrates and disinfested containers/pots and tools; (b) growing plants from seeds or, in case this is not possible, the use of non-infested plant material (confirmed by molecular tests rather than visual inspections); (c) lifting container stands at least 80 cm from the ground with well-drained surfaces underneath to avoid splash infections from the ground; and (d) irrigation with tap water and water from deep wells or, where it cannot be avoided, surface and recirculation water only after filtering with sand, lava grain or pumice filter systems or disinfestation with heat, ultrasonic, chlorination or ozone (Parke & Grünwald, 2012; Pérez-Sierra & Jung, 2013; Stewart-Wade, 2011; Ufer et al., 2008). In addition, the use of fungicides and fungistatic chemicals like potassium phosphite should be avoided as much as possible because their regular use causes the rapid evolution of resistant strains and the suppression of disease symptoms, thus resulting in the spread of visually healthy *Phytophthora*-infested plants (Jung et al., 2016; Migliorini et al., 2015; Pérez-Sierra & Jung, 2013). The Nursery Industry Accreditation Scheme Australia (NIASA; <https://nurseryproductionfms.com.au/niasa-accreditation/>) and the Avocado Nursery Voluntary Accreditation Scheme (ANVAS; <https://avocado.org.au/our-programs/anvas/>) in Australia are excellent examples of how successful accreditation schemes can be (Anonymous, 2018, 2021).

ACKNOWLEDGEMENTS

The authors are grateful to the Fundação para a Ciência e a Tecnologia (FCT) for co-financing the European BiodivERsA project RESIPATH (Responses of European Forests and Society to Invasive Pathogens, BIODIVERSA/0002/2012); the Project *Phytophthora* Research Centre Reg. No. CZ.02.1.01/0.0/0.0/15_003/0000453 co-financed by the Czech Ministry for Education, Youth and Sports and the European Regional Development Fund; the grant MIUR-FIRB 2010 (RBF10PZ4N) 'Metagenomic strategies to assess genetic diversity in soil-borne *Phytophthora* species' from the Italian Ministry of Education, University and Research (MIUR). The authors express their particular gratitude to the Instituto da Conservação da Natureza e das Florestas (ICNF) (<https://www.icnf.pt>) for providing data and knowledge about Portuguese forests, the permission to sample in protected areas and logistic support. We specifically thank Dina Ribeiro (ICNF, Divisão de Fitossanidade Florestal) and numerous ICNF rangers for their dedicated assistance during the field trips.

CONFLICT OF INTEREST STATEMENT

The authors declare that there is no conflict of interest.

DATA AVAILABILITY STATEMENT

For all oomycete taxa found in this study, ITS and *cox1* sequences from representative isolates are available from GenBank at <https://www.ncbi.nlm.nih.gov/genbank/>, and accession numbers are given in Table S1.

ORCID

Marília Horta Jung  <https://orcid.org/0000-0003-2219-8647>

Cristiana Maia  <https://orcid.org/0000-0002-8654-2905>

Beatriz Mora-Sala  <https://orcid.org/0000-0002-9734-9481>

Paloma Abad-Campos  <https://orcid.org/0000-0002-5979-4349>

Leonardo Schena  <https://orcid.org/0000-0002-9737-2593>

Saveria Mosca  <https://orcid.org/0000-0003-4985-8641>

Salvatore Moricca  <https://orcid.org/0000-0003-3097-559X>

Jan Nechwatal  <https://orcid.org/0000-0001-9977-7976>

Lídia Dionísio  <https://orcid.org/0000-0002-0530-9395>

Alfredo Cravador  <https://orcid.org/0000-0002-9831-9815>

Thomas Jung  <https://orcid.org/0000-0003-2034-0718>

REFERENCES

- Abad, Z.G., Burgess, T.I., Bourret, T., Bensch, K., Cacciola, S.O., Scanu, B. et al. (2023) *Phytophthora*: taxonomic and phylogenetic revision of the genus. *Studies in Mycology*, 106, 259–348.
- Aday Kaya, A.G., Lehtijärvi, A., Şaşmaz, Y., Nowakowska, J.A., Oszako, T., Lehtijärvi, H.T.D. et al. (2018) *Phytophthora* species detected in the rhizosphere of *Alnus glutinosa* stands in the floodplain forests of Western Turkey. *Forest Pathology*, 48, e12470.
- Aghighi, S., Hardy, G.E.S.J., Scott, J.K. & Burgess, T.I. (2012) *Phytophthora bilorbang* sp. nov., a new species associated with the decline of *Rubus anglocandicans* (European blackberry) in Western Australia. *European Journal of Plant Pathology*, 133, 841–855.
- Aguayo, J., Adams, G.C., Halkett, F., Catal, M., Husson, C., Nagy, Z.A. et al. (2013) Strong genetic differentiation between north American and European populations of *Phytophthora alni* subsp. *uniformis*. *Phytopathology*, 103, 190–199.
- Anonymous. (2018) *Implementation of recommendations from the avocado nursery voluntary accreditation scheme review*. Sydney: Horticulture Innovation Australia Limited (Hort Innovation).
- Anonymous. (2021) *NIASA Best Management Practice Guidelines*. Sydney: Greenlife Industry Australia.
- Anonymous. (2024) Regulation (EU) 2024/1991 of the European Parliament and of the council of 24 June 2024 on nature restoration and amending regulation (EU) 2022/869. *Official Journal of the European Union, L Series*, 29(7), 2024.
- Arianoutsou, M., Bazos, I., Christopoulou, A., Kokkoris, Y., Zikos, A., Zervou, S. et al. (2021) Alien plants of Europe: introduction pathways, gateways and time trends. *PeerJ*, 9, e11270.
- Báčova, A., Cooke, D.E.L., Milenković, I., Májek, T., Nagy, Z.A., Corcobado, T. et al. (2024) Hidden *Phytophthora* diversity unveiled in tree nurseries of the Czech Republic with traditional and metabarcoding techniques. *European Journal of Plant Pathology*, 170, 131–156.
- Balci, Y., Balci, S., Blair, J.E., Park, S.-Y., Kang, S. & MacDonald, W.L. (2008) *Phytophthora quercetorum* sp. nov., a novel species isolated from eastern and north-central USA oak forest soils. *Mycological Research*, 112, 906–916.
- Balci, Y. & Halmshlager, E. (2003) Incidence of *Phytophthora* species in oak forests in Austria and their possible involvement in oak decline. *Forest Pathology*, 33, 157–174.
- Bienapfl, J.C. & Balci, Y. (2014) Movement of *Phytophthora* spp. in Maryland's nursery trade. *Plant Disease*, 98, 134–144.
- Birmingham, D. (2000) *Trade and empire in the Atlantic, 1400–1600*. London, UK: Routledge.
- Bose, T., Burgess, T.I., Roux, J. & Wingfield, M.J. (2017) *Phytophthora alticola*; revised description based on new collections and a neotype. *Sydowia*, 69, 161–170.
- Bose, T., Wingfield, M.J., Roux, J., Vivas, M. & Burgess, T.I. (2018) Community composition and distribution of *Phytophthora* species across adjacent native and non-native forests of South Africa. *Fungal Ecology*, 36, 17–25.
- Bose, T., Wingfield, M.J., Roux, J., Vivas, M. & Burgess, T.I. (2021) *Phytophthora* species associated with roots of native and non-native trees in natural and managed forests. *Microbial Ecology*, 81, 122–133.
- Bourret, T.A., Fajardo, S.N., Frankel, S.J. & Rizzo, D.M. (2023) Cataloging *Phytophthora* species from agriculture, forests, horticulture and restoration outplantings in California, U.S.A.: a sequence-based meta-analysis. *Plant Disease*, 107, 67–75.
- Bourret, T.B., Fajardo, S.N., Engert, C.P. & Rizzo, D.M. (2022) A barcode-based phylogenetic characterization of *Phytophthora cactorum* identifies two cosmopolitan lineages with distinct host affinities and the first report of *Phytophthora pseudotsugae* in California. *Journal of Fungi*, 8, 303.
- Brasier, C.M. (1978) Stimulation of oospore formation in *Phytophthora* by antagonistic species of *Trichoderma* and its ecological implications. *Annals of Applied Biology*, 89, 135–139.
- Brasier, C.M. (2008) The biosecurity threat to the UK and global environment from international trade in plants. *Plant Pathology*, 57, 792–808.
- Brasier, C.M., Cooke, D.E.L., Duncan, J.M. & Hansen, E.M. (2003) Multiple new phenotypic taxa from trees and riparian ecosystems in *Phytophthora gonapodyides* – *P. megasperma* ITS clade 6, which tend to be high-temperature tolerant and either inbreeding or sterile. *Mycological Research*, 107, 277–290.
- Brasier, C.M., Franceschini, S., Vettraino, A.M., Hansen, E.M., Green, S., Robin, C. et al. (2012) Four phenotypically and phylogenetically distinct lineages in *Phytophthora lateralis*. *Fungal Biology*, 116, 1232–1249.
- Brasier, C.M., Kirk, S.A., Delcan, J., Cooke, D.E.L., Jung, T. & Man in't Veld, W.A. (2004) *Phytophthora alni* sp. nov. and its variants: designation of emerging heteroploid hybrid pathogens spreading on *Alnus* trees. *Mycological Research*, 108, 1172–1184.
- Brasier, C.M., Robredo, F. & Ferraz, J.F.P. (1993) Evidence for *Phytophthora cinnamomi* involvement in Iberian oak decline. *Plant Pathology*, 42, 140–145.
- Brasier, C.M., Sanchez-Hernandez, E. & Kirk, S.A. (2003) *Phytophthora inundata* sp. nov., a part heterothallic pathogen of trees and shrubs in wet or flooded soils. *Mycological Research*, 107, 477–484.
- Brasier, C.M., Scanu, B., Cooke, D. & Jung, T. (2022) *Phytophthora*: an ancient, historic, biologically and structurally cohesive and evolutionarily successful generic concept in need of preservation. *IMA Fungus*, 13, 12.
- Brasier, C.M. & Webber, J. (2010) Sudden larch death. *Nature*, 466, 824–825.
- Bregant, C., Batista, E., Hilário, S., Linaldeddu, B.T. & Alves, A. (2023a) *Phytophthora* species involved in *Alnus glutinosa* decline in Portugal. *Pathogens*, 12, 276.
- Bregant, C., Batista, E., Hilário, S., Linaldeddu, B.T. & Alves, A. (2023b) First report of *Phytophthora hibernalis*, *P. multivora* and *P. niederhauerii* causing root rot and bleeding cankers on *Eucalyptus globulus* in Portugal. *New Disease Reports*, 47, e12171.

- Bregant, C., Sanna, G.P., Bottos, A., Maddau, L., Montecchio, L. & Linaldeddu, B.T. (2020) Diversity and pathogenicity of *Phytophthora* species associated with declining alder trees in Italy and description of *Phytophthora alpina* sp. nov. *Forests*, 11, 848.
- Burgess, T.I. (2015) Molecular characterization of natural hybrids formed between five related indigenous clade 6 *Phytophthora* species. *PLoS One*, 10, e0134225.
- Burgess, T.I., White, D., McDougall, K.M., Garnas, J., Dunstan, W.A., Català, S. et al. (2017) Distribution and diversity of *Phytophthora* across Australia. *Pacific Conservation Biology*, 23, 1–13.
- Caballol, M., Redondo, M.A., Catalán, N., Corcobado, T., Jung, T., Marçais, B. et al. (2024) Climate acts as an environmental filter to plant pathogens. *The ISME Journal*, 18, wrae010.
- Cacciola, S.O., Motta, E., Raudino, F., Chimento, A., Pane, A., di San, M. et al. (2005) *Phytophthora pseudosyringae* the causal agent of bleeding cankers of beech in central Italy. *Journal of Plant Pathology*, 87, 289.
- Català, S., Peréz-Sierra, A. & Abad-Campos, P. (2015) The use of genus-specific amplicon pyrosequencing to assess *Phytophthora* species diversity using eDNA from soil and water in northern Spain. *PLoS One*, 10, e0119311.
- Černý, K., Filipová, N. & Strnadová, V. (2012) Influence of low temperature and frost duration on *Phytophthora alni* subsp. *alni* viability. *Forestry Systems*, 21, 337–342.
- Černý, K. & Strnadová, V. (2010) *Phytophthora* alder decline: disease symptoms, causal agent and its distribution in the Czech Republic. *Plant Protection Science*, 46, 12–18.
- Chen, Q., Bakhshi, M., Balci, Y., Broders, K.D., Cheewangkoon, R., Chen, S.F. et al. (2022) Genera of phytopathogenic fungi: GOPHY 4. *Studies in Mycology*, 101, 417–564.
- Cooke, D.E.L., Drenth, A., Duncan, J.M., Wagels, G. & Brasier, C.M. (2000) A molecular phylogeny of *Phytophthora* and related oomycetes. *Fungal Genetics and Biology*, 30, 17–32.
- Corcobado, T., Cech, T.L., Brandstetter, M., Daxer, A., Hüttler, C., Kudláček, T. et al. (2020) Decline of European beech in Austria: involvement of *Phytophthora* spp. and contributing biotic and abiotic factors. *Forests*, 11, 895.
- Corcobado, T., Cech, T.L., Daxer, A., Ďatková, H., Janoušek, J., Patra, S. et al. (2023) *Phytophthora*, *Nothophytophthora* and *Halophytophthora* diversity in rivers, streams and riparian alder ecosystems of Central Europe. *Mycological Progress*, 22, 50.
- Corcobado, T., Cubera, E., Pérez-Sierra, A., Jung, T. & Solla, A. (2010) First report of *Phytophthora gonapodyides* involved in the decline of *Quercus ilex* in xeric conditions in Spain. *New Disease Reports*, 22, 33.
- Crandall, B.S., Gravatt, G.F. & Ryan, M.M. (1945) Root disease of *Castanea* species and some coniferous and broadleaf nursery stocks, caused by *Phytophthora cinnamomi*. *Phytopathology*, 35, 162–180.
- Crous, P.W., Summerell, B.A., Shivas, R.G., Burgess, T.I., Decock, C.A., Dreyer, L.L. et al. (2012) Fungal planet description sheets: 107–127. *Persoonia*, 28, 138–182.
- Day, W.R. (1938) Root-rot of sweet chestnut and beech caused by species of *Phytophthora*: 1. Cause and symptoms of disease: its relation to soil conditions. *Forestry*, 12, 101–116.
- Diogo, E., Machado, H., Reis, A., Valente, C., Phillips, A.J.L. & Bragança, H. (2023) *Phytophthora alticola* and *Phytophthora cinnamomi* on *Eucalyptus globulus* in Portugal. *European Journal of Plant Pathology*, 165, 255–269.
- Disney, R. (2009) *A history of Portugal and the Portuguese empire, volume 2: the Portuguese empire*. Cambridge, UK: Cambridge University Press.
- Domingues de Almeida, J. & Freitas, H. (2012) Exotic flora of continental Portugal – a new assessment. *Bocconea*, 24, 231–237.
- Drenth, A. & Guest, D.I. (2004) *Diversity and management of Phytophthora in Southeast Asia*. Canberra: Australian Centre for International Agricultural Research, ACIAR Monograph, p. 114.
- Dunstan, W.A., Howard, K., Hardy, G.E.S.J. & Burgess, T.I. (2016) An overview of Australia's *Phytophthora* species assemblage in natural ecosystems recovered from a survey in Victoria. *IMA Fungus*, 7, 47–58.
- Erwin, D.C. & Ribeiro, O.K. (1996) *Phytophthora diseases worldwide*. St Paul, MN: APS Press.
- Frankel, S.J., Conforti, C., Hillman, J., Ingolia, M., Shor, A., Benner, D. et al. (2020) *Phytophthora* introductions in restoration areas: responding to protect California native flora from human-assisted pathogen spread. *Forests*, 11, 1291.
- Gallego, F.J., Perez de Algaba, A. & Fernandez-Escobar, R. (1999) Etiology of oak decline in Spain. *European Journal of Forest Pathology*, 29, 17–27.
- Garbelotto, M.M., Frankel, S. & Scanu, B. (2018) Soil- and waterborne *Phytophthora* species linked to recent outbreaks in northern California restoration sites. *California Agriculture*, 72, 208–216.
- Ginetti, B., Moricca, S., Squires, J.N., Ragazzi, A. & Jung, T. (2014) *Phytophthora acerina* sp. nov., a new species causing bleeding cankers and dieback of *Acer pseudoplatanus* trees in planted forests in northern Italy. *Plant Pathology*, 63, 858–876.
- Gomes, M.J. & Amaro, P.A. (2008) Ocorrência de *Phytophthora ramorum* em Portugal sobre *Viburnum* spp. *Revista de Ciências Agrárias*, 31, 105–111.
- Gomes Marques, I., Vieites-Blanco, C., Rodríguez-González, P.M., Segurado, P., Marques, M., Barrento, M.J. et al. (2024) The ADnet Bayesian belief network for alder decline: integrating empirical data and expert knowledge. *Science of the Total Environment*, 947, 173619.
- Greslebin, A.G., Hansen, E.M. & Sutton, W. (2007) *Phytophthora austrocedrae* sp. nov., a new species associated with *Austrocedrus chilensis* mortality in Patagonia Argentina. *Mycological Research*, 111, 308–316.
- Grünwald, N.J., Garbelotto, M., Goss, E.M., Heungens, K. & Prospero, S. (2012) Emergence of the sudden oak death pathogen *Phytophthora ramorum*. *Trends in Microbiology*, 20, 131–138.
- Hansen, E.M. & Delatour, C. (1999) *Phytophthora* species in oak forests of northeast France. *Annales Des Sciences Forestière*, 56, 539–547.
- Hansen, E.M., Goheen, D.J., Jules, E.S. & Ullian, B. (2000) Managing port-Orford-cedar and the introduced pathogen *Phytophthora lateralis*. *Plant Disease*, 84, 4–14.
- Haque, M.M.U., Hidalgo, E., Martín-García, J., De-Lucas, A.I. & Diez, J.J. (2015) Morphological, physiological and molecular characterization of *Phytophthora alni* isolates from Western Spain. *European Journal of Plant Pathology*, 142, 731–745.
- Hardham, A.R. & Blackman, L.M. (2018) *Phytophthora cinnamomi*. *Molecular Plant Pathology*, 19, 260–285.
- Hong, C.X., Gallegly, M.E., Richardson, P.A., Kong, P., Moorman, G.W., Lea-Cox, J.D. et al. (2010) *Phytophthora hydropathica*, a new pathogen identified from irrigation water, *Rhododendron catawbiense* and *Kalmia latifolia*. *Plant Pathology*, 59, 913–921.
- Huai, W.X., Tian, G., Hansen, E.M., Zhao, W.X., Goheen, E.M., Grünwald, N.J. et al. (2013) Identification of *Phytophthora* species baited and isolated from forest soil and streams in northwestern Yunnan province, China. *Forest Pathology*, 43, 87–103.
- Hüberli, D., Hardy, G.E.S.J., White, D., Williams, N. & Burgess, T.I. (2013) Fishing for *Phytophthora* from Western Australia's waterways: a distribution and diversity survey. *Australasian Plant Pathology*, 42, 251–260.
- Husson, C., Aguayo, J., Revellin, C., Frey, P., Ios, R. & Marçais, B. (2015) Evidence for homoploid speciation in *Phytophthora alni* supports taxonomic reclassification in this species complex. *Fungal Genetics and Biology*, 77, 12–21.
- ICNF. (2019) 6^o Inventário Florestal Nacional (IFN6) – 2015. *Relatório Final*. Lisbon: Instituto da Conservação da Natureza e das Florestas (ICNF).
- Jayasekera, A.U., McComb, J.A., Shearer, B.L. & Hardy, G.E.S.J. (2007) *In planta* selfing and oospore production of *Phytophthora cinnamomi*

- in the presence of *Acacia pulchella*. *Mycological Research*, 111, 355–362.
- Jönsson, U., Lundberg, L., Sonesson, K. & Jung, T. (2003) First records of soilborne *Phytophthora* species in Swedish oak forests. *Forest Pathology*, 33, 175–179.
- Jung, T. (2009) Beech decline in Central Europe driven by the interaction between *Phytophthora* infections and climatic extremes. *Forest Pathology*, 39, 73–94.
- Jung, T., Blaschke, H. & Neumann, P. (1996) Isolation, identification and pathogenicity of *Phytophthora* species from declining oak stands. *European Journal of Forest Pathology*, 26, 253–272.
- Jung, T., Blaschke, H. & Osswald, W. (2000) Involvement of soilborne *Phytophthora* species in central European oak decline and the effect of site factors on the disease. *Plant Pathology*, 49, 706–718.
- Jung, T. & Blaschke, M. (2004) *Phytophthora* root and collar rot of alders in Bavaria: distribution, modes of spread and possible management strategies. *Plant Pathology*, 53, 197–208.
- Jung, T. & Burgess, T.I. (2009) Re-evaluation of *Phytophthora citricola* isolates from multiple woody hosts in Europe and North America reveals a new species, *Phytophthora plurivora* sp. nov. *Persoonia*, 22, 95–110.
- Jung, T., Chang, T.T., Bakonyi, J., Seress, D., Pérez-Sierra, A., Yang, X. et al. (2017) Diversity of *Phytophthora* species in natural ecosystems of Taiwan and association with disease symptoms. *Plant Pathology*, 66, 194–211.
- Jung, T., Colquhoun, I.J. & Hardy, G.E.S.J. (2013) New insights into the survival strategy of the invasive soilborne pathogen *Phytophthora cinnamomi* in different natural ecosystems in Western Australia. *Forest Pathology*, 43, 266–288.
- Jung, T., Cooke, D.E.L., Blaschke, H., Duncan, J.M. & Oßwald, W. (1999) *Phytophthora quercina* sp. nov., causing root rot of European oaks. *Mycological Research*, 103, 785–798.
- Jung, T., Durán, A., Sanfuentes von Stowasser, E., Schena, L., Mosca, S., Fajardo, S. et al. (2018) Diversity of *Phytophthora* species in Valdivian rainforests and association with severe dieback symptoms. *Forest Pathology*, 48, e12443.
- Jung, T., Hansen, E.M., Winton, L., Oßwald, W. & Delatour, C. (2002) Three new species of *Phytophthora* from European oak forests. *Mycological Research*, 106, 397–411.
- Jung, T., Horta Jung, M., Cacciola, S.O., Cech, T., Bakonyi, J., Seress, D. et al. (2017) Multiple new cryptic pathogenic *Phytophthora* species from *Fagaceae* forests in Austria, Italy and Portugal. *IMA Fungus*, 8, 219–244.
- Jung, T., Horta Jung, M., Scanu, B., Seress, D., Kovács, G.M., Maia, C. et al. (2017) Six new *Phytophthora* species from ITS clade 7a including two sexually functional heterothallic hybrid species detected in natural ecosystems in Taiwan. *Persoonia*, 38, 100–135.
- Jung, T., Horta Jung, M., Webber, J.F., Kageyama, K., Hieno, A., Masuya, H. et al. (2021) The destructive tree pathogen *Phytophthora ramorum* originates from the Laurosilva forests of East Asia. *Journal of Fungi*, 7, 226.
- Jung, T., La Spada, F., Pane, A., Aloï, F., Evoli, M., Horta Jung, M. et al. (2019) Diversity and distribution of *Phytophthora* species in protected natural areas in Sicily. *Forests*, 10, 259.
- Jung, T., Milenković, I., Balci, Y., Janoušek, J., Kudláček, T., Nagy, Z.A. et al. (2024) Worldwide forest surveys reveal forty-three new species in *Phytophthora* major clade 2 with fundamental implications for the evolution and biogeography of the genus and global plant biosecurity. *Studies in Mycology*, 107, 251–388.
- Jung, T., Milenković, I., Corcobado, T., Májek, T., Janoušek, J., Kudláček, T. et al. (2022) Extensive morphological and behavioural diversity among fourteen new and seven described species in *Phytophthora* clade 10 and its evolutionary implications. *Persoonia*, 49, 1–57.
- Jung, T. & Nechwatal, J. (2008) *Phytophthora gallica* sp. nov., a new species from rhizosphere soil of declining oak and reed stands in France and Germany. *Mycological Research*, 112, 1195–1205.
- Jung, T., Nechwatal, J., Cooke, D.E.L., Hartmann, G., Blaschke, M., Oßwald, W.F. et al. (2003) *Phytophthora pseudosyringae* sp. nov., a new species causing root and collar rot of deciduous tree species in Europe. *Mycological Research*, 107, 772–789.
- Jung, T., Orlikowski, L., Henricot, B., Abad-Campos, P., Aday, A.G., Aguín Casal, O. et al. (2016) Widespread *Phytophthora* infestations in European nurseries put forest, semi-natural and horticultural ecosystems at high risk of *Phytophthora* diseases. *Forest Pathology*, 46, 134–163.
- Jung, T., Pérez-Sierra, A., Durán, A., Horta Jung, M., Balci, Y. & Scanu, B. (2018) Canker and decline diseases caused by soil- and airborne *Phytophthora* species in forests and woodlands. *Persoonia*, 40, 182–220.
- Jung, T., Scanu, B., Bakonyi, J., Seress, D., Kovács, G.M., Durán, A. et al. (2017) *Nothophytophthora* gen. nov., a new sister genus of *Phytophthora* from natural and semi-natural ecosystems. *Persoonia*, 39, 143–174.
- Jung, T., Scanu, B., Brasier, C.M., Webber, J., Milenković, I., Corcobado, T. et al. (2020) A survey in natural forest ecosystems of Vietnam reveals high diversity of both new and described *Phytophthora* taxa including *P. ramorum*. *Forests*, 11, 93.
- Jung, T., Stukely, M.J.C., Hardy, G.E.S.J., White, D., Paap, T., Dunstan, W.A. et al. (2011) Multiple new *Phytophthora* species from ITS clade 6 associated with natural ecosystems in Australia: evolutionary and ecological implications. *Persoonia*, 26, 13–39.
- Kanoun-Boulé, M., Vasconcelos, T., Gaspar, J., Viera, S., Dias-Ferreira, C. & Husson, C. (2016) *Phytophthora xalni* and *Phytophthora lacustris* associated with common alder decline in Central Portugal. *Forest Pathology*, 46, 174–176.
- Kroon, L.P.N.M., Bakker, F.T., van den Bosch, G.B.M., Bonants, P.J.M. & Flier, W.G. (2004) Phylogenetic analysis of *Phytophthora* species based on mitochondrial and nuclear DNA sequences. *Fungal Genetics and Biology*, 41, 766–782.
- Lamour, K. (2013) *Phytophthora: a global perspective*. Wallingford, UK: CABI.
- Leonberger, A.J., Speers, C., Ruhl, G., Creswell, T. & Beckerman, J.L. (2013) A survey of *Phytophthora* spp. in Midwest nurseries, greenhouses, and landscapes. *Plant Disease*, 97, 635–640.
- Liebold, A.M., Brockerhoff, E.G., Garrett, L.J., Parke, J.L. & Britton, K.O. (2012) Live plant imports: the major pathway for forest insect and pathogen invasions of the US. *Frontiers in Ecology and the Environment*, 10, 135–143.
- Lopes-Pimentel, A.A. (1946) O sobreiro também é parasitado pela *Phytophthora cambivora* (Petri) Buis., agente da “doença da tinta” do castanheiro. Lisboa. *Direcção Geral dos Serviços Florestais e Aquícolas*, 13, 45–49.
- Lopes-Pimentel, A.A. (1947) *Phytophthora cinnamomi* (Rands) um outro agente extremamente virulento da “doença da tinta” do castanheiro. *Agronomia Lusitana*, 9, 181–191.
- Macháčová, M., Tomášková, I., Corcobado, T., Nagy, Z., Milanović, S., Janoušek, J. et al. (2024) Response of *Alnus glutinosa* to *Phytophthora* bark infections at ambient and elevated CO₂ levels. *Frontiers in Forests and Global Change*, 7, 1379791.
- Martin, F.N. & Tooley, P.W. (2003) Phylogenetic relationships among *Phytophthora* species inferred from sequence analysis of mitochondrially encoded cytochrome oxidase I and II genes. *Mycologia*, 95, 269–284.
- Martins, L., Castro, J., Macedo, W., Marques, C. & Abreu, C. (2007) Assessment of the spread of chestnut ink disease using remote sensing and geostatistical methods. *European Journal of Plant Pathology*, 119, 159–164.
- Maseko, B., Coutinho, T.A., Burgess, T.I., Wingfield, B.D. & Wingfield, M.J. (2007) Two new species of *Phytophthora* from south African eucalypt plantations. *Mycological Research*, 111, 1321–1338.
- Migliorini, D., Ghelardini, L., Tondine, E., Luchi, N. & Santini, A. (2015) The potential of symptomless potted plants for carrying

- invasive soilborne plant pathogens. *Diversity and Distributions*, 21, 1218–1229.
- Milenković, I., Keča, N., Karadžić, D., Nowakowska, J.A., Borys, M., Sikora, K. et al. (2012) Incidence of *Phytophthora* species in beech stands in Serbia. *Folia Forestalia Polonica, Series A*, 54, 223–232.
- Moralejo, E., Pérez-Sierra, A., Alvarez, L.A., Belbahri, L., Lefort, F. & Descals, E. (2009) Multiple alien *Phytophthora* taxa discovered on diseased ornamental plants in Spain. *Plant Pathology*, 58, 100–110.
- Mora-Sala, B., León, M., Pérez-Sierra, A. & Abad-Campos, P. (2022) New reports of *Phytophthora* species in plant nurseries in Spain. *Pathogens*, 11, 826.
- Moreira, A.C. & Martins, J.M.S. (2005) Influence of site factors on the impact of *Phytophthora cinnamomi* in cork oak stands in Portugal. *Forest Pathology*, 35, 145–162.
- Mullett, M.S., Harris, A.R., Scanu, B., Van Poucke, K., LeBoldus, J., Stamm, E. et al. (2024) Phylogeography, origin and population structure of the self-fertile emerging plant pathogen *Phytophthora pseudosyringae*. *Molecular Plant Pathology*, 25, e13450.
- Mullett, M.S., Van Poucke, K., Haegemann, A., Focquet, F., Cauldron, N.C., Knaus, B.J. et al. (2023) Phylogeography and population structure of the global, wide host-range hybrid pathogen *Phytophthora x cambivora*. *IMA Fungus*, 14, 4.
- Nagel, J.H., Gryzenhout, M., Slippers, B., Wingfield, M.J., Hardy, G.E.S.J., Stukely, M.J.C. et al. (2013) Characterization of *Phytophthora* hybrids from ITS clade 6 associated with riparian ecosystems in South Africa and Australia. *Fungal Biology*, 117, 329–347.
- Nagy, Z.Á., Bakonyi, J. & Érsek, T. (2003) Standard and Swedish variant types of the hybrid alder *Phytophthora* attacking alder in Hungary. *Pest Management Science*, 59, 484–492.
- Oh, E., Gryzenhout, M., Wingfield, B.D., Wingfield, M.J. & Burgess, T.I. (2013) Surveys of soil and water reveal a goldmine of *Phytophthora* diversity in south African natural ecosystems. *IMA Fungus*, 4, 123–131.
- Page, M. (2013) *The first global village: how Portugal changed the world*, 18th edition. Alfragide, Portugal: Casa das Letras.
- Parke, J.L. & Grünwald, N.J. (2012) A systems approach for management of pests and pathogens of nursery crops. *Plant Disease*, 96, 1236–1244.
- Pérez-Sierra, A., Horta Jung, M. & Jung, T. (2022) Survey and monitoring of *Phytophthora* species in natural ecosystems: methods for sampling, isolation, purification, storage, and pathogenicity tests. *Methods in Molecular Biology*, 2536, 23–49.
- Pérez-Sierra, A. & Jung, T. (2013) *Phytophthora* in woody ornamental nurseries. In: Lamour, K. (Ed.) *Phytophthora: a global perspective*. Wallingford, UK: CABI, pp. 166–177.
- Pérez-Sierra, A., López-García, C., León, M., García-Jiménez, J., Abad-Campos, P. & Jung, T. (2013) Previously unrecorded low temperature *Phytophthora* species associated with *Quercus* decline in a Mediterranean forest in eastern Spain. *Forest Pathology*, 43, 331–339.
- Peters, F.S., Wunderlich, F. & Metzler, B. (2019) First report of *Phytophthora cinnamomi* in forest stands in Germany. *Forest Pathology*, 49, e12485.
- Prigigallo, M.I., Abdelfattah, A., Cacciola, S.O., Faedda, R., Sanzani, S.M., Cooke, D.E.L. et al. (2016) Metabarcoding analysis of *Phytophthora* diversity using genus-specific primers and 454 pyrosequencing. *Phytopathology*, 106, 305–313.
- Prigigallo, M.I., Mosca, S., Cacciola, S.O., Cooke, D.E.L. & Schena, L. (2015) Molecular analysis of *Phytophthora* diversity in nursery-grown ornamental and fruit plants. *Plant Pathology*, 64, 1308–1319.
- Rea, A.J., Burgess, T.I., Hardy, G.E.St.J., Stukely, M.J.C. & Jung, T. (2011) Two novel and potentially endemic species of *Phytophthora* associated with episodic dieback of kwongan vegetation in the southwest of Western Australia. *Plant Pathology*, 60, 1055–1068.
- Redondo, M.A., Boberg, J., Olsson, C.H.B. & Oliva, J. (2015) Winter conditions correlate with *Phytophthora alni* subspecies distribution in southern Sweden. *Phytopathology*, 105, 1191–1197.
- Reeser, P.W., Sutton, W., Hansen, E.M., Remigi, P. & Adams, G.C. (2011) *Phytophthora* species in forest streams in Oregon and Alaska. *Mycologia*, 103, 22–35.
- Riolo, M., Aloï, F., La Spada, F., Sciandrello, S., Moricca, S., Santilli, E. et al. (2020) Diversity of *Phytophthora* communities across different types of Mediterranean vegetation in a nature reserve area. *Forests*, 11, 853.
- Robideau, G.P., De Cock, A.W., Coffey, M.D., Voglmayr, H., Brouwer, H., Bala, K. et al. (2011) DNA barcoding of oomycetes with cytochrome c oxidase subunit I and internal transcribed spacer. *Molecular Ecology Resources*, 11, 1002–1011.
- Robin, C., Desprez-Loustau, M.L., Capron, G. & Delatour, C. (1998) First record of *Phytophthora cinnamomi* on cork and holm oaks in France and evidence of pathogenicity. *Annales des Sciences Forestières*, 55, 869–883.
- Robin, C., Piou, D., Feau, N.F., Douzon, G., Schenck, N., Hansen, E.M. et al. (2011) Root and aerial infections of *Chamaecyparis lawsoniana* by *Phytophthora lateralis*: a new threat for European countries. *Forest Pathology*, 41, 417–424.
- Rooney-Latham, S., Blomquist, C.L., Kosta, K.L., Gou, Y.Y. & Woods, P.W. (2019) *Phytophthora* species are common on nursery stock grown for restoration and revegetation purposes in California. *Plant Disease*, 103, 448–455.
- Sanchez, M.E., Caetano, P., Ferraz, J. & Trapero, A. (2002) *Phytophthora* disease of *Quercus ilex* in southwestern Spain. *Forest Pathology*, 32, 5–18.
- Santini, A., Ghelardini, L., De Pace, C., Desprez-Loustau, M.L., Capretti, P., Chandelier, A. et al. (2013) Biogeographic patterns and determinants of invasion by alien forest pathogens in Europe. *New Phytologist*, 197, 238–250.
- Scanu, B., Hunter, G.C., Linaldeddu, B.T., Franceschini, A., Maddau, L., Jung, T. et al. (2014) A taxonomic re-evaluation reveals that *Phytophthora cinnamomi* and *P. cinnamomi* var. *parvispora* are separate species. *Forest Pathology*, 44, 1–20.
- Scanu, B., Linaldeddu, B.T., Deidda, A. & Jung, T. (2015) Diversity of *Phytophthora* species from declining Mediterranean maquis vegetation, including two new species, *Phytophthora crassamura* and *P. ornamentata* sp. nov. *PLoS One*, 10, e0143234.
- Scanu, B., Linaldeddu, B.T., Pérez-Sierra, A., Deidda, A. & Franceschini, A. (2014) *Phytophthora ilicis* as a leaf and stem pathogen of *Ilex aquifolium* in Mediterranean islands. *Phytopathologia Mediterranea*, 53, 480–490.
- Schenck, N., Saurat, C., Guinet, C., Fourrier-Jeandel, C., Roche, L., Bouvet, A. et al. (2018) First report of *Phytophthora ramorum* causing Japanese larch dieback in France. *Plant Disease*, 102, 2045.
- Schumacher, J., Leonhard, S., Grundmann, B.M. & Roloff, A. (2006) New alder disease in Spreewald biosphere reserve – causes and incidental factors of an epidemic. *Nachrichtenblatt Deutscher Pflanzenschutzdienst*, 58, 141–147.
- Scott, P.M., Burgess, T.I., Barber, P.A., Shearer, B.L., Stukely, M.J.C., Hardy, G.E.S.J. et al. (2009) *Phytophthora multivora* sp. nov., a new species recovered from declining *Eucalyptus*, *Banksia*, *Agonis* and other plant species in Western Australia. *Persoonia*, 22, 1–13.
- Seddaiu, S., Brandano, A., Ruiu, P.A., Sechi, C. & Scanu, B. (2020) An overview of *Phytophthora* species inhabiting declining *Quercus suber* stands in Sardinia (Italy). *Forests*, 11, 971.
- Shakya, S.K., Grünwald, N.J., Fieland, V.J., Knaus, B.J., Weiland, J.E., Maia, C. et al. (2021) Phylogeography of the wide-host range panglobal plant pathogen *Phytophthora cinnamomi*. *Molecular Ecology*, 30, 5164–5178.
- Shrestha, S.K., Zhou, Y. & Lamour, K. (2013) Oomycetes baited from streams in Tennessee 2010–2012. *Mycologia*, 105, 1516–1523.

- Simamora, A.V., Stukely, M.J.C., Hardy, G.E.S.J. & Burgess, T.I. (2015) *Phytophthora boodjera* sp. nov., a damping-off pathogen in production nurseries and from urban and natural landscapes, with an update on the status of *P. alticola*. *IMA Fungus*, 6, 319–335.
- Sims, L.L. & Garbelotto, M. (2021) *Phytophthora* species repeatedly introduced in northern California through restoration projects can spread into adjacent sites. *Biological Invasions*, 23, 2173–2190.
- Sims, L.L., Sutton, W., Reeser, P. & Hansen, E.M. (2015) The *Phytophthora* species assemblage and diversity in riparian alder ecosystems of western Oregon, USA. *Mycologia*, 107, 889–902.
- Solla, A., Pérez-Sierra, A., Corcobado, T., Haque, M.M., Diez, J.J. & Jung, T. (2010) *Phytophthora alni* on *Alnus glutinosa* reported for the first time in Spain. *Plant Pathology*, 59, 798.
- Stępniewska, H. & Dłuszyński, J. (2010) Incidence of *Phytophthora cambivora* in bleeding lesions on beech stems in selected forest stands in south-eastern Poland. *Phytopathologia*, 56, 39–51.
- Stewart-Wade, S.M. (2011) Plant pathogens in recycled irrigation water in commercial plant nurseries and greenhouses: their detection and management. *Irrigation Science*, 29, 267–297.
- Štraus, D., Caballol, M., Serradó, F., Oliveras, J., Ramis, X. & Oliva, J. (2023) Distribution of *Phytophthora* species within recreational chestnut, beech and cork oak forests. *Forest Ecology and Management*, 529, 120674.
- Thoirain, B., Husson, C. & Marçais, B. (2007) Risk factors for the *Phytophthora*-induced decline of alder in north-eastern France. *Phytopathology*, 97, 99–105.
- Tkaczyk, M., Sikora, K., Galko, J. & Kunca, A. (2023) Occurrence of *Phytophthora* species in riparian stands of black alder (*Alnus glutinosa*) in Slovakia. *Forest Pathology*, 53, e12800.
- Trzewik, A., Orlikowski, L.B., Oszako, T., Nowakowska, J.A. & Orlikowska, T. (2015) The characterization of *Phytophthora* isolates obtained from diseased *Alnus glutinosa* in Poland. *Baltic Forestry*, 21, 44–50.
- Tsykun, T., Prospero, S., Schoebel, C.N., Rea, A. & Burgess, T.I. (2022) Global invasion history of the emerging plant pathogen *Phytophthora multivora*. *BMC Genomics*, 23, 153.
- Ufer, T., Werres, S., Posner, M. & Wessels, H.-P. (2008) Filtration to eliminate *Phytophthora* spp. from recirculating water systems in commercial nurseries. *Plant Health Progress*, 9. Available from: <https://doi.org/10.1094/PHP-2008-0314-01-RS>
- Van Poucke, K., Haegeman, A., Goedefroit, T., Focquet, F., Leus, L., Horta Jung, M. et al. (2021) Unravelling hybridization in *Phytophthora* using phylogenomics and genome size estimation. *IMA Fungus*, 12, 16.
- Vettrano, A.M., Barzanti, G.P., Bianco, M.C., Ragazzi, A., Capretti, P., Paoletti, E. et al. (2002) Occurrence of *Phytophthora* species in oak stands in Italy and their association with declining oak trees. *Forest Pathology*, 32, 19–28.
- Vettrano, A.M., Morel, O., Perlerou, C., Robin, C., Diamandis, S. & Vannini, A. (2005) Occurrence and distribution of *Phytophthora* species associated with ink disease of chestnut in Europe. *European Journal of Plant Pathology*, 111, 169–180.
- White, T.J., Bruns, T., Lee, S. & Taylor, J. (1990) Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: Innes, M.A., Gelfand, D.H., Sninsky, J.J. & White, T.J. (Eds.) *PCR protocols: a guide to methods and applications*. San Diego, CA, USA: Academic Press, pp. 315–322.
- Yang, X., Richardson, P.A. & Hong, C. (2014) *Phytophthora xstagnum* nothosp. nov., a new hybrid from irrigation reservoirs at ornamental plant nurseries in Virginia. *PLoS One*, 9, e103450.
- Yang, X., Tyler, B.M. & Hong, C. (2017) An expanded phylogeny for the genus *Phytophthora*. *IMA Fungus*, 8, 355–384.

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

How to cite this article: Horta Jung, M., Maia, C., Mora-Sala, B., Abad-Campos, P., Schena, L., Mosca, S. et al. (2024) High diversity of *Phytophthora* species in natural ecosystems and nurseries of Portugal: Detrimental side effect of plant introductions from the age of discovery to modern globalization. *Plant Pathology*, 00, 1–33. Available from: <https://doi.org/10.1111/ppa.14022>