

ECSA 41ST INTERNATIONAL CONFERENCE
MEASURING AND MANAGING
CHANGES IN ESTUARIES AND LAGOONS

CONTRIBUTION INFORMATION

AUTHORS: Caeiro, S.; Costa, M. H.; Goovaerts, P.; Martins, F.

TITLE: BENTHIC BIOTOPE INDEX FOR CLASSIFYING ESTUARINE HABITATS. A CASE STUDY

(Please tick the appropriate box)

Oral presentation

Poster

Corresponding author:

Name: Sandra Caeiro

Address: IMAR, Department of Exact and Technological Sciences of the Portuguese Distance Learning University, R. Escola Politecnica, 147, 1269-001 Lisbon, Portugal.

E-mail: scaeiro@univ-ab.pt

BENTHIC BIOTOPE INDEX FOR CLASSIFYING ESTUARINE HABITATS. A CASE STUDY¹

CAEIRO, S.^{1*}; COSTA, M. H.²; GOOVAERTS, P.³; MARTINS, F.⁴

1. IMAR, Department of Exact and Technological Sciences of the Portuguese Distance Learning University, R. Escola Politecnica, 147, 1269-001 Lisbon, Portugal.

2. IMAR, Faculty of Science and Technology of the New University of Lisbon, Quinta da Torre 2829-516 Caparica, Portugal,

3. BioMedware, Inc. 516 North State Street, Ann Arbor, USA;

4. EST, Universidade do Algarve, Campus da Penha, 8000 Faro;

* scaeiro@univ-ab.pt

An integration of sediment physical, chemical, biological, and toxicity data is necessary for a meaningful interpretation of the complex sediment conditions in the marine environment. Assessment of benthic community is a vital component for that interpretation, yet their evaluation is complex and requires a large expenditure of time and funds. Thus, there is a need for new tools that are less expensive and more understandable for managers. This paper presents a benthic biotope index to predict from physical and chemical variables the occurrence of macrobenthic habitats, applied to Sado Estuary, as a case of study. This estuary is the second largest in Portugal and most of it is classified as a natural reserve, but with many industries, harbor associated activities and with agriculture, aquaculture and tourism pressures. Parameters such as sediment type, organic matter, depth, and hydrodynamic parameters were selected, through a forward stepwise discriminant analysis, to compute the index. A dataset of 131 stations were used. Other authors have used multivariate methods to determine the benthic biotopes to Sado Estuary. These biotopes have a very good agreement between the conclusions drawn from the analysis of the biological data alone and those from the imposed variability of the physical and chemical variables listed above. The model was tested and validated using field macrobenthic data. The index permits to predict the occurrence of benthic biotopes at unsampled locations in a cost-effective way and proved to be a valid tool to classify and assess the spatial patterns of benthic habitat and to identify stress biotope gradients.

¹ Most part of this work has been published in *Marine Environmental Research* 60(2005) 570 – 593.