




## Correction to: Characterization of a fatty acid-binding protein from the Pacific oyster (*Crassostrea gigas*): pharmaceutical and toxicological implications

Juliana F. Tisca<sup>1,2</sup> · Karin dos Santos<sup>2,3</sup> · Tomás B. Pessati<sup>1,2</sup> · Flávia L. Zacchi<sup>1,2</sup> · Fabíola S. Soares<sup>2</sup> · Vanessa A. Oliveira<sup>4</sup> · Maria J. A. F. Bebianno<sup>5</sup> · Afonso C. D. Bainy<sup>1,2</sup> · Guilherme Razzera<sup>1,2,3</sup> 

© Springer-Verlag GmbH Germany, part of Springer Nature 2021

### Correction to: Environmental Science and Pollution Research

<https://doi.org/10.1007/s11356-021-12645-y>

The correct Fig. 3 is presented in this paper.  
The original article has been corrected.

---


The online version of the original article can be found at <https://doi.org/10.1007/s11356-021-12645-y>

---

✉ Afonso C. D. Bainy  
afonso.bainy@ufsc.br

✉ Guilherme Razzera  
guilherme.razzera@ufsc.br

- <sup>1</sup> Programa de Pós-Graduação em Bioquímica, Departamento de Bioquímica, Universidade Federal de Santa Catarina, Florianópolis, SC 88040-900, Brazil
- <sup>2</sup> Laboratório de Biomarcadores de Contaminação Aquática e Imunoquímica, Universidade Federal de Santa Catarina, Florianópolis, SC 88040-900, Brazil
- <sup>3</sup> Núcleo de Bioinformática e Biologia Computacional da UFSC-NuBioinfo, Universidade Federal de Santa Catarina, Florianópolis, SC 88040-900, Brazil
- <sup>4</sup> Centro de Biologia Molecular Estrutural, Universidade Federal de Santa Catarina, Florianópolis, SC 88040-900, Brazil
- <sup>5</sup> Centre for Marine and Environmental Research, CIMA, University of Algarve, Campus de Gambelas, 8000-139 Faro, Portugal



**Fig. 3** Circular dichroism spectra of rCgFABP2 (5.0  $\mu\text{M}$ ) in phosphate buffer (0.75 mM pH 7.4) in the presence of increasing amounts of fatty acids **(a)** Palmitic acid; **(b)** Oleic acid; and **(c)** Arachidonic acid. Titration stoichiometry ratios:  $r = [\text{ligand}]/[\text{CgFABP2}] = 0.00, 0.20, 1.00, 2.60,$

5.00, 7.41, 9.81, 12.21, 16.21 and 20.22. **(d)** The  $\alpha$ -helical secondary structural content at 195 nm as a function of ligand concentration. Halfmaximum values ( $\text{EC}_{50}$ ) are shown under the curves

**Publisher's note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.