


Livro de resumos das comunicações do  
Libro de resúmenes de comunicaciones del

# 4º Encontro Ibérico de Ecologia 4º Congreso Ibérico de Ecología

Coimbra, Portugal • 16 a 19 Junho de 2015 / 16 a 19 de Junio de 2015



## A Ecologia e os Desafios Societais La Ecología y los Retos Sociales



Sociedade Portuguesa de Ecologia



ASOCIACIÓN ESPAÑOLA  
DE ECOLOGÍA TERRESTRE



Centre for  
Functional  
Ecology



MARE



• U C •

FCTUC FACULDADE DE CIÊNCIAS  
E TECNOLOGIA  
UNIVERSIDADE DE COIMBRA  
DEPARTAMENTO DE  
CIÊNCIAS DA VIDA



CÂMARA MUNICIPAL  
COIMBRA

Livro de resumos das comunicações do 4º Encontro Ibérico de Ecologia,  
Coimbra, Portugal, 16-19 de Junho de 2015

Libro de resúmenes de comunicaciones del 4º Congreso Ibérico de Ecología,  
Coimbra, Portugal, 16-19 junio de 2015

Publicação eletrónica publicada em Open Access de acordó com Creative Commons  
Attribution Non Commercial Licence

Publicación electrónica publicada en Open Access bajo los términos de Creative  
Commons attribution Non Comercial License

DOI: 10.7818/4IberianEcologicalCongress.2015

Deadline editorial: 01/06/2015

Publica:

Asociación Española de Ecología Terrestre (AEET)  
C/ Tulipán s/n, 28399, Móstoles, Madrid, Spain

Elaboração/Maquetación: Susana Rodríguez Echeverría, João Loureiro, Sílvia Castro.

### S5. Oral

#### **Large-scale prediction of seagrass distribution: the case of *Cymodocea nodosa* (Mediterranean-Atlantic)**

Chefaoui, R.<sup>1</sup>, Assis, J.<sup>1</sup>, Duarte, C.M.<sup>2</sup>, Serrão, E.A.<sup>1</sup>

(1) Centro de Ciências do Mar (CCMAR), Universidade do Algarve, (2) Red Sea Research Center (RSRC), King Abdullah University of Science.

Understanding the factors that affect seagrass meadows encompassing their entire range of distribution is challenging yet important for their conservation. We model the environmental niche of *Cymodocea nodosa* using a combination of environmental variables and landscape metrics to examine factors defining its distribution and find suitable habitats for the species. The most relevant environmental variables defining the distribution of *C. nodosa* were sea surface temperature (SST) and salinity. We found suitable habitats at SST from 5.8 °C to 26.4 °C and salinity ranging from 17.5 to 39.3. Optimal values of mean winter wave height ranged between 1.2 m and 1.5 m, while waves higher than 2.5 m seemed to limit the presence of the species. The influence of nutrients and pH, despite having weight on the models, was not so clear in terms of ranges that confine the distribution of the species. Landscape metrics able to capture variation in the coastline enhanced significantly the accuracy of the models, despite the limitations caused by the scale of the study. By contrasting predictive approaches, we defined the variables affecting the distributional areas that seem unsuitable for *C. nodosa* as well as those suitable habitats not occupied by the species. These findings are encouraging for its use in future studies on climate-related marine range shifts and meadow restoration projects of these fragile ecosystems.