



## Corrigendum to “Innovative approach in sustainable agriculture: Harnessing microalgae potential via subcritical water extraction” [Environ. Technol. Innov. 36, 103797]

Alice Ferreira <sup>a,b</sup>, Jelena Vladic <sup>c</sup>, Diego de Oliveira Corrêa <sup>a,d</sup>,  
Valéria Louzada Leal Butzke <sup>e</sup>, Pedro L. Martins <sup>a,b</sup>, Belina Ribeiro <sup>a</sup>,  
Cláudia Marques-dos-Santos <sup>b</sup>, F. Gabriel Acién <sup>f</sup>, Luisa Gouveia <sup>a,g</sup>

<sup>a</sup> LNEG, National Laboratory of Energy and Geology I.P., Bioenergy and Biorefineries Unit, Estrada do Paço do Lumiar 22, Lisbon 1649-038, Portugal

<sup>b</sup> Forest Research Centre, Associate Laboratory TERRA, School of Agriculture, University of Lisbon, Tapada da Ajuda, Lisboa 1349-017, Portugal

<sup>c</sup> NOVA School of Science and Technology, NOVA University Lisbon, Caparica 2829-516, Portugal

<sup>d</sup> Departamento de Bioquímica e Biologia Molecular, Universidade Federal do Paraná (UFPR), PO Box 19046, Curitiba, Paraná CEP 81531-980, Brazil

<sup>e</sup> Postgraduate Program in Environmental Technology, University of Santa Cruz do Sul (UNISC), Santa Cruz do Sul, Rio Grande do Sul, Brazil

<sup>f</sup> Chemical Engineering Department, University of Almeria, Ctra. Sacramento, s/n, Almería 04120, Spain

<sup>g</sup> GreenCoLab, Green Ocean Technologies and Products Collaborative Laboratory, University of Algarve, Campus de Gambelas, Faro 8005-139, Portugal

The authors regret that the values of **Table 6** are in the incorrect units. The corrected version of **Table 6** is enclosed herewith. Furthermore, the author Jelena Vladic would like to add the DOI to her CEEC grant.

### Funding

Jelena Vladic would like to acknowledge her CEEC grant CEECIND/04909/2022 awarded by Fundação para a Ciência e Tecnologia at program CEEC IND5ed (<https://doi.org/10.54499/2022.04909.CEECIND/CP1725/CT0014>).

The authors would like to apologise for any inconvenience caused.

DOI of original article: <https://doi.org/10.1016/j.eti.2024.103797>.

E-mail address: [alice.ferreira@lneg.pt](mailto:alice.ferreira@lneg.pt) (A. Ferreira).

<https://doi.org/10.1016/j.eti.2024.103934>

Available online 18 December 2024

2352-1864/© 2024 The Author(s).

Published by Elsevier B.V. This is an open access article under the CC BY license

(<http://creativecommons.org/licenses/by/4.0/>).

**Table 6**

Macro and micronutrient contents of residues (spent biomass) of *Tetradesmus obliquus* (grown in Bristol medium and piggery wastewater, PWW) after subcritical water extraction at different temperatures (120, 170, and 220 °C) and biomass loading (1:10 and 1:80). Commercial mineral and organic fertilisers from SIRO® were used for comparison. NPK is given as percentage of N, P (as P<sub>2</sub>O<sub>5</sub>), and K (as K<sub>2</sub>O).

Nutrients	Bristol				PWW				Mineral Fertiliser SIRO®	Organic Fertiliser SIRO®
	WB	120 °C	170 °C	220 °C	WB	120 °C	170 °C	220 °C		
<b>Macronutrients (g/kg)</b>										
C	372	416	428	412	423	531	575	647	-	276
N	68.8	68.9	58.2	40.1	58.7	47.9	45.9	44.9	120	60
P	42.5	45.2	53.4	66.3	6.88	5.17	7.95	9.56	13.1	8
K	3.21	1.60	1.32	1.41	6.92	4.99	3.00	2.84	70.5	33
S	5.40	9.50	6.40	4.70	3.70	4.70	4.30	4.00	24	-
Ca	49.9	52.7	66.7	83.9	71.9	13.2	15.4	17.4	-	-
Mg	7.46	7.30	8.15	11.0	12.2	15.1	15.2	15.6	12	-
Ratio C/N	5	6	7	10	7	11	13	14	-	4.6
NPK	5-39-0.8	7-41-0.4	6-49-0.3	4-61-0.3	7-6-2	5-5-1	5-7-1	5-9-1	12-12-17	6-7-8
<b>Micronutrients (mg/kg)</b>										
Zn	149	214	262	323	652	548	698	834	100	-
Cu	69.8	78.7	100	124	93.4	124	162	203	-	-
Fe	21200	21400	25800	34300	2010	731	734	1010	-	-
Mn	1470	1430	1740	2290	428	158	183	192	-	-