

The dynamics of the reaction between O (1D) and DH and the isotopic effect

Carolina M. A. Rio^a and J. Brandão^a

^a Departamento de Química e Bioquímica, Faculdade de Ciências e Tecnologia, Universidade do Algarve, Campus de Gambelas, 8005-139 Faro, Portugal

The reaction $O(^1D) + H_2 \rightarrow OH + H$ and its isotopic variants plays an important role in atmospheric chemistry and also is important in combustion chemistry. It has been taken as a prototype for insertion reactions. As a result, it has been subject of several experimental studies. Also theoretical studies of its dynamics have been carried out.

In absence of energy barrier, the long-range interactions [1] between reactants should play an important role on the dynamics of the reaction of O (1D) with H_2 that mainly occurs in the ground state potential energy surface for H_2O [2]. Quasiclassical and capture studies [3] on a new potential energy surface that carefully reproduces these interactions are in close agreement with the most recent estimates for the thermal rate constant, when the effects of the excited surfaces is taken into account.

The isotope effect in these reactions is an additional test to the accuracy of the potential energy surface.

In this work we study the dynamics of the O (1D) + DH reaction at fixed collision energies [4,5] using quasiclassical trajectories calculations on a double-valued potential energy surface for H_2O [2].

In order to compare with experiment we have carried out quasiclassical trajectories calculations on O (1D) + DH reaction at the fixed translational energies of 2.05, 3.7, and 4.55 kcal mol⁻¹.

In particular, we analyse the reactive cross section, the opacity function, and the differential cross sections. In this work we also study the energy distribution of the products for the O (1D) + DH reaction. We also analyse the H/D ratio, $\Gamma_{H/D}$, where there are larger divergences between the results from different potential energy surfaces and experiment.

The agreement found between our QCT results [3,4] and experiment reinforces the accuracy of the H_2O potential energy surface used.

This work was supported by:

- FCT under the POCTI/CTA/41252/2001 research project, co-financed by the European community fund, FEDER.
- Centro Multidisciplinar de Química do Ambiente (CMQA).

References

- [1] João Brandão and Carolina M. A. Rio, *Chem. Phys. Lett.* (2003), 372(4), 866-872.
- [2] João Brandão and Carolina M. A. Rio, *J. Chem. Phys.* (2003), 119 (6), 3148-3159.
- [3] João Brandão and Carolina M. A. Rio, *Chem. Phys. Lett.* (2003), 377, 523-529.
- [4] Carolina M. A. Rio and João Brandão, "Isotopic effect of O (1D) + H_2 / D_2 / DH reactions using a double-valued DMBE potential energy surface for H_2O " (to be published).