

<b>DIAMETER OF INHIBITION ZONES (mm)</b>			
<b>ANTIBIOTIC</b>	<b>RESISTANT</b>	<b>INTERMEDIARY</b>	<b>SENSITIVE</b>
<b>Ampicilin AM – 10</b>	11	12 – 13	14
<b>Bacitracin B – 10</b>	8	9 – 12	13
<b>Penicilin PEN – 10</b>	<18	21 – 18	>29
<b>Eritomicin E – 15</b>	13	14 – 17	18
<b>Gentamicina GM – 10</b>	12	----	13
<b>Kanamycin K – 30</b>	13	14 – 17	18
<b>Nalidixic Acid NA – 30</b>	13	14 – 18	19
<b>Neomicin NEO – 30</b>	<12	13 – 16	>17
<b>Novobiocin NB – 30</b>	17	18 – 21	22
<b>Polymyxin B PB – 300</b>	8	9 – 11	12
<b>Streptomycin S – 10</b>	11	12 – 14	15
<b>Tetracycline TE – 30</b>	14	15 – 18	19
<b>Doxycycline D – 30</b>	12	13 – 15	16
<b>Sulfamethoxazole / trimethoprim SXT</b>	16	19 – 17	>20
<b>Oxytetracycline OT – 30</b>	<14	18 – 15	>19
<b>Oxolinic acid OA – 2</b>	10	16 – 20	30 – 40
<b>Flumequine AR – 30</b>	10	16 – 20	30 – 40
<b>Enrofloxacin ENRO – 5</b>	≤21	16 – 20	<21