

Amrit Kumar Mishra

## **Global change effects on seagrass ecosystem**

Thesis submitted for the degree of “Doctor of Philosophy” (Marine Sciences)  
Erasmus Mundus Program in Marine Ecosystem Health and Conservation  
(MARES)

### **Supervisors**

Professor Rui Santos

Professor Jason Hall Spencer

Dr. Joao Silva

Dr. Eugenia Apostolaki



**UAlg**

UNIVERSIDADE DO ALGARVE

**May 2017**



## **DECLARATION OF AUTHORSHIP**

This work is original research work and has not previously been submitted for a degree in any University. To the best of my knowledge and belief, the thesis contains no material previously published or written by another individual except where due reference is made in the thesis itself.

Amrit Kumar Mishra



## **COPYRIGHT**

The University of Algarve has the perpetual right with no geographical boundaries to archive and publicize this work through printed copies in paper or digitized or in any other format known or that becomes invented to broadcast through scientific repositories and to allow its copy and distribution for educational research and non-commercial purposes, with prior consent to its author and both the author and editor are credited in the published work.



SUPPORT



**FCT**

Fundação para a Ciência e a Tecnologia  
MINISTÉRIO DA EDUCAÇÃO E CIÊNCIA

**RESEARCH  
WITH  
PLYMOUTH  
UNIVERSITY**



Hellenic Center of Marine Research



This thesis and research work is dedicated to my late father Professor Laxminarayan Mishra, who always inspired and supported me to pursue a career in Science and work for the betterment of human society.

He always said to me,

“Education is not an end, but means to an end”



“Don’t follow the path everyone walks through,  
Find your own path through hard work and dedication, and  
Leave a trail behind”

“L. N. Mishra”



## Contents

<b>Abbreviations list.....</b>	<b>xiii-xiv</b>
<b>Acknowledgements.....</b>	<b>xv-xvi</b>
<b>Authors Declaration.....</b>	<b>xviii</b>
<b>Chapter 1. General Introduction.....</b>	<b>1-17</b>
1.1. Global changes .....	3-4
1.2. Seagrass ecosystem and carbon sequestration.....	4-6
1.3. Global change effects on seagrass.....	6-10
1.4. Main objectives.....	11
1.5. References.....	11-17
<b>Chapter 2. Short-term effects of CO<sub>2</sub> on the diel and seasonal response of <i>Zostera noltii</i> community metabolism to light</b>	<b>19-45</b>
2.1. Introduction.....	23-25
2.2. Methods.....	25-28
2.3. Results.....	28-34
2.4. Discussion.....	35-40
2.5. References.....	40-45
<b>Chapter 3. Population dynamics of <i>Cymodocea nodosa</i> in the vicinity of volcanic CO<sub>2</sub> seeps</b>	<b>47-79</b>
3.1. Introduction.....	51-53
3.2. Methods.....	53-56
3.3. Results.....	57-67
3.4. Discussion.....	68-71
3.5. References.....	71-74
3.6. Appendix.....	72-79
<b>Chapter 4. Elevated trace elements in sediments and seagrass at six Mediterranean volcanic seeps</b>	<b>81-122</b>
4.1. Introduction.....	85-88
4.2. Methods.....	88-92

4.3. Results.....	92-108
4.4. Discussion.....	119-114
4.5. References.....	115-122
<b>Chapter 5. General Discussion</b>	<b>123-133</b>
5.1. Main findings and implications for seagrass ecosystem.....	125-128
5.2. Summary and Main Conclusions.....	129-131
5.3. References.....	131-133

## Abbreviations list

**ANOVA**-Analysis of Variance

**BEDS**-Biological Effects Database for Sediments

**BSAF**-Bio- sediment accumulation factor

**CH<sub>4</sub>**-Methane

**Cu**-Copper

**Co**-Cobalt

**Cd**-Cadmium

**C**-Carbon

**CR**-Community Respiration

**Cs**-Caesium

**CO<sub>2</sub>**-Carbon dioxide

**CO<sub>3</sub><sup>2-</sup>**- Carbonate

**CO<sub>2</sub>Sys**- Carbon Dioxide System calculation

**DIC**-Dissolved inorganic carbon

**DW**-Dry weight

**ERL**- Effects Range Low

**ERM**- Effects Range Median

**Fe**-Ferrous (Iron)

**H<sub>2</sub>S**-Hydrogen Sulphide

**HCO<sub>3</sub><sup>-</sup>**-Bicarbonate

**HCl**- Hydrogen chloride

**Hg**-Mercury

**HNO<sub>3</sub>**- Nitric acid

**GCP**-Gross community production

**IRGA**- Infrared Gas Analyzer

**IPCC**- International Panel for Climate Change

**ICP-MS**- Induced couple plasms- Mass Spectrometry

**ICP-AES**- Induced Couple Plasma-Atomic Emission Spectroscopy

**LCP**-Light Compensation point

**LGC**-Laboratory for Government Chemistry

**M**- Mortality

**Mn**-Manganese

**N**-Nitrogen

**NE**-North East

**NW**-North West

**Ni**-Nickel

**NCP**-Net community production

**OA**-Ocean acidification

**PAR**-Photosynthetic active radiation

**PEL**-Probable Effects Level

**Pb**-Lead

**$p\text{CO}_2$** -partial pressure of Carbon dioxide

**PI**-Plastochrome interval

**R**- Recruitment rate (long-term)

**R<sub>0</sub>**- Recruitment rate (Present)

**SCUBA**-Self Contained Underwater Breathing Apparatus

**SE**-South East

**SW**-South West

**SQG**- Sediment Quality Guidelines

**SQG-Q**-Sediment Quality Guidelines Quotient

**Sr**-Strontium

**TA**-Total Alkalinity

**TE**- Trace elements

**TEL**-Threshold Effects Level

**Zn**-Zinc

## **Acknowledgements**

This dissertation would not be possible without the assistance of many people during these four years of my life. I would like to thank my supervisors Prof. Rui Santos, Prof. Jason Hall Spencer, Dr. Joao Silva and Dr. Eugenia Apostolaki, who all helped me during this thesis with their expertise, designing of the experiments and completing the task for which I would be grateful.

I am immensely grateful for all the help I have received during my field work from Ricardo Guerreiro in Portugal, Thanos Dailianis, Julius Glampedakis and Veronica Santinelli in field sampling and laboratory works respectively at HCMR, Greece. I would like to thank Dr. Eugenia Apostolaki in providing me necessary facilities during my sampling work at Greece. I am very grateful to Prof. Jason Hall Spencer, Dr. Marco Milazzo in helping me during my sampling at Vulcano, Italy. I would also like to thank Dr. Joao Silva and Dr. Irene Olive in helping me collect samples from Ischia and Panarea in Italy and Monya Costa in helping me with the laboratory works at CCMAR, Portugal.

I am also very grateful to Prof. Jason Hall Spencer and Andrew Fisher, Andrew Tonkin and Robert Clough at Plymouth University, UK in helping me with official proceedings and laboratory works respectively. I am very grateful to Prof. Murray Brown at the University of Plymouth for his guidance and support during my time at Plymouth.

I am very thankful to Prof. Rui Santos, Dr. Joao Silva and our ALGAE group and my buddy Laura Sordo for providing support during field work and logistics, and Carmen Santos for helping me with the Ria Formosa tidal model data at University of Algarve, Portugal.

I am very grateful to Hania Zwierzchowska for helping me during my work in making digital maps. I am very thankful to Thanos Dailianis for the underwater pictures used in the dissertation work.

Thank you, MARES, for funding the project and to Tim Deprez, Pieter Blondeel and Veerle Rogge for overseeing the administrative aspects which was quite challenging at times. At the University of Algarve, Prof. Adelino Canario and Mercedes Covas at the international mobility office also greatly assisted in the administrative aspects. Thanks to CCMAR for allowing me to use their facilities.

Finally, I am grateful to my Mother (Basant Manjari Mishra) and family to provide me the support and strength during my research work and all the time being away from them and Anchal Paramguru for her unconditional support during my last part of dissertation work.

I would like to thank all my friends who have supported me during my doctoral work and encouraged me to pursue a career in science.

## **AUTHOR'S DECLARATION**

At no time during the registration for the degree of Doctor of Philosophy has the author been registered for any other University award without prior agreement of the Graduate committee. Work submitted for this research degree at the University of Algarve has not formed of any other degree either at University of Algarve or at another establishment.

This study was financed with the aid of a studentship from the Joint Doctorate program MARES selected under Erasmus Mundus, co-ordinated by Ghent University (FPA 2011-16) and carried out in collaboration with University of Plymouth, UK and Hellenic Centre for Marine Research, Greece. Plymouth Marine Science and Education Foundation, Plymouth Marine Institute, Challenger Society, UK, Partnership for Observation of Global Oceans, Nippon foundation, European Space Agency and International Council for the Exploration of Sea have provided me travel bursaries to attend international conferences and training programs.

Relevant scientific seminars and conferences were regularly attended at which research outcomes were often presented.



## Short Abstract

Long and short effects of increased CO<sub>2</sub> on seagrass carbon metabolism were investigated. *Cymodocea nodosa* meadows, exposed to long term elevated CO<sub>2</sub> at shallow volcanic CO<sub>2</sub> seeps off Greece and Italy were investigated using population reconstruction techniques. Growth, morphometry, density, biomass and age structure was affected by high CO<sub>2</sub>. Above to below ground biomass ratio of *C. nodosa* were higher at CO<sub>2</sub> seeps. *C. nodosa* population grew faster but longevity of plants was lower at the seeps. The present recruitment (sampled year) of the seagrass was higher near the seeps. Carbon to nitrogen ratios (%DW) in leaves and annual leaf production of *C. nodosa* were higher at seeps. Elements (Fe and Cd, Cu, Co, Hg, Pb, Mn, Ni and Zn) were analysed from sediments and *P. oceanica* and *C. nodosa* compartments from CO<sub>2</sub> seeps off Greece and Italy. Sediment Quality Guidelines Quotient pollution index indicated higher element level seeps than reference sites with possibility of moderate to adverse biological impacts. Element concentration in the sediments and seagrass compartments were higher at seeps. Higher accumulation of elements in the seagrass compartments were observed. Net community production (NCP) and community respiration (CR) of intertidal *Z. noltii* and unvegetated sediment communities were measured under air exposed and CO<sub>2</sub> enriched conditions seasonally. Increase in NCP of *Z. noltii* and sediment community were observed with elevated CO<sub>2</sub> concentrations than natural conditions within similar light range. NCP of *Z. noltii* was higher than sediment community in both summer and winter seasons under CO<sub>2</sub> enriched conditions. CR of both communities were less affected under CO<sub>2</sub> enriched conditions. Light compensation point of *Z. noltii* were lower than sediment communities in summer season with elevated CO<sub>2</sub> levels. Seasonal community production of *Z. noltii* were higher than sediment communities. Significant effect of light on NCP was observed from linear regression model.

Keywords: Seagrass, community production, reconstruction techniques, element toxicity, CO<sub>2</sub> seeps



## Resumo

Foram investigados efeitos a curto e longo prazo do CO<sub>2</sub> no metabolismo do carbono das ervas marinhas. As pradarias de *Cymodocea nodosa*, expostas durante um longo período a CO<sub>2</sub> elevado, no vento vulcânico de CO<sub>2</sub> ao largo da Grécia e da Itália, foram investigadas utilizando técnicas de reconstrução populacional. O Crescimento, a morfometria, a densidade, a biomassa e a estrutura etária foram afetados pelo alto teor de CO<sub>2</sub>. A razão entre a biomassa da parte aérea e parte subterrânea de *C. nodosa* foi superior nos ventos de CO<sub>2</sub>. A população de *C. nodosa* cresceu mais rapidamente, mas a longevidade das plantas foi menor nos ventos de CO<sub>2</sub>. O recrutamento atual (ano amostrado) da erva marinha foi maior perto das infiltrações. As taxas de carbono - azoto (% PS) nas folhas e a produção foliar anual de *C. nodosa* foram maiores nos ventos de CO<sub>2</sub>. Os elementos (Fe e Cd, Cu, Co, Hg, Pb, Mn, Ni e Zn) foram analisados a partir de sedimentos e das várias partes de *P. oceanica* e *C. nodosa* dos ventos de CO<sub>2</sub> ao largo da Grécia e da Itália. O Índice de Poluição do Quociente das Diretivas de Qualidade do Sedimento indicou que o nível de elemento é mais alto nos sítios de CO<sub>2</sub> do que nos de referência com possibilidade de impactos biológicos moderados a adversos. A concentração de elementos nos sedimentos e nas várias partes das plantas foi maior nos sítios de CO<sub>2</sub>. Foi observada uma maior acumulação de elementos nas das ervas marinhas. A produção líquida da comunidade (NCP) e a respiração da comunidade (CR) das populações intertidais de *Z. noltei* e de zonas não sedimentadas foram medidas sob condições expostas ao ar e enriquecidas com CO<sub>2</sub>, sazonalmente. O NCP de *Z. noltei* e dos sedimentos comunidade foi mais elevado em concentrações elevadas de CO<sub>2</sub> do que as condições naturais, para um intervalo de luz semelhante. O NCP de *Z. noltei* foi maior do que no sedimento da comunidade nas estações de verão e inverno, em condições enriquecidas com CO<sub>2</sub>. A CR de ambas as comunidades foi menos afetada em condições enriquecidas com CO<sub>2</sub>. O ponto de compensação para a luz de *Z. noltei* foi menor do que o dos sedimentos das comunidades na estação do verão com elevados níveis de CO<sub>2</sub>. A produção comunitária sazonal de *Z. noltei* foi maior do que no sedimento das comunidades. O efeito significativo da luz no NCP foi observado a partir do modelo de regressão linear.

Palavras-chave: erva marinha, produção comunitária, técnicas de reconstrução, toxicidade de elementos, vazamentos de CO<sub>2</sub>



## Long Abstract

Rising carbon dioxide (CO<sub>2</sub>) concentrations in the atmosphere will increase the average *p*CO<sub>2</sub> level in the world oceans, which will have a knock-on effect on the marine ecosystem. Coastal seagrass communities one of the most productive marine ecosystems are predicted to benefit from the increase in CO<sub>2</sub> levels, but long term effects of elevated CO<sub>2</sub> on seagrass communities are less understood. Population reconstruction techniques was used to investigate the population dynamics of *Cymodocea nodosa* meadows, exposed to long term elevated CO<sub>2</sub> at volcanic seeps off Greece and Italy. Effect of elevated CO<sub>2</sub> was noticed on the growth, morphometry, density, biomass and age structure at CO<sub>2</sub> seeps. Above to below ground biomass ratio of *C. nodosa* were higher at CO<sub>2</sub> seeps than at reference sites. The plastochrome interval were similar at all CO<sub>2</sub> seeps. The shoot age and shoot longevity of plants were lower at seeps than reference sites. The present recruitment (sampled year) of the seagrass were higher than long-term average recruitment of the communities near the seeps. Carbon to nitrogen ratios (%DW) of *C. nodosa* were higher in leaves at seeps. Annual leaf production was higher near the seeps. This study suggests increased production of *C. nodosa* under elevated CO<sub>2</sub> levels, but other co-factors such as nutrients, trace metal toxicity must also be taken into consideration while predicting effects of future CO<sub>2</sub> concentrations.

Volcanic CO<sub>2</sub> seeps are now being used as natural analogues for ocean acidification studies although these areas can be affected by trace element input and may alter ecosystem responses to gradient in carbonate chemistry. Here Fe and a range of trace elements (Cd, Co, Cu, Hg, Mn, Pb, Ni and Zn) were analysed from sediments and from the roots, rhizomes and leaves of seagrass at six CO<sub>2</sub> seeps and reference sites off Greece and Italy. There were higher metal levels in sediment and seagrasses at all CO<sub>2</sub> seeps than reference sites. Sediment Quality Guideline Quotient, a commonly used pollution index, indicated that some of the metals (Cd, Cu, Hg, Ni) were in high enough concentrations to have adverse biological effects, such as Cu at Ischia site and Hg at Vulcano. Higher accumulation of elements from sediments in roots and leaves at CO<sub>2</sub> seeps were found from Bio Sediment Accumulation Factor index. There were higher levels of Cu, Fe, Mn and Zn in leaves and rhizomes for *P. oceanica* and higher levels of Cd, Co, Cu, Fe and Zn in *C. nodosa* compartments at CO<sub>2</sub> seeps. Fe and Mn were found with positive correlation within sediment-roots and sediment-rhizomes, whereas Cd, Co and Pb were found with positive correlation in compartments of *C. nodosa*. In *P. oceanica* positive correlation were only observed for Cd within sediment-roots and plant compartments. Low pH and ocean acidification increased the concentration of elements at CO<sub>2</sub> seeps than

reference sites. Thus, caution is needed, when using volcanic seep systems as analogue for the effects of rising CO<sub>2</sub>, as metals can reach levels that are toxic to seagrass, masking any potential benefits of increased levels of carbon dioxide for seagrass productivity.

Net community production (NCP) and community respiration (CR) were measured under air exposed and CO<sub>2</sub> enriched conditions for intertidal *Z. noltei* meadows and unvegetated sediment communities during emersion in summer and winter seasons. Community production and respiration were measured in-situ using benthic chambers. CO<sub>2</sub> flux under air and CO<sub>2</sub> enriched conditions were measured over a series of short term incubations (30min) using an infra-red gas analyser. Incident photosynthetic active radiation (PAR) was recorded during the incubations covering the daily and seasonal variation. Linear regression model was used to test the effects of irradiance on net community production. NCP of *Z. noltei* community were higher under CO<sub>2</sub> enriched conditions than air exposed conditions in both summer and winter seasons. There was no effect of CO<sub>2</sub> on the CR rate of *Z. noltei* community in summer season. NCP of sediment community were higher in summer season and winter season under CO<sub>2</sub> enriched conditions. Sediment CR rates were higher in winter than summer season. The light compensation point of *Z. noltei* and sediment community were lower in both seasons under CO<sub>2</sub> enriched conditions. Seasonal budget of community production was higher in *Z. noltei* than sediment communities. A clear effect of PAR was noticed on the net community production of both communities. Higher PAR intensities resulted in higher NCP under CO<sub>2</sub> enriched conditions for both communities. CO<sub>2</sub> enrichment will have a positive effect on the intertidal communities during emersion.

## Résumé

O aumento das concentrações de dióxido de carbono (CO<sub>2</sub>) na atmosfera aumentará o nível médio de pCO<sub>2</sub> nos oceanos mundiais, o que afetará o ecossistema marinho. As comunidades de ervas marinhas costeiras, um dos ecossistemas marinhos mais produtivos, prevêem beneficiar do aumento dos níveis de CO<sub>2</sub>, mas os efeitos a longo prazo de CO<sub>2</sub> elevado nas comunidades de ervas marinhas são menos compreendidos. As técnicas de reconstrução populacional foram utilizadas para investigar a dinâmica populacional dos prados de *Cymodocea nodosa*, expostos ao CO<sub>2</sub> elevado a longo prazo em infiltrações vulcânicas na Grécia e na Itália. O efeito do CO<sub>2</sub> elevado foi notado no crescimento, morfometria, densidade, biomassa e estrutura etária nas infiltrações de CO<sub>2</sub>. Acima da relação de biomassa abaixo do solo de *C. nodosa* foi maior nas infiltrações de CO<sub>2</sub> do que nos locais de referência. O intervalo de plastocromo foi semelhante em todas as infiltrações de CO<sub>2</sub>. A idade do tiroteio e a longevidade das plantas foram mais baixas nas infiltrações do que nos sites de referência. O recrutamento atual (ano amostrado) das ervas marinhas foi maior do que o recrutamento médio a longo prazo das comunidades perto das infiltrações. As proporções de carbono para nitrogênio (% DW) de *C. nodosa* foram maiores nas folhas nas infiltrações. A produção anual de folhas foi maior perto das infiltrações. Este estudo sugere uma produção aumentada de *C. nodosa* sob níveis elevados de CO<sub>2</sub>, mas outros cofactores como nutrientes, toxicidade de rótulos metálicos também devem ser levados em consideração ao prever os efeitos das futuras concentrações de CO<sub>2</sub>. As infiltrações vulcânicas de CO<sub>2</sub> agora estão sendo usadas como análises naturais para estudos de acidificação dos oceanos, embora essas áreas possam ser afetadas pela entrada de oligoelementos e podem alterar as respostas do ecossistema ao gradiente na química do carbonato. Aqui foram analisados Fe e uma série de oligoelementos (Cd, Co, Cu, Hg, Mn, Pb, Ni e Zn) a partir de sedimentos e raízes, rizomas e folhas de ervas marinhas em seis infiltrações de CO<sub>2</sub> e locais de referência na Grécia e na Itália. Houve maiores níveis de metais em sedimentos e seagrasses em todas as infiltrações de CO<sub>2</sub> do que nos sites de referência. O quociente de orientação de qualidade de sedimento, um índice de poluição comumente usado, indicou que alguns dos metais (Cd, Cu, Hg, Ni) estavam em concentrações suficientemente altas para ter efeitos biológicos adversos, como Cu no local de Ischia e Hg em Vulcano. Maior acumulação de elementos de sedimentos em raízes e folhas em vazões de CO<sub>2</sub> foi encontrada a partir do índice Biocentro de acumulação de sedimentos. Houve níveis mais elevados de Cu, Fe, Mn e Zn nas folhas e rizomas para *P. oceanica* e níveis mais elevados de Cd, Co, Cu, Fe e

Zn nos compartimentos de *C. nodosa* às infiltrações de CO<sub>2</sub>. Fe e Mn foram encontrados com correlação positiva dentro de raízes de sedimentos e rizomas de sedimentos, enquanto Cd, Co e Pb foram encontrados com correlação positiva em compartimentos de *C. nodosa*. Na correlação positiva de *P. oceanica* foram observados apenas Cd dentro de raízes de sedimentos e compartimentos de plantas. O baixo pH e a acidificação do oceano aumentaram a concentração de elementos nas infiltrações de CO<sub>2</sub> do que nos sites de referência. Assim, é necessário ter cuidado ao usar sistemas de infiltração volcânica como analógicos para os efeitos do aumento do CO<sub>2</sub>, pois os metais podem atingir níveis tóxicos para as ervas marinhas, mascarando quaisquer benefícios potenciais do aumento do nível de dióxido de carbono para a produtividade das ervas marinhas. A produção comunitária líquida (NCP) e a respiração comunitária (CR) foram medidas sob condições expostas ao ar e enriquecidas com CO<sub>2</sub> para prados intermicionais de *Z. noltei* e comunidades de sedimentos não estimados durante a emersão nas estações de verão e inverno. A produção comunitária e a respiração foram medidas in situ usando câmaras bentônicas. O fluxo de CO<sub>2</sub> no ar e as condições enriquecidas em CO<sub>2</sub> foram medidas ao longo de uma série de incubações de curto prazo (30 min) usando um analisador de gás infravermelho. A incidente de radiação fotossintética ativa (PAR) foi registrada durante as incubações cobrindo a variação diária e sazonal. O modelo de regressão linear foi utilizado para testar os efeitos da irradiância na produção comunitária líquida. NCP da comunidade *Z. noltei* foi maior em condições enriquecidas em CO<sub>2</sub> do que as condições expostas ao ar, tanto nas estações de verão como no inverno. Não houve efeito de CO<sub>2</sub> na taxa CR da comunidade *Z. noltei* no verão. O NCP da comunidade de sedimentos foi maior na temporada de verão e no inverno em condições de CO<sub>2</sub> enriquecidas. As taxas de CR de sedimento foram maiores no inverno do que na temporada de verão. O ponto de compensação de luz de *Z. noltei* e comunidade de sedimentos foi menor em ambas as estações sob condições enriquecidas em CO<sub>2</sub>. O orçamento sazonal da produção comunitária foi maior em *Z. noltei* do que as comunidades de sedimentos. Um efeito claro do PAR foi notado na produção comunitária líquida de ambas as comunidades. Maiores intensidades de PAR resultaram em maior NCP sob condições enriquecidas em CO<sub>2</sub> para ambas as comunidades. O enriquecimento de CO<sub>2</sub> terá um efeito positivo nas comunidades intermareais durante a emergência.





---

# General Introduction





### General Introduction

#### 1.1. Global changes

Many changes are underway on Earth including increased land and sea temperatures and increasing sea level, coupled with a host of secondary changes (IPCC, 2014). These changes are a result of increasing anthropogenic carbon dioxide levels (CO<sub>2</sub>) in the Earth's atmosphere due to human activities such as deforestation, cement production and fossil fuel burning. Atmospheric CO<sub>2</sub> have increased 25% since the Industrial Revolution and include the concentration of other greenhouse gasses (e.g. methane, nitrous oxide, chlorofluorocarbons) which are also rising (Solomon et al., 2007). The ocean is the largest carbon reservoir with nearly fifty times more carbon storage capacity than either the atmosphere or terrestrial reservoirs (Solomon et al., 2007). Anthropogenic CO<sub>2</sub> enters the ocean across the air-sea interface and this facilitates the sequestration of anthropogenic carbon emissions through physical, chemical and biological processes (McLeod et al., 2011) into the ocean's carbon sink (Feely et al., 2001). Once assimilated into the ocean, carbon availability and water circulation partially regulate the biological productivity and biochemistry of the oceans globally (Libes, 2009). The role of the ocean as a sink for CO<sub>2</sub> is driven by two processes; the solubility pump and the biological pump (Sabina and Feely, 2007; Libes, 2009; McLeod et al., 2011). The solubility pump works because CO<sub>2</sub> is more soluble in cold seawater than warm. Cold deep waters are generally rich in dissolved inorganic carbon (DIC) and when these deep waters upwell into warmer equatorial regions, extensive outgassing of CO<sub>2</sub> occurs to the atmosphere due to reduced solubility (Le Quéré et al., 2009; McLeod et al., 2011). The biological carbon pump refers to the uptake of CO<sub>2</sub> by marine plankton from surface water through photosynthesis, thus a small fraction (1/3) of algal biomass produced is transferred to the deep ocean and buried in sediments (Boyce et al., 2010; McLeod et al., 2011). Both carbon pumps help in regulate the uptake of carbon from the atmosphere by the oceans.

Over 35 giga tonnes per year of anthropogenic CO<sub>2</sub> produced due to human activities ends up in the world oceans and this is rapidly changing seawater carbonate chemistry (Royal Society, 2005; IPCC, 2014.). When CO<sub>2</sub> dissolves in water, it forms carbonic acid (H<sub>2</sub>CO<sub>3</sub>), which further dissociates into hydrogen (H<sup>+</sup>), bicarbonate (HCO<sub>3</sub><sup>-</sup>) and carbonate (CO<sub>3</sub><sup>2-</sup>) ions. The availability of more H<sup>+</sup> ions increases the acidity of the seawater and is known as “ocean acidification”, (Caldeira & Wickett, 2003) which has reduced the pH levels of surface seawater by an average of 0.1 units, with a predicted further decrease of 0.3-0.4 units by the end of the

2100 century (IPCC, 2014). The oceans are well buffered through changes in the relative proportions of  $\text{CO}_3^{2-}$ ,  $\text{HCO}_3^-$ , and  $\text{CO}_2$ . This DIC maintains the seawater pH within narrow limits although this buffering capacity diminishes as increased amounts of  $\text{CO}_2$  are absorbed by the ocean. This is because when  $\text{CO}_2$  dissolves, the chemical process that takes place reduces some carbonate ions ( $\text{HCO}_3^-$ ) required for the ocean pH buffer (Royal Society, 2005).

Elevated atmospheric  $\text{CO}_2$  is reducing carbonate and pH levels and increasing the concentration of aqueous  $\text{CO}_2$  and bicarbonate (Doney et al., 2009). The decrease in surface pH is happening so rapidly that this may make it difficult for many organisms to adapt to a changing chemical environment (Milazzo et al., 2016). Reduced carbonate ion concentrations are known to impact calcified organisms with decreased aragonite and calcite saturation states causing skeletal or shell dissolution in extreme cases, so calcifying organisms are expected to be especially vulnerable to rising  $\text{CO}_2$  levels (Kleypas et al., 1999; Andersson et al., 2007).

### 1.2. Seagrass ecosystem and Carbon sequestration

The coastal zones are characterized by several different ecosystems, each dominated by different suites of primary producers making coastal habitats one of the most productive regions in the world. Primary productivity in coastal systems can be dominated by phytoplankton, benthic microalgae, seaweeds, kelp, seagrasses, tidal marsh plants or mangroves (Thom, 2001). Along with high production rates coastal zones are also characterized for nutrient cycling and biodiversity (Thom, 2001). These habitats are important for the ocean's biogeochemical cycles, and they play a crucial role in providing ecosystem services (Gattuso et al., 1998; Erwin et al., 2009). Their physical structure provides breeding grounds and nurseries for crustaceans, finfish and shellfish, as well as food for reptiles, fish, birds and mammals (Coles et al., 2007). Coastal marine algae and plants are the basis of a detrital food chain, they process nutrients and contaminants, stabilize sediments and act as storm buffers. Economically, they support food webs and contribute to the local economy (Conservation International, 2008)

One of the most important, productive and dynamic types of coastal vegetation are seagrasses. Seagrasses are a relatively small group of flowering plants with about 50 species in 13 Genera and 5 Families (Coles et al., 2007) that are found all over the world. Seagrass growth is characterized by repeated structural units that in combination form the body of the plant (Hemminga & Duarte, 2000). These units also make up the rhizomes (root- like stems) that extend below the sediments and create thick networks of detritus, leaves, old rhizomes and

other organic matter, which can extend several meters below the surface. It is in these networks that carbon and nutrients are stored by seagrasses which makes seagrasses potential carbon sink (Hemminga & Duarte, 2000; Borum et al., 2004). These networks are called mattes which are mostly found in *P. oceanica*.

Seagrasses cover a very small area in the coastal systems and have low biomass with an average of  $184 \text{ gCm}^{-2}$  (Laffoley et al., 2009), but due to their high primary production they contribute 1% of the net primary production and 12% of the net ecosystem production in the ocean, around  $600 \text{ gCm}^{-2}$  (Fourqurean et al., 2012). This high productivity, combined with their global distribution, makes them stand out from other coastal systems and they bury  $48 - 112 \text{ Tg C y}^{-1}$  globally (McLeod et al., 2011), with some estimates that they account for nearly 10% of the yearly carbon burial in the oceans (Fourqurean et al., 2012). For example, *Posidonia oceanica*, is thought to be responsible for long term, large-scale carbon burial due to its long survival rate and large mattes formation (Borum et al., 2004), this species is endemic to and widespread throughout most of the coastal Mediterranean, and contains one of the largest coastal reservoirs of carbon in world (Lo-Iacono et al., 2008).

Seagrass meadows are thought to play an important role in the sequestration of carbon that would otherwise remain in the atmosphere and exacerbate climate change (Nellemann et al., 2010; Duarte et al., 2010; Kennedy et al., 2010). Seagrasses sequester carbon within their underlying sediments, within living biomass aboveground (roots), and within non-living biomass (leaf litter and dead shoots) and account for as much as 71% of the carbon storage in ocean sediments along with mangroves and saltmarsh ecosystems (Nellemann et al., 2009). “Blue carbon” is the carbon that is captured by the living organisms in the ocean and coastal ecosystems and stored in the form of biomass and in sediments of the oceans. This carbon is sequestered both short term (decennial) in biomass and long term (millennial) time scales in sediments (Duarte et al., 2005a; Lo Iacono et al., 2008, McLeod et al., 2011). The carbon burial in the seagrass ecosystem is higher than other coastal ecosystems (Duarte et al., 2013) and due to their anaerobic sediment conditions and sediment dynamics, the buried carbon stays buried for millennia (Laffoley et al., 2009), for instance in *Posidonia oceanica* meadows, which are extremely efficient at both burying and storing carbon for millennia (Lo-Iacono et al., 2008; Fourqurean et al., 2012). The rate and success of carbon sequestration varies between and within seagrass species based on a whole suite of natural processes including herbivory, export of nutrients and decomposition (Gattuso et al., 1998).

Seagrass ecosystems will not become saturated with C deposits, as sediments will accrete vertically with response to rising sea levels, assuming ecosystem health is maintained (McKee et al., 2007). Therefore, the rate of sediment C sequestration and the size of the sediment C sink may continue to increase over time (Chmura et al., 2003), for example in *P. oceanica* meadows in Portlligat Bay, Spain (McKee et al., 2007). The global changes are likely to affect the C sequestration capacity of seagrass ecosystem, but exact prediction is difficult (McLeod et al., 2011). As such, global storage estimates differ between species, sources and measuring methods, despite these uncertainties, seagrass contribute a large portion of the ocean's ability to sequester carbon (McLeod et al., 2011; Mazarrasa et al., 2015).

### 1.3. Global Change Effects on Seagrasses

Global change is widely recognized as a real threat to our natural systems (Laffoley et al., 2009). Coastal zones and seagrasses are no exception. Seagrass meadows, while supporting a huge array of diversity, are often themselves not particularly diverse. Large areas of seagrass are often dominated by a single species (Green and Short, 2003; Ehlers et al., 2008). They are also exceptionally vulnerable to changes in water clarity, light availability and temperature (Bjork et al., 2008; Laffoley et al., 2009). The combination of genetic homogeneity and vulnerability to changes in climatic and environmental factors makes seagrasses particularly vulnerable to global warming (Ehlers et al., 2008; Jorda et al., 2012). In fact, some studies suggest that they are more vulnerable than many other ocean systems (Nellmann et al., 2009). Seagrass ecosystems can shift from being a net sink of C to a source of C because of changes in climate (e.g. changes in precipitation and temperature), atmospheric composition effects (e.g. CO<sub>2</sub> fertilization, nutrient deposition, damage by pollution), and land-use-change effects (e.g. deforestation, afforestation, agricultural practices; IPCC, 2007)

Climate change may exacerbate these factors and seagrass meadows are already under stress due to climate change (Short and Neckles, 1999; Duarte et al., 2008b; Waycott et al., 2009), particularly where maximum temperatures already approach their physiological limit (Short and Neckles, 1999). Increases in water temperature will directly affect seagrass metabolism and the maintenance of a positive carbon balance (Evans et al., 1986; Zimmerman et al., 1989; Collier and Waycott, 2014) thus affecting the growth rates (Koch et al., 2013). Temperature increase effects will be species -specific and result in increase/decrease of production at thermal tolerance levels (Short and Neckles, 1999; Collier and Waycott, 2014).

Heatwaves can lead to widespread seagrass shoot mortality (Reusch et al., 2005; Diaz-Almela et al., 2009; Rasheed and Unsworth, 2011; Repolho et al., 2017), for instance shoot mortality in *P. oceanica* in Mediterranean Sea (Marbà and Duarte, 2010). Increased temperature will also affect the seagrass distribution and abundance through direct effects on flowering events of seagrass (Duarko and Moffler, 1987; Short and Neckles, 1999; Marbà and Duarte, 2010; Diaz-Almela et al., 2007), whereas in other studies warming have shown to increase seagrass flowering but the larger recruitment rate is insufficient to compensate for the losses resulting from elevated temperature (Diaz-Almela et al., 2009). Before temperature tolerance levels are reached, increase in water temperature may increase productivity, but it may also increase the productivity of algae that shades seagrasses from sunlight (Watkiss & Bonjean, 2012; Collier and Waycott, 2014). Along with coastal developments, eutrophication events will increase turbidity (Burkholder et al. 2007), hence decreasing light intensities will have consequences on seagrass productivity (Short and Neckles, 1999; Bjork et al., 2008) and colonization depth (Duarte, 1991). Finally, an increase in the frequency or severity of storms may cause significant additional mechanical damage to the seagrass beds (Connolly, 2009; Watkiss & Bonjean, 2012).

Though other global change factors will affect seagrass ecosystem, increased CO<sub>2</sub> concentrations will play a major role on their growth and productivity and carbon cycling capacity in coastal areas (Duarte and Chiscano 1999; Hemminga and Duarte, 2000). Seagrass meadows are reported as one of few ecosystems that may benefit from rising CO<sub>2</sub> levels because their photosynthetic rates have been considered Ci-limited at current oceanic CO<sub>2</sub> levels (Beer and Koch 1996; Thom 1996; Zimmerman et al., 1997; Invers et al., 2001). Consequently, increase in seagrass production and growth may occur in a future high-CO<sub>2</sub> scenario. CO<sub>2</sub> enrichment may also affect nitrogen uptake and the assimilation process, as growth enhancement at high-CO<sub>2</sub> concentrations is expected to increase the nitrogen demand of plants (Stitt and Krapp, 1999; Ow et al., 2016). In addition, the relative uptake rates of ammonium and nitrate may be altered by the acidification of the seawater resulting from CO<sub>2</sub> enrichment (Short and Neckles, 1999; Ow et al., 2016), due to involvement of protons (H<sup>+</sup>) in nitrogen transport across plasma membrane. From ionic balance perspective, lower pH levels in seawater may reduce the ammonium uptake rates of seagrasses, whereas nitrate uptake rates may be unaffected or even increased (Short and Neckles, 1999; Ow et al., 2016) or decreased (Alexandre et al., 2012)

Both short and long term experiments of elevated CO<sub>2</sub> conditions on seagrass have shown that net photosynthetic rate and photosynthetic efficiency was positively affected by CO<sub>2</sub> enrichment of seawater for *Z. noltii* (Welsh et al., 1997; Invers et al., 2001; Alexandre et al., 2012) and *Z. marina* (Zimmerman et al. 1997; Palacios and Zimmerman, 2007; Pajusalu et al., 2016), whereas no direct effects of high CO<sub>2</sub> on plant biochemistry or population level traits were found for *Z. noltii* (Martinez-Crego et al., 2014). Plants exposed to CO<sub>2</sub>-enriched conditions showed higher photosynthetic rates at saturating irradiances and were photosynthetically more efficient with higher reproductive output, below ground to above ground biomass and vegetative proliferation at limiting light intensities, when compared with plants exposed to the current CO<sub>2</sub> concentration (Welsh et al. 1997; Palacios and Zimmerman, 2007). This results are due to a higher carboxylation activity relatively to oxygenation activity from Rubisco in presence of a higher CO<sub>2</sub>/O<sub>2</sub> ratio, as CO<sub>2</sub> and O<sub>2</sub> are competitive inhibitors for Rubisco's active site (Furbank 1998; Taiz and Zeiger, 2002, Pajusalu et al., 2016; Repolho et al., 2017). Significant effects of high CO<sub>2</sub>, when combined with different factors (light, temperature and nutrients) have yielded different variation in results for *Z. noltii* (Alexandre et al., 2012; Martinez-Crego et al., 2014).

Experimental studies on *Cymodocea nodosa* and *Posidonia oceanica* have indicated, that increase in CO<sub>2</sub> concentrations will favour their utilization of inorganic carbon sources for higher photosynthetic activity (Invers et al., 2001, Apostolaki et al., 2014), whereas gene expression of *C. nodosa* photosynthetic genes expressed a down regulation and decrease in net primary production at elevated CO<sub>2</sub> levels (Olive et al., 2017). Short term mesocosm experiments with elevated CO<sub>2</sub> on *C. nodosa* have been observed with lower net community production (Khan et al., 2016). Lower density, biomass and above to below ground biomass was observed for *C. nodosa* at high CO<sub>2</sub> concentrations (Apostolaki et al., 2014), whereas increased density and biomass was observed for *C. rotundata* and *C. serrulata* (Takahashi et al., 2016). Carbon content and  $\delta^{13}\text{C}$  of leaves declined with increasing  $p\text{CO}_2$  for *C. nodosa* (Apostolaki et al., 2014) and *C. serrulata* and *C. rotundata* (Takahashi et al., 2016). High CO<sub>2</sub> and low pH conditions also reduces the concentration of phenolic protective substances of *C. nodosa* followed by higher grazing rates, which can alter the coastal carbon flux of seagrasses (Arnold et al., 2012). In case of *Posidonia oceanica* increase in CO<sub>2</sub> concentration was observed with increase in net carbon balance, which can also increase the depth limit of the seagrass (Invers et al., 2002; Vizzini et al., 2010) and shoot density (Hall Spencer et al., 2008),

whereas limited effects on shoot productivity (Cox et al., 2015) and no effects of high CO<sub>2</sub> were observed on leaf biomass, biomass and photosynthesis (Cox et al., 2016).

Most seagrass species possess the ability to utilize HCO<sub>3</sub><sup>-</sup> (Koch et al., 2013), as HCO<sub>3</sub><sup>-</sup> represents higher percentage of the total DIC pool in seawater than CO<sub>2</sub> (Stum and Morgan, 1981). So, carbon limitation of seagrasses in the current scenario is mitigated by shifting towards use of HCO<sub>3</sub><sup>-</sup>, although use of HCO<sub>3</sub><sup>-</sup> is species specific for seagrasses (Invers et al., 2001). Irrespective of photosynthetic mechanisms, a key issue in predicting the response to rising CO<sub>2</sub> is whether photosynthesis and growth are saturated by seawater DIC under present-day conditions of inorganic C speciation, pH, O<sub>2</sub> and temperature. In many cases an increase in DIC will result in higher photosynthetic and growth rates (Invers et al., 2001)

Seagrass photosynthesis appears limited by current DIC, due to the slow diffusive supply of CO<sub>2</sub> to the leaves, and possibly a less effective use of HCO<sub>3</sub><sup>-</sup> when compared to many macroalgae (Beer et al. 1989, Beer and Rehnberg, 1997; Zimmerman et al., 1997), for instance *Posidonia* and *Cymodocea* sp. with higher capacity for HCO<sub>3</sub><sup>-</sup> use can have higher production than compared to *Zostera* sp. with lower efficiency for HCO<sub>3</sub><sup>-</sup> utilization (Duarte, 1991; Invers et al. 2001; Beer et al., 2002). Several seagrass species have also exhibited higher photosynthesis, increased reproduction, below ground biomass, and greater production of non-structural carbohydrates, as well as lower leaf-N and chlorophyll under elevated CO<sub>2</sub> (Durako, 1993; Zimmerman et al., 1997; Beer & Koch, 1996; Palacios & Zimmerman, 2007; Jiang et al., 2010). Thus, CO<sub>2</sub> enrichment can shift C allocation to carbohydrates and away from N-containing compounds, such as Rubisco while increasing efficiency in seagrass production (Hemminga and Duarte, 2000; Vizzini et al., 2010), like findings in terrestrial C<sub>3</sub> species (Leakey et al., 2009). Consequently, an increase in dissolved CO<sub>2</sub> could be positive for photosynthesis and growth in seagrasses that are presently under saturated with respect to DIC, regardless of their capacity to utilize HCO<sub>3</sub><sup>-</sup>. Seagrasses through their metabolic activity can alter the chemical properties of the meadow (Hendriks et al., 2014a), thus making the organisms within the meadow less vulnerable to ocean acidification due to large diel pH changes. The increased productivity during the day may buffer the impacts of ocean acidification, particularly for calcifiers by providing a daily window of maximum calcium carbonate saturation where calcification can be more efficient and finally ocean acidification could stimulate seagrass productivity and thus increase buffering capacity (Shaw et al., 2013; Hendriks et al., 2013; Hendriks et al., 2014b).

Saying that studies on short term effects of CO<sub>2</sub> enrichment on seagrasses have mostly done under experimental conditions and using ex-situ results for predicting the possible impacts of elevated CO<sub>2</sub> on seagrass is difficult as there are interactions between various factors (light, temperature, nutrients, epiphytic growth on leaf surface of seagrass) in-situ which determines the response of seagrass to CO<sub>2</sub> enrichments. This increases the need of short term CO<sub>2</sub> enrichment experiments in situ on seagrasses to have a better understanding on the effects of elevated CO<sub>2</sub>. Similarly, to understand long term elevated CO<sub>2</sub> effects most studies have preferred volcanic CO<sub>2</sub> seeps where seagrass ecosystems are found. The response observed from various studies on seagrass at these CO<sub>2</sub> seeps are mostly species specific with some seagrasses showing positive responses, for example *C. serrulata* at CO<sub>2</sub> seeps of Papua New Guinea (Russel et al., 2013) whereas negative responses were observed for *C. nodosa* at Vulcano CO<sub>2</sub> seeps (Vizzini et al., 2013; Apostolaki et al., 2014). Both positive and negative responses at these CO<sub>2</sub> seeps can be related to other factor such as trace element levels which are higher at these CO<sub>2</sub> seeps due to continuous input from the seeps and can act as a confounding factor to seagrass ecosystem and associated biota due to their toxic effects. These element concentrations, their bioavailability and possible toxicity is going to increase in future ocean conditions due to low pH (Millero et al., 2009) and this can have negative impacts on the seagrass response to elevated CO<sub>2</sub> levels. So, while using natural CO<sub>2</sub> seeps for effects of elevated CO<sub>2</sub> levels trace elements levels should be taken into consideration

### Main Objectives

Considering the predicted changes to the marine environment, the importance of seagrass to the coastal ecosystem, the carbon metabolism of seagrass and the short and long term impacts of increased CO<sub>2</sub> on seagrass ecosystem, the main objectives of the thesis are

- i) to understand the short-term impacts of elevated CO<sub>2</sub> on the diel and seasonal variations on the community metabolism intertidal seagrass *Zostera noltii* community with respect to light
- ii) to understand the long -term impacts of elevated CO<sub>2</sub> on the growth, biomass, density, morphometry of seagrass *Cymodocea nodosa* ecosystem at CO<sub>2</sub> seeps using reconstruction techniques
- iii) to understand the trace elements variation within sediments, their accumulation within the seagrass *Cymodocea nodosa* and *Posidonia oceanica*, possible impacts on seagrass with respect to future ocean conditions of elevated CO<sub>2</sub> and low pH at CO<sub>2</sub> seeps

### 1.4. References

- Alexandre, A., Silva, J., Buapet, P., Bjork, M., et al., 2012. Effects of CO<sub>2</sub> enrichment on photosynthesis, growth and nitrogen metabolism of the seagrass *Zostera noltii*. *Ecol. and Evol.* 2:2626-2635
- Andersson, A.J., Bates, N.R., and Mackenzie, F.T., 2007. Dissolution of carbonate sediments under rising pCO<sub>2</sub> and ocean acidification: observations from Devil's Hole, Bermuda." *Aquat. Geochem.* 13: 237-264.
- Apostolaki, E.T., Vizzini, S., Hendriks, I.E., Oslon, Y.S., 2014 Seagrass ecosystem response to long term high CO<sub>2</sub> in a Mediterranean volcanic vent. *Marine Environ. Res.* 99: 9-15
- Arnold, T., Mealey, H., Leahey, H., Miller, W., et al., 2012. Ocean acidification and loss of Phenolic substances in marine plants. *PLoS ONE*.4: 35107
- Beer, S., 1989. Photosynthesis and photorespiration in marine angiosperms. *Aquat. Bot.* 34: 153-166
- Beer, S. and Rehnberg, J., 1997. The acquisition of inorganic carbon by the seagrass *Zostera marina*. *Aquatic Botany* 56: 277-283
- Beer, S., Koch, E., 1996. Photosynthesis of marine macroalgae and seagrasses in globally changing CO<sub>2</sub> environments. *Mar. Ecol. Prog. Ser.* 144: 199–204.
- Beer, S., Bjork, M., Hellblom, F., Axelsson, L., 2002. Inorganic carbon utilization in marine angiosperms (seagrasses). *Fun. Plant Biol.*, 29:1–6.

- Bjork, M., Short, F., Mcleod, E. & Beer, S., 2008. Managing Seagrasses for Resilience to Climate Change. Gland, Switzerland: IUCN.
- Borum, Jens., Duarte, C., Krause, D., Greve, T., 2004. An introduction to European seagrass monitoring and management
- Boyce, D.G., Lewis, M.R., Worm, B., 2010., Global phytoplankton decline over the past century. *Nature*. 466: 591-596
- Caldeira, K., Wickett, M. E., 2003. Oceanography: anthropogenic carbon and ocean pH. *Nature*. 425: 365
- Chmura, G.L., Anisfeld, S.C., Cahoon, D.R., and Lynch, J.C., 2003. Global carbon sequestration in tidal, saline wetland soils. *Global Biogeochem. Cy.* 17: 1111
- Coles, R. G., McKenzie, L. J., Rasheed, M. A., Mellors, J. E., et al., 2007. Status and Trends of Seagrass Habitats in the Great Barrier Reef World Heritage Area. Report to the Marine and Tropical Sciences Research Facility. Reef and Rainforest Research Centre Limited, Cairns (122 pp.).
- Collier, C., Waycott, M., 2014. Temperature extremes reduce seagrass growth and induce mortality. *Mar. Pol. Bull.* 83:483-490
- Conservation International., 2008. Economic Values of Coral Reefs, Mangroves, and Seagrasses: A Global Compilation. Centre for Applied Biodiversity Science, Conservation International, Arlington, VA, USA.
- Connolly, R.M., 2009. Seagrass. In a marine climate change impacts and adaptation report card for Australia 2009 (Eds. E. S. Poloczanska, A. J. Hobday and A. J. Richardson), NCCARF Publication 05/09
- Cox, T. E., Gazeau, F., Alliouane, S., Hendriks, I. E., et al., 2016. Effects of in situ CO<sub>2</sub> enrichment on structural characteristics, photosynthesis, and growth of the Mediterranean seagrass *Posidonia oceanica*. *Biogeosciences* 13:2179–2194
- Diaz- Almela, E., Marba, N., Duarte, C.M., 2007. Consequences of Mediterranean warming events in seagrass (*Posidonia oceanica*) flowering records. *Glo. change Biol.* 13: 224-235.
- Diaz-Almela, E., Marba, N., Martinez, R., Santiago, R., et al., 2009. Seasonal dynamics of *Posidonia oceanica* in Magalluf Bay (Mallorca, Spain): temperature effects on seagrass mortality. *Lim. and Oceano.*54:2170-2182
- Doney, S.C., Fabry, V.J., Feely, R.A., Kleypas, J.A., 2009. Ocean acidification: The other CO<sub>2</sub> problem. *Annu. Rev. Mar. Sci.* 1:169-192
- Duarte, C.M., 1991. Allometric scaling of seagrass form and productivity. *Mar. Ecol. Prog. Ser.*77:289-300.
- Duarte, Carlos M., 1999. Seagrass Biomass and Production: a Reassessment. *Aq. Bot.* 65:159-74
- Duarte, C.M., Chiscano, C.L., 1999. Seagrass biomass and production: a reassessment. *Aq. Bot.* 65:159-174

Duarte, C.M., Borum, J., Short, F.T., Walker, D.I., 2008b. Seagrass ecosystems: their global status and prospects. In: Aquatic Ecosystems (ed Polunin NVC), pp. 281–294. Cambridge University Press, Cambridge.

Duarte, C.M., Núria, M., Gacia, E., Fourqurean, J.W., et al., 2010. Seagrass community metabolism: Assessing the carbon sink capacity of seagrass meadows. *Glo. Biogeo. Cyc.* 24: GB4032

Duarte, C.M., Kennedy, H., Marbà, N. and Hendriks, I., 2013. Assessing the capacity of seagrass meadows for carbon burial: Current limitations and future strategies. *Oc. & Co. Man.* 83: 32-38.

Duarte, C.M., Borum, J., Short, F.T., and Walker, D.I., 2005b. Seagrass ecosystems: their global status and prospects. In: Polunin NVC (Ed). *Aquatic ecosystems: trends and global prospects*. Cambridge, UK: Cambridge University Press

Duarte, C.M., Middelburg, J., and Caraco, N., 2005a. Major role of marine vegetation on the oceanic carbon cycle. *Biogeosciences* 2: 1–8.

Duarko, M.J., Moffler, M.D., 1987. Factors affecting the reproductive ecology of *Thalassia testudinum* Hydrocharitaceae. *Aquat. Bot.* 27: 79-96

Duarko, M.J., 1993. Photosynthetic utilization of CO<sub>2</sub> and HCO<sub>3</sub><sup>-</sup> in *Thalassia testudinum* (Hydrocharitaceae). *Mar. Biol.* 115: 373-380

Erwin, K. L., 2009. Wetlands and Global Climate Change: The Role of Wetland Restoration in a Changing World." *Wet. Ecol. and Man.* 17: 71-84.

Ehlers, A., Worm, B., Reusch, T., 2008. Importance of genetic diversity in eelgrass *Zostera marina* for its resilience to global warming. *Mar. Ecol. Prog. Ser.* 355: 1–7

Evans, A.S., Webb, K.L., Penhale, P.A., 1986. Photosynthetic temperature acclimation in two co-existing seagrass *Zostera marina* L. and *Ruppia maritima* L. *Aquat. Bot.* 24: 185-197

Falkowski, P., Scholes, R. J., Boyle, E., Canadell, J., et al., 2000. The global carbon cycle: a test of our knowledge of Earth as a system. *Science*, 290: 291–296.

Feely, R.A., Sabine, C.L., Takahashi, T., Wanninkhof, R., 2001. The uptake and storage of carbon dioxide in the ocean: The global CO<sub>2</sub> survey. *Oceanography*. 14:18-32.

Fourqurean, J.W., Duarte, C.M., Kennedy, H., Marbà, N, et al., 2012. Seagrass ecosystem as a globally significant carbon stock. *Nat. Geo.* 5:505-509.

Furbank, R. T., 1998. Photosynthesis, a comprehensive treatise. Cambridge Univ. Press, Cambridge, U.K. C4 pathway. Pp. 123–135 in A. S. Raghavendra, ed.

Gattuso, J.-P., Frankignoulle, M., Bourge, I., Romaine, S., et al., 1998. Effect of calcium carbonate saturation of seawater on coral calcification. *Glob. Planet. Change.* 18: 37-46

Gattuso, J. P., 1998. Carbon and Carbonate Metabolism in Coastal Aquatic Ecosystems." *Annual Review of Ecology, Evol. and System.* 29: 405-34.

Green, E.P., Short, F.T., 2003. *World Atlas of Seagrasses*. California University Press, 310 pp

- Hall-Spencer, J.M., Rodolfo-Metalpa, R., Martin, S., Ransome, E., et al., 2008. Volcanic carbon dioxide vents show ecosystem effects of ocean acidification. *Nature*. 454: 96–99.
- Hemminga, M., Duarte, C.M., 2000. *Seagrass Ecology*. Cambridge University Press. New York, New York
- Hendriks, I. E., Olsen, Y. S., Ramajo, L., Basso, L., et al., 2014a. Photosynthetic activity buffers ocean acidification in seagrass meadows, *Biogeosciences*. 11: 333–346
- Hendriks, I. E., Duarte, C. M., Olsen, Y. S., Steckbauer, A., et al., 2014b. Biological mechanisms supporting adaptation to ocean acidification in coastal ecosystems. *Estuar. Coast. Shelf S.* 152: 1–8
- Houghton, R.A., Woodwell, G.M., 1989. Global climate change. *Sci. Amer.* 260:36-44.
- Invers, O., Tomas, F., Perez, M., Romero, J., 2002. Potential effect of increased global CO<sub>2</sub> availability on the depth distribution of the seagrass *Posidonia oceanica* (L.) Delile: a tentative assessment using a carbon balance model. 73:1191-1198
- Invers, O., Zimmerman, R.C., Alberte, R.S., Perez, M., et al., 2001. Inorganic carbon sources for seagrass photosynthesis: An experimental evaluation of bicarbonate use in species inhabiting temperate waters. *Jour. Of Exper. Mar. Bio. And Ecol.* 265:203-217
- IPCC., 2007. *Climate change: The Physical Science Basis.*” Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge University Press: Cambridge, UK and New York, NY, USA
- IPCC (International Panel for Climate Change)., 2014. *Fifth Assessment Report on Climate change: Impacts, Adaptation and Vulnerability*. New York
- Jorda, G., Marba, N., Duarte, C.M., 2012. Mediterranean Seagrass vulnerable to regional climate warming. *Nat. Cli. Change*. 2: 821-824.
- Jiang, Z., Huang, X., Zhang, J., 2010. Effects of CO<sub>2</sub> enrichment on photosynthesis, growth, and biochemical composition of seagrass *Thalassia hemprichii* (Ehrenb.) Aschers. *J. Integr. Plant. Biol.* 52: 904-913.
- Kennedy, H., Beggins, J., Duarte, C.M., Fourqurean, J.W., et al., 2010. Seagrass sediments as a global carbon sink: isotopic constraints. *Glo. Biogeochemi. Cy.* 24: 1-8
- Kleypas, J.A., Buddemeier, R.W., Archer, D., Gattuso, J.P., et al., 1999. Geochemical consequences of increased atmospheric carbon dioxide on coral reefs. *Science*. 284: 118–120
- Khan, F., Alexandre, A., Ullah, H., Santos, R., 2016. Effects of elevated CO<sub>2</sub> and nutrients on the community metabolism of a *Cymodocea nodosa* bed. *Turk. J. of Bot.* 3: 250-257
- Koch, M., Bowes, G., Ross, C., Zhang, X.H., 2013. Climate change and ocean acidification effects on seagrasses and marine macroalgae. *Glo. Change Biol.* 19: 103-132.
- Le Quéré, C., Raupach, M.R., Canadell, J.G., Marland, G. et al. 2009. Trends in the sources and sinks of carbon dioxide. *Nat. Geo.* 2:831-836.

- Laffoley, Dan., 2009. The Management of Natural Coastal Carbon Sinks. Gland, Switzerland: International Union for Conservation of Nature and Natural Resources.
- Libes, S.M., 2009. Introduction to Marine Biogeochemistry. Second Edition
- Lo Iacono, C., Mateo, M.A., Gracia, E., Guasch, L., et al., 2008. Very high- resolution seismo-acoustic imaging of seagrass meadows (Mediterranean Sea): implications for carbon sink estimates. *Geophys. Res. Lett.* 35: L18601
- Leakey, A.D.B., Ainsworth, E.A., Bernacchi, C.J., Rogers, A., et al., 2009. Elevated CO<sub>2</sub> effects on plant carbon, nitrogen, and water relations: six important lessons from FACE. *J. of Exp. Bot.* 60: 2859–2876
- Marba, N., Duarte, C.M., 2010. Mediterranean warming triggers seagrass (*Posidonia oceanica*) shoot mortality. *Glo. Change Bio.* 8:2366-2375
- Martinez-Crego, B., Olive, I., Santos, R., 2014., CO<sub>2</sub> and nutrient driven changes across multiple levels of organization in *Zostera noltii* ecosystems. *Biogeo. Discus.* 11: 5239-5274
- Mazarrasa, I., Marba, N., Lovelock, C.E., Serrano, O., et al., 2015. Seagrass meadows as a globally significant carbonate reservoir. *12: 4993-5003*
- McKee, K.L., Cahoon, D.R., and Feller, I., 2007. Caribbean mangroves adjust to rising sea level through biotic controls on change in soil elevation. *Global Ecol. Biogeogr.* 16: 545–56
- McLeod, E., Chmura, G., Bouillon, S., Salm, R., et al., 2011. A blueprint for carbon: Toward an improved understanding of the role of vegetated coastal habitats in the sequestering CO<sub>2</sub>. *Front. In Eco. And the Env.* 10:552-560
- Mitchell, J.F.B., 1989. The greenhouse effect and climate change. *Rev. Geophys.* 27: 115-139.
- Millero, F.J., Woosley, R., DiTrollo, B., Waters, J., 2009. Effect of ocean acidification on the speciation of metals in seawater. *Oceanography.* 22:72–85.
- Nellemann, C., Corcoran, E., Duarte, C.M., Valdes, L., et al., 2009. Blue carbon. A rapid response assessment. GRID-Arendal: United Nations Environment Programme.
- Olive, I., Silva, J., Lauritano, C., Costa, M.M., et al., 2017. Linking gene expression to productivity to unravel long and short term responses of seagrasses exposed to CO<sub>2</sub> in volcanic vents. *Sci. Rep.* 7:42278
- Orr, J.C., Fabry, V.J., Aumont, O., Bopp, L. et al., 2005. Anthropogenic ocean acidification over the twenty-first century and its impact on calcifying organisms. *Nature.* 437: 681-686
- Ow, Y.X., Vogel, N., Collier, C.J., Holtum, J.A.M., et al., 2016. Nitrate fertilisation does not enhance CO<sub>2</sub> responses in two tropical seagrass species. *Sci. Rep.* 6: 23093.
- Palacios, S.L., Zimmerman, R. 2007. Response of eelgrass *Zostera marina* to CO<sub>2</sub> enrichment: Possible impacts of climate change and potential for remediation of coastal habitats. *Mar. Ecol. Prog. Ser.* 344:1-13

- Pajusalu, L., Martin, G., Pollumae, A., Paalme, T., 2016. The influence of CO<sub>2</sub> on Net photosynthesis of Seagrass *Zostera marina* in a Brackish water environment. *Front. In Mar. Sci.* 3:1-10
- Rasheed, M.A., Unsworth, R., 2011. Long term climate associated dynamics of a tropical seagrass meadow: implications for the future. *Mar. Ecol. Prog. Ser.* 422:93-103
- Reusch, T.B.H., Ehlers, A., Hammerli, A., Worm, B., 2005. Ecosystem recovery after climatic extremes enhanced by genotypic diversity. *Proceedings National Academy of Science USA*, 102: 2826–2831
- Repolho, T., Duarte, B., Dionisio, G., Paula, J.R., et al., 2017. Seagrass ecophysiological performance under ocean warming and acidification. *Sci. Rep.*7:41443
- Riebesell, U., Fabry, V.J., Hansson, L., Gattuso, J.P., 2010. Guide to best practices for ocean acidification research and data reporting, 260p. Luxembourg: Publications office of the European Union.
- Royal Society., 2005. Ocean acidification due to increasing atmospheric carbon dioxide. London, The Royal Society U.K.
- Rodolfo-Metalpa, R., Houlbrèque, F., Tambutté, E., Boisson, F., et al., 2011. Coral and mollusc resistance to ocean acidification adversely affected by warming. *Nat. Cli. Change.* 1: 308-312.
- Russell, B.D., Connell, S.D., Uthicke, S., Muehllehner, N. et al., 2013. Future seagrass beds: can increased productivity lead to increased carbon storage? *Marine Pollution Bulletin.* 1-7
- Sabina, C.L. and R.A. Feely., 2007. The oceanic sink for carbon dioxide. In *Greenhouse Gas Sinks*. Eds. D. Reay, N Hewitt, J Grace, and K Smith. Oxfordshire: CABI Publishing
- Short, F.T., Neckles, H.A., 1999. The effects of global climate change on seagrasses. *Aq. Bot.* 63:169–196
- Siegenhaller, U. and J.L. Sarmiento., 1993. Atmospheric carbon dioxide and the ocean. *Nature.* 365:119-125
- Seibel, B.A., Walsh, P.J. 2001., Potential impacts of CO<sub>2</sub> injection on deep sea biota. *Science* 294: 319-320
- Shaw, E. C., McNeil, B. I., Tilbrook, B., Matear, R., et al., 2013. Anthropogenic changes to seawater buffer capacity combined with natural reef metabolism induce extreme future coral reef CO<sub>2</sub> conditions, *Glob. Change Biol.* 19:1632–1641
- Solomon S, Qin D, Manning, M, et al., 2007. Technical summary. In: Solomon S, Qin D, Manning M, et al. (Eds). *Climate change: the physical science basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change.* Cambridge, UK, and New York, NY: Cambridge University Press
- Stitt, M., and A. Krapp. 1999. The interaction between elevated carbon dioxide and nitrogen nutrition: the physiological and molecular background. *Plant Cell Environ.* 22:583–621
- Stumm, W., Morgan, J.J., 1981. *An Introduction emphasizing chemical equilibria in natural waters.* Wiley, New York. Aq. Chem. 780 pp

Taiz, L., Zeiger, E., 2002. 3rd ed. Sinauer Associates, Inc., Publishers, Sunderland, MA. *Plant Physiol.*

Takahashi, M., Noonan, S.H.C., Fabricus, K.E., Collier, C.J., 2016. The effects of long term in situ CO<sub>2</sub> enrichment on tropical seagrass communities at volcanic vents. *ICES J. of Mar. Sci.* 73: 876-886.

Thom, R. M., 2001. Carbon Sinks in Nearshore Marine Vegetated Ecosystems. Pacific Northwest Laboratory, Marine Science Laboratory

Vizzini, S., Tomasello, A., Maida, G.D., Pirrotta, M., et al., 2010. Effect of explosive shallow hydrothermal vents on  $\delta^{13}\text{C}$  and growth performance in the seagrass *Posidonia oceanica*. *J. Of Ecol.* 6:1284-1291

Vizzini, S, Di Leonardo, R., Costa, V., Tramati, C.D. et al., 2013. Trace element bias in the use of CO<sub>2</sub> seeps as analogues for low pH environments: Implications for contamination level in acidified oceans. *Estu. coastal and Shelf Sci.* 1-12

Watkiss, P. & Bonjean, M., 2012. Projections of Climate Change and Sea Level Rise for Zanzibar. Watkiss, P. et al. *The Econ. of Cli. Change in Zanzibar.* 1–36

Waycott, M., Duarte, C.M., Carruthers, T.J.B., Orth, R.J., et al. 2009 Accelerating loss of seagrasses across the globe threatens coastal ecosystems. *Proc. of the Nat. Acad. of Sci. of the USA*, 106: 12377–12381

Welsh, D., Bourgues, S., de Wit, R., Auby, I. 1997. Effect of plant photosynthesis, carbon sources and ammonium availability on nitrogen fixation rates in the rhizosphere of *Zostera noltii*. *Aq. Micro. Eco.* 12:285-290

Zimmerman, R.C., Smith, R.D., Alberte, R.S., 1989. Thermal acclimation and whole plant carbon balance in *Zostera marina* L.(eelgrass). *J. Exp. Mar. Biol. Ecol.* 130: 93-109

Zimmerman, R.C., Kohrs, D.G., Steller, D.L., Alberte, R.S., 1997. Impacts of CO<sub>2</sub> enrichment of productivity and light requirements of eelgrass. *Plant Physiol.* 115:599-607



**Short term effects of CO<sub>2</sub> on the diel and seasonal responses of *Zostera noltii* community metabolism to light**



Mishra, A.K., Silva, J., Santos, R.

(To be submitted)



---

### Short term effects of CO<sub>2</sub> on the diel and seasonal responses of *Zostera noltei* community metabolism to light

#### Abstract:

Net community production (NCP) and community respiration (CR) were measured under air exposed and CO<sub>2</sub> enriched conditions for intertidal *Z. noltei* meadows and unvegetated sediment communities during emersion seasonally. Community production and respiration were measured in-situ using benthic chambers under air and CO<sub>2</sub> enriched conditions over a series of short term incubations (30min) using an infra-red gas analyser. Incident photosynthetic active radiation (PAR) was recorded during the incubations covering the daily and seasonal variation. Linear regression model was used to test the effects of irradiance on net community production. NCP of *Z. noltei* community were higher under CO<sub>2</sub> enriched conditions than air exposed conditions in both summer and winter seasons. There was no effect of CO<sub>2</sub> on the CR rate of *Z. noltei* community in summer season. NCP of sediment community were higher in summer season and winter season under CO<sub>2</sub> enriched conditions. Sediment CR rates were higher in winter than summer season. The light compensation point of *Z. noltei* and sediment community were lower in both seasons under CO<sub>2</sub> enriched conditions. Seasonal budget of community production was higher in *Z. noltei* than sediment communities. A clear effect of PAR was noticed on the net community production of both communities. Higher PAR intensities resulted in higher NCP under CO<sub>2</sub> enriched conditions for both communities. CO<sub>2</sub> enrichment can have a positive effect on the intertidal communities during emersion.

Keywords: Intertidal, emersion, unvegetated sediments, regression model, CO<sub>2</sub> enrichment



### 2.1. Introduction

Current concentration of atmospheric carbon dioxide (CO<sub>2</sub>) is predicted to be doubled by end of this century (IPCC, 2007). Rising atmospheric CO<sub>2</sub> will lead to an increase in oceanic CO<sub>2</sub> levels, as 30% of the emitted anthropogenic CO<sub>2</sub> is absorbed in the surface oceans (IPCC, 2014), causing an increased interest of the direct impacts of elevated CO<sub>2</sub> on the marine ecosystem (Gattuso et al., 1998; Feely et al., 2004; Guinotte and Fabry 2008; Hall-Spencer et al., 2008; Portner, 2008; Porzio et al., 2011). The increase of CO<sub>2</sub> in the world oceans is changing the carbonate chemistry of the seawater, lowering the pH resulting in ocean acidification. A further decrease of pH by 0.3–0.4 units is expected by the end of year 2100 (Royal Society, 2005; IPCC, 2014). The low pH of seawater will alter the carbonate chemistry in the relative proportions of the dissolved inorganic carbon species, i.e. carbon dioxide (CO<sub>2</sub>), bicarbonate (HCO<sub>3</sub><sup>-</sup>), and carbonate (CO<sub>3</sub><sup>2-</sup>), shifting the total dissolved inorganic carbon away from CO<sub>3</sub><sup>2-</sup> toward more HCO<sub>3</sub><sup>-</sup> and CO<sub>2</sub> (aq) (Riebesell et al., 2007). This shift toward more HCO<sub>3</sub><sup>-</sup> is expected to benefit species that use it as a carbon source for photosynthesis in addition to CO<sub>2</sub> (Beer et al., 2002; Mercado et al., 2003)

Seagrass ecosystems are distributed worldwide and play an important role in the carbon cycle of coastal areas (Duarte and Chiscano 1999; Hemminga and Duarte 2000). They act as marine carbon sinks and store about 12% of the total carbon storage in the ocean (Duarte et al., 1999, McLeod et al., 2011; Fourqurean et al., 2012). Saying that, the response of seagrass ecosystem to the rising CO<sub>2</sub> levels will be very important, as seagrasses are predicted to benefit from the increasing CO<sub>2</sub> levels (Beer and Koch 1996; Zimmerman et al., 1997; Invers et al., 2001) and overcome their photosynthetic limitation to under current carbon dioxide concentrations.

Most of the studies on elevated CO<sub>2</sub> levels on seagrass have focused on the effects of elevated CO<sub>2</sub> concentrations on seagrass productivity and light requirements (Beer and Koch, 1996; Thom, 1996; Zimmerman et al., 1997; Palacios and Zimmerman, 2007), their photosynthetic and leaf growth rate (Alexandre et al., 2012;) and the uptake rate of ammonium and nitrate (Short and Neckles, 1999; Alexandre et al., 2012; Ow et al., 2016).

However, these experiments of elevated CO<sub>2</sub> levels were carried out more on submerged seagrass ecosystems than air exposed and results have suggested that, seagrass meadows can utilize the increased CO<sub>2</sub> concentration from the water column and enhance their photosynthetic activity and community metabolism, through a regime shift towards utilizing more HCO<sub>3</sub><sup>-</sup> (Frankignoulle and Distèche, 1984; Frankignoulle and Bouquegneau 1990; Invers

et al., 1997). Higher photosynthetic activity of submerged meadows was observed in shallow sites, influenced by light penetration in water column and diurnal variation (Leuschner et al., 1998; Runice and Duarko, 2004). The productivity decreased with increase in depth from shallow sites implying light availability being the important factor for seagrass productivity (Biber et al., 2005) along with other factors such as temperature, diurnal variations and nutrient availability (Lee et al., 2007) in both shallow and intertidal habitats (Jimenez et al., 1987, Kenworthy and Fonseca, 1996; Bach et al., 1998)

However, seagrass meadows in the intertidal habitats, are subjected to pronounced shifts in light and temperature, produced by the combined rhythms of tides and daily irradiance (Short et al., 2001; Vermaat, 2009; Van der Heide et al., 2010). In these intertidal flats, during emersion, light intensity is always higher than during immersion, whereas the nutrient sources are reduced to sediment pore water (Davoult et al. 2011). Intertidal seagrasses are also exposed to varying concentration of inorganic carbon such as gaseous CO<sub>2</sub> during emersion and dissolved CO<sub>2</sub>, HCO<sub>3</sub><sup>-</sup> and CO<sub>3</sub><sup>2-</sup> during immersion (Ouisse et al. 2011; Bahlmann et al., 2015). The seagrass *Zostera noltei* that inhabits the low intertidal zone, experience alternate daily periods of immersion and exposure to air for several hours especially during spring tides (Silva and Santos, 2003; Coyer et al., 2004; Ouisse et al. 2011). In this environment, the *Z. noltei* community have shown considerable photosynthetic activity, when exposed to the air during low tide periods (Leuschner et al., 1998; Silva et al., 2005).

Most of community-level photosynthetic activity estimates of intertidal macrophytes has been assessed from gas exchange methods based on infrared gas analysis (IRGA) of carbon dioxide (CO<sub>2</sub>) fluxes, using flow- through systems or closed chambers for longer periods of time (Streever et al. 1998). Studies of community-level productivity involving CO<sub>2</sub> flux measurements are more common in salt marsh communities than in seagrass meadows. In the former systems, carbon fluxes have been measured either in individual leaves, incubated on special leaf chambers with temperature control and artificial light or in whole plants, using larger incubation chambers (Streever et al., 1998), but less in intact communities. Carbon flux measurements using IRGA in intertidal seagrasses are scarce, particularly on *Zostera noltei*. Most studies using carbon flux measurements on *Z. noltei* are performed in lab conditions (Leuschner and Rees, 1993, Perez-Llorens and Neil, 1994; Leuschner et al. 1998), whereas few measurements have been performed in-situ (Leuschner et al., 1998; Silva et al., 2005; Ouisse et al., 2010, 2011). So, in this context, we aim to measure the effects of high CO<sub>2</sub> concentration (future scenarios of 750 ppm) on the net community production (NCP) and community

respiration (CR) of air exposed *Z. noltei* community at Ria Formosa Lagoon, Portugal, where the species dominates the intertidal zone. The effects of vegetation on the intertidal ecosystem metabolism will be assessed by comparison with unvegetated areas.

### 2.2. Materials and methods

#### *Study sites:*

Ria Formosa lagoon (37°00'N, 7°58' W) is a mesotidal system located in the southern coast of Portugal, characterized by large intertidal flats (Andrade, 1990). *Zostera noltei* is the most abundant seagrass species in Ria Formosa coastal lagoon in the lower intertidal zone, distributed within a bathymetric range of about 2 m and plays a major role in the lagoon's metabolism (Santos et al. 2004). The *Z. noltei* meadows regularly experience alternate daily periods of immersion and exposure (Site A, Fig 1) withstanding long periods of air exposure up to 4.5 to 6 hours in each tidal cycle. The unvegetated sediment beds are at the same bathymetric zone experiencing the daily tidal fluctuations. These beds are located on the north-west direction of seagrass beds approximately 600 m in distance (Site B, Fig 1).

#### *Gas Exchange Measurements:*

Measurements were made in-situ, through carbon flux measurements for short durations, using Infrared gas analyser (IRGA) and Plexiglas chambers, with respect to light variations daily and seasonally and the community response of *Z. noltei* was compared with adjacent un-vegetated sediments. This experiment was carried out in summer (June, July, August) 2014 and winter (November, December, January) 2014-15 seasons. Incubation chambers made of UV-transparent Plexiglas, providing closed environment were used for incubations. These chambers comprised of two separated parts, a basal ring (30 cm in diameter) and a dome-shaped lid to close the chamber. The basal ring was sharpened for easier sediment insertion, usually down to 5–10 cm deep. Once the lid was set in place, both parts were secured by power clamps (Fig. 2). A special latex ring installed in a flange insures the chamber was sealed. The chambers were provided with three ports, one in the ring (2), one on the top of the dome (1) and the third (3) on the base 5-10 cm above the above the sediment insertion line for the CO<sub>2</sub> gas to be fluxed into the chamber. All ports were fitted with Tygon® ST gas-impermeable tubing. The global operating volume of the chambers was between 22 and 26 L, depending on how deep the ring was inserted into the sediment.

Short term field incubations using these chambers were conducted in *Z. noltei* meadows and bare sediment beds during several low tides at different times of day along the season, so that the diel variation of light was captured. Each incubation lasted for a maximum of 30 minutes. Each incubation was first conducted under natural atmospheric CO<sub>2</sub> concentration (397-399 ppm) followed by CO<sub>2</sub> enrichment to 750 ppm, of the same *Z. noltei* community. During the incubation air entered the chamber through the ring port (2) and was out through the dome port (1), allowing the homogenization of air within the chamber medium. A low-volume (approx. 25 ml) closed air circuit is established between the chamber and an external non-dispersive infrared gas analyser (IRGA) (EGM-4, PP Systems, UK) that measures CO<sub>2</sub> concentrations with a precision of 1 µatm. The gas analyser had a built-in pump that pulled the air from the chamber, passed through a desiccation column (Anhydrous calcium sulphate, Drierite) which removed humidity from the air to avoid interferences in CO<sub>2</sub> measurements and sends it back after passing through the infrared cell, keeping a continuous air flow between the incubation chamber and the analyser. The CO<sub>2</sub> partial pressure inside the chamber is then continuously monitored (1 min steps) over a period, typically between 5 and 15 min for each incubation, or enough for a 30–40 µatm drop in the CO<sub>2</sub> partial pressure. During this time the port (3) was clamped, so that no air passes through the tubes. After completion of the air incubation, the clamp was removed and CO<sub>2</sub> gas was fluxed slowly into the chamber through the port (3), from a CO<sub>2</sub> gas enriched tank till the CO<sub>2</sub> concentration inside the chamber reached 750 ppm, recorded by IRGA. The clamp was closed again. The CO<sub>2</sub> partial pressure was measured, same as described previously. Following each incubation, the seagrass was collected, washed and dried at 60° C for 24 hours and the dry weight was recorded. Photosynthetic active radiation (PAR) inside the chambers was measured with a Li-Cor system (Li-Cor, 250 A meter, USA) at both the start and end of each incubation. Instantaneous PAR measurements were continuously done at the Ramalhete field station (Fig.1) every 30-minute using the same sensor to record daily and seasonal light variation. Light attenuation inside the chambers was estimated to be approximately 10%. Temperature was recorded using a thermometer at start and end of each air and CO<sub>2</sub> incubation (Fig. 2). Similar incubations were also performed for unvegetated sediment beds.

---

*NCP and CR measurements*

Light and dark incubations were conducted for both net community production (NCP) and community respiration (CR) respectively. The net flux (F) of CO<sub>2</sub> (μmol C m<sup>-2</sup> h<sup>-1</sup>) for both air and CO<sub>2</sub> enriched incubations was computed as:

$$F = s \times mv \times V/A \times 60/1000$$

where  $s$  (ppm CO<sub>2</sub> min<sup>-1</sup> = μmol CO<sub>2</sub> mol air<sup>-1</sup> min<sup>-1</sup>) is the slope of the linear change in CO<sub>2</sub> concentration over time during the incubation period,  $V$  (L) is the chamber volume,  $A$  (m<sup>2</sup>) is the benthic exchange area, and  $mv$  (mol L<sup>-1</sup>) is the molar volume or number of gas mole per volume unit, calculated from the ideal gas law ( $mv = P/(R \times T)$ ), where  $P$  (atm) is the gas pressure,  $T$  (K) the temperature, and  $R$  (0.082 atm LK<sup>-1</sup> mol<sup>-1</sup>) is the universal gas constant. Thus, the net CO<sub>2</sub> flux will be a direct measure of the NCP for the given light and temperature conditions. Similarly, the net flux of CO<sub>2</sub> in dark represents the community respiration, CR.

A linear regression model was fitted to the NCP versus PAR data to assess the effects of PAR on NCP,

$$Y = -b x + a,$$

Where  $Y$  is NCP,  $-b$  represent the rate of CO<sub>2</sub> uptake by the community,  $X$  is the PAR value and 'a' intercept, representing community respiration. The daily PAR variation reaching the *Z. noltei* community when exposed to the air, was used to estimate the daily NCP variation using the above model. Tidal variations (emersion time) for the study site (*Z. noltei* meadows and sediment beds) for each season (summer and winter) was estimated using a tidal model (Carrasco et al. unpublished data). The emersion time derived from the model for each day along the month and seasons for seagrass meadows and sediment beds was used to calculate the daily NCP, GCP and CR for each season.

Linear regression was used to model the effects of PAR on NCP. The summer and winter NCPs were calculated based on the average daily PAR of the period the plants were exposed to the air and the linear models of NCP response to irradiance. Seasonal NCP (g C m<sup>-2</sup> season<sup>-1</sup>) is the average of all summer or winter days.

The daily NCP and CR during emersion was estimated by adding every 30 minute NCP and CR values along the day when the seagrass beds were exposed to the air. Gross community production (GCP) was obtained by adding the NCP and CR during light hours. The seasonal NCP, CR and GCP were obtained by considering the variability of exposure times related to the alternation of spring and neap tidal cycles and the daily cycle of irradiance.

Results are expressed as mean  $\pm$  Standard error (S.E). A t-test was used to detect the significant differences of NCP, GCP and CR between natural and CO<sub>2</sub> enriched conditions. Data was tested for normality and constant variance before testing for significant difference. Significant levels were tested at  $p < 0.05$ .

### 2.3. Results

NCP of *Zostera noltei* and sediment communities were higher under CO<sub>2</sub> enriched conditions than natural conditions in both summer and winter seasons (Fig. 3). The differences were only apparent at light intensities above the light compensation for community production (higher than 200  $\mu\text{mol m}^{-2} \text{s}^{-1}$ ), i.e. when production is higher than respiration. NCP of sediments was lower than *Z. noltei* within PAR ranges of 800  $\mu\text{mol m}^{-2} \text{s}^{-1}$  to 1600  $\mu\text{mol m}^{-2} \text{s}^{-1}$ , which was most evident in the summer (Fig. 4). In the summer season under higher light (1680  $\mu\text{mol m}^{-2} \text{s}^{-1}$ ) the NCP of sediments was higher than *Z. noltei* under natural conditions, whereas under CO<sub>2</sub> enriched conditions the NCP of *Z. noltei* was higher than unvegetated sediment community (Fig. 4). Mean plant biomass of *Z. noltei* was  $1.80 \pm 0.89 \text{ g DW m}^{-2}$  in summer and  $1.68 \pm 0.80 \text{ g DW m}^{-2}$  in winter season.

Significant differences were observed in the net community production of *Z. noltei* and sediment communities under natural and CO<sub>2</sub> enriched conditions (Fig. 5). The NCP of *Z. noltei* was higher in both seasons under CO<sub>2</sub> enriched conditions and community respiration was lower and consequently gross community production was even higher under CO<sub>2</sub> enriched conditions than natural conditions (Fig. 5a). Similarly, under CO<sub>2</sub> enriched conditions sediment NCP was higher in both season, whereas the CR was higher in summer and lower in winter (Fig. 5b). GCP of sediment community was higher in summer and lower in winter season under CO<sub>2</sub> enriched conditions (Fig. 5b).

Seasonal budget of *Z. noltei* and unvegetated sediment community were higher under CO<sub>2</sub> enriched conditions than natural conditions in both seasons (Table 1). However, seasonal budget of NCP for *Z. noltei* was 3.9 -fold higher and seasonal GCP 4.9-fold higher in summer season than winter season under CO<sub>2</sub> enriched conditions (Table 1). In case of sediment communities, the increase of seasonal budget of NCP was 3.5-fold in summer season, whereas the decrease of GCP was 7.5-fold in winter season under CO<sub>2</sub> enriched conditions (Table 1). Similar seasonal budget of NCP for *Zostera noltei* and sediment community was observed in winter season under CO<sub>2</sub> enriched conditions, whereas seasonal GCP of *Z. noltei* was higher

than sediments communities (Table 1). Seasonal budget of CR was higher in sediment communities than *Z. noltei* under CO<sub>2</sub> enriched conditions in winter season (Table 1).




Fig. 1. Study area showing *Zostera noltei* beds and unvegetated sediment beds at Ria Formosa Lagoon.




Fig. 2. A diagrammatic representation of the incubation work. 1. Port one on top of dome. 2. Port two on ring. 3. Port three for CO<sub>2</sub> flux




Fig. 3. Response of net community production (NCP) to irradiance (PAR) under natural and CO<sub>2</sub> enriched conditions in summer and winter for *Z. noltei*, (a, b) and sediments (c, d) obtained from linear regression.




Fig. 4. Comparison of community production and respiration of *Z. nolteii* and sediment community for summer (a, b) and winter seasons (c, d) under natural (air) and  $\text{CO}_2$  enriched conditions




Fig. 5. Mean ( $\pm$  se) community production and respiration ( $\text{mg C m}^{-2} \text{ day}^{-1}$ ) of *Z. noltei* (a, b) and unvegetated sediment community (c, d) in summer and winter seasons under natural and CO<sub>2</sub> enriched conditions. Different letters indicate significant difference between natural (air) and CO<sub>2</sub> enriched conditions. Error bars represent standard errors

Table 1.

Seasonal budget of NCP ( $\text{g C m}^{-2} \text{ season}^{-1}$ ), CR ( $\text{g C m}^{-2} \text{ season}^{-1}$ ) and GCP ( $\text{g C m}^{-2} \text{ season}^{-1}$ ) of *Z. noltei* and unvegetated sediment community under natural (air) and  $\text{CO}_2$  enriched conditions in summer and winter seasons

Season	Condition	<i>Z. noltei</i> community		
		NCP	CR	GCP
Summer	Air	78.21 $\pm$ 5.58	16.99 $\pm$ 0.85	59.56 $\pm$ 5.19
	$\text{CO}_2$	115.54 $\pm$ 7.58	12.12 $\pm$ 0.67	98.65 $\pm$ 7.44
Winter	Air	20.13 $\pm$ 3.02	26.86 $\pm$ 1.52	12.15 $\pm$ 2.30
	$\text{CO}_2$	29.45 $\pm$ 4.04	12.30 $\pm$ 0.74	19.91 $\pm$ 3.67
<b>Sediment community</b>				
Summer	Air	72.26 $\pm$ 5.08	13.24 $\pm$ 0.74	56.51 $\pm$ 4.87
	$\text{CO}_2$	91.28 $\pm$ 6.32	14.55 $\pm$ 0.81	61.57 $\pm$ 5.94
Winter	Air	19.78 $\pm$ 2.98	24.36 $\pm$ 1.50	8.75 $\pm$ 1.99
	$\text{CO}_2$	25.83 $\pm$ 4.01	18.45 $\pm$ 1.27	8.60 $\pm$ 3.54

Significant effect of PAR was observed on the NCP of *Z. noltei* and unvegetated sediments under natural and  $\text{CO}_2$  enriched conditions in summer and winter seasons (Table 2). The slope, intercept and  $R^2$  values of the linear regression model for both conditions and seasons are presented in Table 2. The light compensation point (LCP) were lower under  $\text{CO}_2$  enriched conditions than natural conditions for *Z. noltei* and sediment community (Table 2). LCP of *Z. noltei* community was lower than unvegetated sediment community in both seasons under  $\text{CO}_2$  enriched conditions (Table 2). Highest PAR recorded during the field experiments was 1840.45 to 1895.56  $\text{mol m}^{-2} \text{ s}^{-1}$  in summer season, and between 1522  $\mu \text{mol m}^{-2} \text{ s}^{-1}$  to 1594  $\mu \text{mol m}^{-2} \text{ s}^{-1}$  in winter season for both communities (Fig. 3).

Table 2.

Slope (b), intercept (a), p, R<sup>2</sup> and light compensation point (LCP,  $\mu\text{ mol m}^{-2}\text{ s}^{-1}$ ) values derived from linear regression model between PAR and NCP under natural and CO<sub>2</sub> enriched conditions for *Z. noltei* community and unvegetated sediments community.

Season	Condition	b	<i>Z. noltei</i> community		R <sup>2</sup>	LCP
			a	PAR vs NCP (p value)		
Summer	Air	0.0069	-1.622	<0.001	0.87	247.50
	CO <sub>2</sub>	0.0092	-1.233	<0.001	0.95	134.02
Winter	Air	0.0064	-1.796	<0.001	0.96	280.65
	CO <sub>2</sub>	0.0068	-0.882	<0.001	0.90	129.70
<b>Sediment community</b>						
Summer	Air	0.0062	-1.281	<0.001	0.95	206.61
	CO <sub>2</sub>	0.0077	-1.436	<0.001	0.90	186.49
Winter	Air	0.0062	-1.707	<0.001	0.89	275.32
	CO <sub>2</sub>	0.0070	-1.391	<0.001	0.86	198.71

### 2.4. Discussion

In our study, the linear regression model used to test the effect of irradiance on the net community production showed a significant correlation with both natural and CO<sub>2</sub> enriched conditions. The high coefficient of determination ( $R^2$ ) in our datasets under natural and CO<sub>2</sub> enriched conditions (Table 1) indicates a closed fit of the model to the data, suggesting irradiance is key driver of NCP. The NCP followed a linear pattern of increase with increasing PAR variation and CO<sub>2</sub> enrichment, indicating higher CO<sub>2</sub> concentrations with higher irradiance levels will increase NCP of *Z. noltei* and sediment communities under future high CO<sub>2</sub> conditions. Lower light compensation point obtained from the model suggests that high CO<sub>2</sub> will compensate for low light and increase the carbon utilizing capacity of *Z. noltei* during low light conditions. The linear increasing trend of community production versus irradiance observed in our model differs from irradiance versus net community production models observed for *Z. noltei* elsewhere (Cavier et al., 2011, 2014) and unvegetated sediment community (Uthicke et al., 1998; Cavier et al., 2014).

The light compensation point derived from the model showed lower LCP for CO<sub>2</sub> enriched conditions and higher under natural conditions. The LCP for *Z. noltei* was in between 134  $\mu\text{mol quanta m}^{-2} \text{s}^{-1}$  to 247  $\mu\text{mol quanta m}^{-2} \text{s}^{-1}$  for summer season and between 129  $\mu\text{mol quanta m}^{-2} \text{s}^{-1}$  to 280  $\mu\text{mol quanta m}^{-2} \text{s}^{-1}$  for winter season under CO<sub>2</sub> enriched and natural conditions respectively (Table 1). The LCP observed in our results were higher than previously reported for *Z. noltei* for both seasons under natural conditions (Gacia et al., 2005; Ouisse et al., 2010) and lower for CO<sub>2</sub> enriched conditions. The LCP for sediment community were between 186  $\mu\text{mol quanta m}^{-2} \text{s}^{-1}$  to 275  $\mu\text{mol quanta m}^{-2} \text{s}^{-1}$  for both seasons under CO<sub>2</sub> enriched and natural conditions, which were higher than previously observed for sediment communities (Uthicke & Klumpp, 1998; Clavier & Garrigue, 1999). The light compensation point of *Z. noltei* values for both summer and winter seasons suggest an effect of light on net community production for both *Z. noltei* and sediment community in the Ria Formosa. The LCP for CO<sub>2</sub> enriched experiments during emersion are presented for the first time for *Z. noltei* and sediment community.

Measurements of NCP vs PAR during emersion highlighted the continuous increase in net community production with PAR variation for both conditions in summer season, similar results were obtained for *Z. noltei* with no decrease in community production under natural conditions (Ouisse et al., 2010, 2011; Clavier et al., 2011). The increase in community

production at higher light intensities can be related to tolerance of high light intensities by *Z. noltei* without damaging its pigment pool and dynamically down regulating the photosynthetic apparatus (Jimenez et al., 1987). This phenomenon was also observed in *Z. marina* (Zimmerman et al., 1997) and *Z. noltei* (Alexandre et al., 2012) where increase in CO<sub>2</sub> concentrations increased the light utilizing capacity of individual plants.

The lower range of LCP in CO<sub>2</sub> enriched conditions and higher net community production for *Z. noltei* in summer suggests that the community is more efficient in utilizing low light (Silva & Santos, 2003) and can start utilizing CO<sub>2</sub> for production in low light conditions without community production compensating for community respiration, similar pattern was observed for *Z. marina* in natural conditions (Ouisse et al., 2011). In case of sediment community, the LCP were higher than *Z. noltei* under CO<sub>2</sub> enriched conditions, but the community production was not, which suggests that even with higher CO<sub>2</sub> concentrations the microphytobenthos can't utilize carbon still a certain level of irradiance level is reached.

Net community production and community respiration of sediment community was lower than *Z. noltei* community in both summer and winter seasons under natural conditions (Fig. 5). Similar results were observed for *Z. noltei* community and sediment community at Bay of Morlaix (Ouisse et al., 2010) and Banco de Mauritania (Clavier et al., 2014).

The short-term CO<sub>2</sub> enrichment in our studies resulted in increase of net community production of *Z. noltei* community along the daily variation of PAR. During summer, the net community production of *Z. noltei* were higher and community respiration rates lower in CO<sub>2</sub> enriched conditions. This study showed that net community production of *Z. noltei* is affected by CO<sub>2</sub> enrichment and that PAR variation along the day and season plays an important role in the community production. Seagrass community exposed to CO<sub>2</sub> enriched conditions showed higher community production and were more efficient at low light intensities, when compared with seagrass community exposed to the current CO<sub>2</sub> concentration.

For summer season, NCP of *Z. noltei* community in air exposed community was higher than previously reported for *Z. noltei* and *Z. marina* air exposed community (Ouisse et al., 2011), whereas the CR observed in our results under natural conditions was lower than *Z. marina* community and higher than *Z. noltei* community response observed by Ouisse et al., (2011). Similarly, NCP under CO<sub>2</sub> enrichments experiments was higher than results reported for air exposed *Z. noltei* and *Z. marina* community (Ouisse et al., 2010, 2011; Clavier et al., 2014).

For summer season, GCP under natural conditions were lower to GCP obtained for *Z. noltei* community and higher to *Z. marina* community (Ouisse et al., 2011). Similarly, GCP under CO<sub>2</sub> enrichment was higher than values observed for air exposed *Z. noltei* and *Z. marina* community (Ouisse et al., 2011; Clavier et al., 2014). The GCP and CR of *Z. noltei* observed under air exposed conditions in winter season, were higher than previously reported for *Z. noltei* community by Clavier et al., (2011,2014) and Ouisse et al., (2011). Higher community production can be due to higher irradiance (1522 to 1594  $\mu\text{mol m}^{-2}\text{ s}^{-1}$ ) levels observed in our studies than irradiance (829  $\mu\text{mol m}^{-2}\text{ s}^{-1}$ ) level observed by Ouisse et al., (2011) contributing to the community production and higher respiratory demand of the extended rhizome network of *Z. noltei* which requires higher production (Vermaat et al., 1987). Gross community production of *Z. noltei* communities were higher than previously reported by Silva et al., (2005) in Ria Formosa, under natural and CO<sub>2</sub> enriched conditions.

The community respiration of *Z. noltei* community under natural conditions was higher in winter than summer season (Fig.5), opposite trend of higher respiration rates in summer than winter was observed for *Z. noltei* community elsewhere (Plus et al., 2005; Ouisse et al., 2010). Higher community respiration rates in *Z. noltei* in winter can be due to low irradiance levels observed in our studies, followed by low community production, thus community respiration takes over the community production with CO<sub>2</sub> release and oxygen consumption, similar pattern of higher respiration was observed in *Z. marina* elsewhere (Rasmusson, 2015).

Community production under natural conditions during emersion always varies with the community production during immersion. Higher community production during emersion than immersion was observed in the same study site for *Z. noltei* (Silva et al., 2005), whereas community production of *Z. noltei* lower during emersion and higher during immersion was observed elsewhere (Ouisse et al., 2011; Clavier et al., 2011).

The short-term gas exchange measurements with CO<sub>2</sub> enrichment conditions for *Z. noltei* community clearly indicated that community production (NCP and GCP) around similar irradiance level increased 1.5-fold and 1.6-fold respectively under CO<sub>2</sub> enriched conditions than natural conditions (Fig. 5). Higher community production under CO<sub>2</sub> enrichment suggest that community production of *Z. noltei* are currently carbon limited under the current atmospheric CO<sub>2</sub> concentration, confirming the same conclusions previously obtained for *Z. noltei* (Alexandre et al., 2012) and for *Z. marina* (Beer and Koch, 1996; Zimmerman et al., 1997; Invers et al., 2001). This results also suggest that *Z. noltei* community may benefit from

future CO<sub>2</sub> enrichment by enhancing their net photosynthetic rates at higher CO<sub>2</sub> concentrations. The CO<sub>2</sub> stimulated increase in community production found here for *Z. noltei* is consistent with the findings reported for the temperate and tropical seagrass species, where positive community production responses to CO<sub>2</sub> enrichment were found *Z. marina* (Beer & Koch, 1996; Thom, 1996; Zimmerman et al., 1997) and *Thalassia hemprichii* (Jiang et al., 2010), whereas no response to CO<sub>2</sub> enrichment was observed in *Halophila uninervis* (Ow et al., 2016).

The key factor during emersion for higher production under CO<sub>2</sub> enriched conditions can be higher light availability during emersion periods and abundant CO<sub>2</sub> as substrate from the CO<sub>2</sub> enrichment. Increase in CO<sub>2</sub> as substrate resulting in higher production efficiency was observed in *Z. marina* where elevated CO<sub>2</sub> levels increased plant production (Zimmerman et al., 1997). Secondly sediment microtopography of *Z. noltei* community plays an important role in maintaining hydration for the whole emersion period, due to numerous small depressions in the sediment retaining water (Silva et al., 2005). Thirdly during emersion, a thin layer of water is trapped between the leaves of *Z. noltei* due to higher leaf density (Silva & Santos, 2003), helping the plants remain moist throughout the emersion and this thin layer of water also helps in transfer of CO<sub>2</sub> from the atmosphere to the leaf surface (Leuschner and Rees, 1993), whereas in our case this transfer of CO<sub>2</sub> takes place inside the chamber from CO<sub>2</sub> enrichment. Simultaneously, the desiccations in *Z. noltei* meadows are not severe (Silva & Santos, 2003) and the *Z. noltei* root system also supplies water to the leaves, which helps in maintaining a thin layer of water at the leaf surface (Leuschner et al., 1998). In these favourable conditions when CO<sub>2</sub> enrichment happens, a rapid diffusion of CO<sub>2</sub> across the thin film of water around the seagrass leaves occurs due to air-water CO<sub>2</sub> gradient, making the CO<sub>2</sub> readily available to the plants (Leuschner et al., 1998). Therefore, high CO<sub>2</sub> concentrations along with high irradiance levels during air exposure of the *Z. noltei* community, creates the ideal condition for high community production during emersion (Leuschner & Rees, 1993; Silva et al., 2005; Touchette & Burkholder, 2007).

Net community production of sediment community (785 mg C m<sup>-2</sup> day<sup>-1</sup>) in our studies under natural conditions in summer was higher than previously reported NCP (681.5 mg C m<sup>-2</sup> day<sup>-1</sup>) in natural at Tagus estuary, Portugal in summer season (Uthicke et al., 1998), whereas NCP (212.52 mg C m<sup>-2</sup> day<sup>-1</sup>) of sediment community in winter season in our results was lower than previously reported NCP (255 mg C m<sup>-2</sup> day<sup>-1</sup>). Average NCP and CR observed in our

results for sediment community in winter season were higher than NCP and CR rates observed in the intertidal sediment community at Banco di Marituania (Clavier et al., 2014). Sediment community NCP were higher with CO<sub>2</sub> enrichment incubations in summer and winter than air incubations, which can be related to the presence of micro-phytobenthos in the sediments during intertidal exposure (Uthicke & Klumpp, 1998; Hubas et al., 2006; Clavier et al., 2014). The intertidal desiccation of the sediments was avoided due to the presence of a thin film of water trapped by the benthos community and water trapped by intertidal burrows (Michael & McIntire, 1983).

Seasonal (summer and winter) variation of GCP (614.23 to 93.64 mg C m<sup>-2</sup> day<sup>-1</sup>) were higher than seasonal GCP (38 to 151 mg C m<sup>-2</sup> day<sup>-1</sup>) reported by Hubas et al., (2006). GCP of unvegetated sediment communities were lower in winter season than summer under CO<sub>2</sub> enriched conditions, due to higher community respiration rates (Fig.5). Higher community respiration can be related to the presence of other bacteria and infauna not contributing to the community production but respiration (Uthicke & Klumpp, 1998). Secondly lower irradiance intensities in winter means, less penetration of light into the sediment resulting in decrease of microphytobenthos biomass in the sediment (Pinckney & Zingmark, 1993b) and finally the presence of large content of detritus on the unvegetated sediment surface presents a shading effect on the fewer micro-phytobenthos community present in winter season (Ouisse et al., 2010). The incident PAR was also influenced by the length of the day in winter (11-hour daylight compared to 13-hour daylight in summer) and cloud cover which could have also affected the sediment community production (Parsons et al., 1984; Migné et al., 2004; Hubas et al., 2006).

Overall the short-term incubation method based on CO<sub>2</sub> enrichment using benthic chambers described here is reported for the first time for intertidal *Z. noltei* community. This method can be an important tool for intertidal lagoon community productivity. It provides fast and precise values of community carbon fixation during emersion periods for air exposed plants and sediment community. The use of small chambers and short-term incubation periods is an important feature in these experiments as the temperature and humidity levels remain constant. This method also allows repeatable in situ measurements of carbon uptake with minimum community disturbance. For these reasons, we recommend the use of this method to assess the contribution of different communities to the metabolism and carbon budget of complex tidal systems.

The results of a good correspondence observed from the linear regression model between carbon metabolism measurements of the *Z. noltei* community and sediment community with respect to light intensities across summer and winter seasons, provides evidence that community production rates derived from CO<sub>2</sub> enriched incubations are reflection of photosynthetic activity at ecologically relevant scales for future conditions of carbon dioxide. Positive response to elevated CO<sub>2</sub> levels and lower light requirements as observed in our results may at least partially offset negative responses to decreased light availability from sea level rise associated with global climate change.

### 2.5. References:

- Alexandre, A, Silva, J., Buapet, P., Bjork, M., Santos, R., 2012. Effects of CO<sub>2</sub> enrichment on photosynthesis, growth and nitrogen metabolism of the seagrass *Zostera noltei*. *Ecology and Evolution*. 2: 2620-2630.
- Andrade, C.F., 1990. O ambiente barreira da Ria Formosa, Algarve, Portugal. PhD Thesis, University of Lisbon, Lisbon
- Bach, S.S., Borum, J., Fortes, M.D., Duarte, C.M., 1998. Species composition and plant performance of mixed seagrass beds along a siltation gradient at Cape Bolinao, The Philippines. *Mar. Ecol. Prog. Ser.* 174: 247–256.
- Bahlmann, E., Weinberg, I., Lavric, V.J., Eckhardt, T., Michaelis, W., et al., 2015. Tidal controls on trace gas dynamics in a seagrass meadow of the Ria Formosa lagoon (Southern Portugal). *Biogeosciences*. 12:1683-1696
- Beer, S., Koch, E., 1996. Photosynthesis of marine macroalgae and seagrasses in globally changing CO<sub>2</sub> environments. *Mar. Ecol. Prog. Ser.* 141:199–204
- Beer, S., Bjork, M., Hellblom, F., Axelsson, L., 2002. Inorganic carbon utilization in marine angiosperms (seagrasses). *Funct. Plant Biol.* 29: 349–354.
- Biber, P.D., Paerl, H.W., Gallegos, C.L., Kenworthy, W.J., 2005. Evaluating indicators of seagrass stress to light. In: Bortone, S.A. (Ed.), *Estuarine Indicators*. CRC Press, Boca Raton, FL, pp. 193-210
- Bjork, M., Uku, J., Weil, A., Beer, S., 1999. Photosynthetic tolerances to desiccation of tropical intertidal seagrasses. *Mar. Ecol. Prog. Ser.* 191: 121–126.
- Caldeira, K., and Wickett, M.E., 2003. Oceanography: anthropogenic carbon and ocean pH. *Nature*. 425:365.
- Carrasco, A.R., Plomaritis, H., Reynolds, J., Ferreira, O., Roelvink, D. Tide circulation patterns under sea -level rise scenarios: Ria Formosa coastal lagoon. *Estu. Coastal and Shelf Sci.* (in revision)

- Clavier, J., Chauvaud, L., Carlier, A., Amice, E., Geest, M., et al., 2011. Aerial and underwater carbon metabolism of a *Zostera noltei* seagrass bed in the Banc d'Arguin, Mauritania. *Aq. Bot.* 1: 24-30
- Clavier, J., Chauvaud, L., Amice, E., Lazure, P., Geest, M., et al., 2014. Benthic metabolism in shallow coastal ecosystems of the Banc d'Arguin, Mauritania. *Mar. Ecol. Prog. Ser.* 501: 11-23
- Coyer, J.A., Diekmann, O.E., Serrao, E.A., Procaccini, G., Milchakova, N., et al., 2004. Population genetics of dwarf eelgrass *Zostera noltei* throughout its biogeographic range. *Mar. Ecol. Prog. Ser.* 281: 51–62.
- Duarte, C. M. 1990. Seagrass nutrient content. *Mar. Ecol. Prog. Ser.* 67:201–207.
- Duarte, C. M., and C. L. Chiscano. 1999. Seagrass biomass and production: a reassessment. *Aq. Bot.* 65:159–164
- Enríquez, S., Merino, M., Iglesias-Prieto, R., 2002. Variations in the photosynthetic performance along the leaves of the tropical seagrass *Thalassia testudinum*. *Mar. Biol.*140: 891-900
- Fourqurean, J.W., Duarte, C.M., Kennedy, H., Marba, N, Holmer, M. et al., 2012. Seagrass ecosystem as a globally significant carbon stock. *Nat. Geosci.* 5:505-509.
- Frankignoulle, M. and Bouqueneau J. M.,1990. Daily and yearly variations of total inorganic carbon in a productive coastal area, *Estuar. Coast. Shelf Sci.*30: 79–89
- Frankignoulle, M. and Disteche, A., 1984. CO<sub>2</sub> chemistry in the water column above a *Posidonia* seagrass bed and related air-sea exchanges, *Oceanol. Ac.*7: 209–219
- Feely, R. A., Sabine, C.L., Lee, K., Berelson, W., Kleypas, J., et al. 2004. Impact of anthropogenic CO<sub>2</sub> on the CaCO<sub>3</sub> system in the oceans. *Science.* 305:362–366
- Gacia, E., Duarte, C.M., Middelburg, J.J., 2002. Carbon and nutrient deposition in a Mediterranean seagrass (*Posidonia oceanica*) meadow. *Limnol. Oceanogr.*47:23-32
- Gacia, E, Kennedy, H., Duarte, C.M., Terrados, J., Marba, N., et al., 2005. Light dependence of the metabolic balance of a highly productive Philippine seagrass community. *J. of Expt. Mar. Biol. and Ecol.* 316: 55-67.
- Gazeau, F., Quiblier, C., Jansen, J. M., Gattuso, J. P., Middelburg, J. J., et al., 2007: Impact of elevated CO<sub>2</sub> on shellfish calcification, *Geophys. Res. Lett.*, 34.
- Gazeau, F., Duarte, C., Gattuso, J., Barron, C., Navarro, N., et al. 2004. Whole system metabolism and CO<sub>2</sub> fluxes in a Mediterranean Bay dominated by seagrass beds (Palma, Bay, NW Mediterranean). 1: 755-802
- Gattuso, J.P., Frankignoulle, M., Bourge, I., Romaine, S., Buddemeier, R. W., 1998. Effect of calcium carbonate saturation of seawater on coral calcification. *Glob. Planet. Change.* 18: 37–46.
- Guinotte, J. M., and Fabry, V.J., 2008. Ocean acidification and its potential effects on marine ecosystems. *Ann. N. Y. Acad. Sci.* 1134:320–342.

- Hall-Spencer, J. M., Rodolfo-Metalpa, R., Martin, S., Ransome, E., Fine, M., et al., 2008. Volcanic carbon dioxide vents show ecosystem effects of ocean acidification. *Nature*. 454:96–99
- Hemminga, M. A., and Duarte, C.M., 2000. *Seagrass ecology*. Cambridge Univ. Press, Cambridge, U.K.
- Hubas, C., Davoult, T., Cariou, T., 2006. Factors controlling benthic metabolism during low tide along a granulometric gradient in an intertidal bay (Roscoff Aber Bay, France). *Mar. Ecol. Prog. Ser.* 316: 53-68.
- Invers, O., Zimmerman, R.C., Alberte, R.S., Perez, M., Romero, R., 2001. Inorganic carbon sources for seagrass photosynthesis: an experimental evaluation of bicarbonate use in species inhabiting temperate waters. *J. Exp. Mar. Biol. Ecol.* 265:203–217
- IPCC (Intergovernmental Panel on Climate Change). 2007. *Climate change 2007 synthesis report*. Cambridge Univ. Press, New York.
- IPCC. 2014. *Fifth Assessment Report on Climate change: Impacts, Adaptation and Vulnerability*.
- Jiménez, C., Niell, F.X., Algarra, P., 1987. Photosynthetic adaptation of *Zostera noltei* Hornem. *Aq. Bot.* 29:217-26
- Jacques. C, Garrigue, C.,1999. Annual sediment primary production and respiration in a large coral reef lagoon. *Mar. Ecol. Prog. Ser.* 191:79-89.
- Jiang, Z. J., Huang, X.P., Zhang, J.P., 2010. Effects of CO<sub>2</sub> enrichment on photosynthesis, growth and biochemical composition of seagrass *Thalassia hemprichii* (Ehrenb.) Aschers. *J. Integr. Plant Biol.* 52: 904–913
- Jiménez, C., Niell, F.X., Algarra, P., 1987. Photosynthetic adaptation of *Zostera noltei* Hornem. *Aq. Bot.* 29: 217–226
- Khan, F., Alexandre, A., Ullah, H., Santos, R., 2016. Effects of elevated CO<sub>2</sub> and nutrients on the community metabolism of a *Cymodocea nodosa* bed. *Turk. J. of Bot.* 3: 250-257
- Kenworthy, W.J., Fonseca, M.S., 1996. Light requirements of seagrasses *Halodule wrightii* and *Syringodium filiforme* derived from the relationship between diffuse light attenuation and maximum depth distribution. *Estuaries.* 19:740–750
- Leuschner, C., Rees, U., 1993. CO<sub>2</sub> gas exchange of two intertidal seagrass species, *Zostera marina* L. and *Zostera noltei* Hornem., during emersion. *Aq. Bot.* 45: 53–62.
- Leuschner, C., Landwehr, S., Mehlig, U., 1998. Limitation of carbon assimilation of intertidal *Zostera noltei* and *Zostera marina* by desiccation at low tide. *Aq. Bot.* 62: 171-176
- Leuschner, C., Landwehr, S., Mehlig, U., 1998. Limitation of carbon assimilation of intertidal *Zostera noltei* and *Z. marina* by desiccation at low tide. *Aq. Bot.* 62: 171–176.
- Lee, K.S., Park, S.R., Kim, Y.K., 2007. Effects of irradiance, temperature, and nutrient on growth dynamics of seagrasses: a review. *J. Exp. Mar. Biol. Ecol.* 350: 144–175

- 
- Massa, S.I., Arnaud-Haond, S., Pearson, G.A., Serrão, E.A., 2009. Temperature tolerance and survival of intertidal populations of the seagrass *Zostera noltei* (Hornemann) in Southern Europe (Ria Formosa, Portugal). *Hydrobiologia*. 619, 195–201.
- McLeod, E., Chmura, G., Bouillon, S., Salm, R., Bjork, M., 2011. A blueprint for carbon: Toward an improved understanding of the role of vegetated coastal habitats in the sequestering CO<sub>2</sub>. *Front. In Eco. And the Env.* 10:552-560
- Michael, W.D., McIntir, D., 1983. Effects of physical gradients on the production dynamics of sediment- associated algae. *Mar. Ecol. Prog. Ser.* Vol. 13: 103-114
- Mercado, J. M., Niell, F.X., Silva, J., Santos, R., 2003. Use of light and inorganic carbon acquisition by two morphotypes of *Zostera noltei* Hornem. *J. Exp. Mar. Biol. Ecol.* 297:71–84
- Migné, A., Davoult, D., Spilmont, N., Menu, D., Boucher, G., et al., 2002. A closed chamber CO<sub>2</sub> flux method for estimating primary production and respiration under emersed conditions. *Mar. Biol.* 4: 865-869
- Ouisse, V., Migné, A., Davoult, D., 2010. Seasonal variations of community production, respiration and biomass of different primary producers in an intertidal *Zostera noltei* bed (Western English Channel, France). *Hydrobiologia*. 649: 3-11
- Ouisse, V., Davoult, D., 2011. Community level carbon flux variability over a tidal cycle in *Zostera marina* and *Z. noltei* beds. *Mar. Ecol. Prog. Ser.* 437: 79-87.
- Ouisse, V., Migné, A., Davoult, D., 2014. Comparative study of methodologies to measure in-situ the intertidal benthic community metabolism during immersion. *Est. Coast. and Shelf Sci.* 136: 19-25
- Ow, Y.X., Vogel, N., Collier, C.J., Holtum, J.A.M., Flores, F. et al., 2016. Nitrate fertilisation does not enhance CO<sub>2</sub> responses in two tropical seagrass species. *Sci. Rep.* 6: 23093.
- Parsons, T.R., Takahashi, M., Hargrave, B., 1984. *Biological oceanographic processes*, 3<sup>rd</sup> edition. Pergamon Press, London.
- Perez-Loren, J.L., Niell, F.X., 1994. Photosynthesis in air: comparative responses to different temperatures of two morphotypes of *Zostera noltei* Hornem. from Palmones River estuary (southern Spain). *Verh. Internat. Verein. Limnol.* 25: 2265–2269.
- Plus, M., Deslous-Paoli, J.M., Auby, I., Dagault, F., 2001. Factors influencing primary production of seagrass beds (*Zostera noltei* Hornem.) in the Thau Lagoon (French Mediterranean coast). *J. Exp. Mar. Biol. Ecol.* 259: 63–84
- Plus, M., Auby, I., Verlaque, M., Levavasseur, G., 2005. Seasonal variations in photosynthetic irradiance response curves of macrophytes from a Mediterranean Coastal lagoon. *Aq. Bot.* 81: 157-173
- Portner, H. O. 2008. Ecosystem effects of ocean acidification in times of ocean warming: a physiologist's view. *Mar. Ecol. Prog. Ser.* 373: 203–217.
- Porzio, L., Buia, M. C., Hall-Spencer, J.M., 2011. Effects of ocean acidification on macroalgal communities. *J. Exp. Mar. Biol. Ecol.* 400: 278–287.
-

- Rasmusson, M.L. 2015. Seagrass Respiration. An assessment of oxygen consumption patterns in temperate marine macrophytes. Stockholm University, Sweden.
- Royal Society., 2005. Ocean acidification due to increasing atmospheric carbon dioxide. London, The Royal Society U.K.
- Runcie, J.W., Durako, M.J., 2004. Among-shoot variability and leaf specific absorption characteristics affect diel estimates of in situ electron transport in *Posidonia australis*. *Aq. Bot.* 80: 209-220.
- Riebesell, U., Schulz, K. G., Bellerby, R. G. J., Botros, M., Fritsche, P., et al., 2007. Enhanced biological carbon consumption in a high CO<sub>2</sub> ocean. *Nature* 450: 545–548.
- Schwarz, A.M., 2008. Contributions of Photosynthetic gains during tidal emersion to production of *Zostera capricorni* in a North Island, New Zealand Estuary. *New Zealand J. of Mar. and Freshwater Res.* 38: 809-818.
- Santos, R., Silva, J., Alexandre, A., Navarro, R., Barron, C., et al., 2004. Ecosystem metabolism and carbon fluxes of a tidally-dominated coastal lagoon. *Estuaries* 27: 977–985.
- Short, F.T., Neckles, H.A., 1999. The effects of global climate change on seagrasses. *Aquat. Bot.* 63, 169–196
- Short, F.T., Coles, R.G., Pergent-Martini, C., 2001. Global seagrass distribution. In: Short, F.T., Coles, R.G. (Eds.), *Global Seagrass Research Methods*. Elsevier, Amsterdam, pp. 5–31.
- Silva, J., and Santos, R., 2003. Daily variation patterns in seagrass photosynthesis along a vertical gradient. *Mar. Ecol. Prog. Ser.* 257: 37-44
- Silva, J., Santos, R., 2008. Underwater measurements of carbon dioxide evolution in marine plant communities: A new method. *Estu. and Coastal Shelf Sci.* 78. 827-830
- Silva, J., Santos, R., Calleja, M.L., Duarte, C., 2005. Submerged versus air exposed intertidal macrophyte productivity: from physiological to community level assessments. *J. of Exp. Mar. Bio. and Ecol.* 317: 87-95.
- Streever, W.J., Genders, A.J., Cole, M.A., 1998. A closed chamber CO<sub>2</sub> flux method for estimating marsh productivity. *Aquat. Bot.* 62, 33–44
- Sven, U., Klumpp, D.W., 1998. Microphytobenthos community production at a near shore coral reef: seasonal variation response to ammonium recycled by holothurians. *Mar. Eco. Prog. Ser.* 169: 1-11
- Thom, R. M., 1996. CO<sub>2</sub> enrichment effect on eelgrass (*Zostera marina* L.) and bull kelp (*Nereocystis luetkeana* (MERT.) P. & R.). *Water Air Soil Pollut.* 88:383–391.
- Touchette, B.W., Burkholder, J.M., 2007. Carbon and nitrogen metabolism in the seagrass, *Zostera marina* L.: environmental control of enzymes involved in carbon allocation and nitrogen assimilation. *J. Exp. Mar. Biol. Ecol.* 350: 216–233
- Uthicke, S., Klumpp, D.W., 1998. Microphytobenthos community production at a near shore coral reef: Seasonal variation and response to ammonium recycled by holothurians. *Mar. Ecol. Prog. Ser.* 169: 1-11

Van Es, F.B., 1982. Community metabolism of intertidal flats in the Ems-Dollard Estuary. *Mar. Biol.* 66: 95-108

Viaroli, P., Bartoli, M., Bondavalli, C., Christian, R.R., Giordani, G., et al., 1996. Macrophyte communities and their impact on benthic fluxes of oxygen, sulphide and nutrients in shallow eutrophic environments. *Hydrobiologia* 329:

Van der Heide, T., Bouma, T.J., van Nes, E.H., van de Koppel, J., Scheffer, M., et al., 2010. Spatial self-organized patterning in seagrasses along a depth gradient of an intertidal ecosystem. *Ecology* 91. 362–369

Vermaat, J.E., Verhagen, F.C.A., 1996. Seasonal variation in the intertidal seagrass *Zostera noltei* Hornem.: coupling demographic and physiological patterns. *Aquat. Bot.* 52: 259-281

Vermaat, J.E., 2009. Linking clonal growth patterns and ecophysiology allows the prediction of meadow-scale dynamics of seagrass beds. *Perspect. Plant Ecol. Evolut. Syst.* 11: 137–155

Zimmerman, R. C., Kohrs, D. G., Steller, D. L., Alberte, R. S., 1997. Impacts of CO<sub>2</sub> enrichment on productivity and light requirements of eelgrass. *Plant Physiol.* 115: 599–607.



## Population dynamics of *Cymodocea nodosa* in the vicinity of volcanic CO<sub>2</sub> seeps



Mishra, A.K., Cabaço, S., Apostolaki, E., Vizzini, S., Santos, R

(To be submitted)



### Population dynamics of *Cymodocea nodosa* in the vicinity of volcanic CO<sub>2</sub> seeps.

#### Abstract

Rising carbon dioxide (CO<sub>2</sub>) concentrations in the atmosphere will increase the average  $p\text{CO}_2$  level in the world oceans, which will have a knock-on effect on the marine ecosystem. Coastal seagrass communities are predicted to benefit from the increase in CO<sub>2</sub> levels, but long term effects of elevated CO<sub>2</sub> on seagrass communities are less understood. Population reconstruction techniques were used to investigate the population dynamics of *Cymodocea nodosa* meadows, exposed to long term elevated CO<sub>2</sub> at volcanic seeps off Greece and Italy. Effect of elevated CO<sub>2</sub> was noticed on the growth, morphometry, density, biomass and age structure at CO<sub>2</sub> seeps than reference sites. Above to below ground biomass ratio of *C. nodosa* were higher at CO<sub>2</sub> seeps. The plastochrome interval were similar at all CO<sub>2</sub> seeps. The shoot age and shoot longevity of plants were lower at seeps. The present recruitment (sampled year) of the seagrass were higher than long-term average recruitment of the communities near the seeps. Carbon to nitrogen ratios (%DW) of *C. nodosa* were higher in leaves at seeps. Annual leaf production was higher near the seeps. This study suggests increased production of *C. nodosa* under elevated CO<sub>2</sub> levels, but other co-factors such as nutrients, trace metal toxicity must also be taken into consideration while predicting effects of future CO<sub>2</sub> concentrations.

**Keywords:** Reconstruction techniques, plastochrome interval, elevated CO<sub>2</sub>, carbon metabolism



### 3.1. Introduction

The ocean's absorption of anthropogenic CO<sub>2</sub> emissions (over 35 Giga tonnes of CO<sub>2</sub> per year, (IPCC, 2014) has already lowered the mean ocean surface pH by 0.1 units since preindustrial values, with a predicted further decrease of 0.3- 0.4 units by the end of this century (IPCC, 2014). These on-going changes are expected to intensify in the future with potentially significant, but variable effects on marine organisms depending on their sensitivity (Hendrick et al., 2010; Kroeker et al., 2010). Calcifying organisms are more susceptible to ocean acidification than non-calcifying organisms (Suggett et al., 2012), even though their responses to ocean acidification are also dependent on the taxonomic group and their developmental stages (Hendrick et al., 2010; Kroeker et al., 2010). Realisation of the key role of seagrass in coastal ecosystems has fostered ever growing efforts to quantify their annual productivity and growth dynamics. (Duarte et al., 1994). Seagrasses, as many macro algal species, are notably tolerant to CO<sub>2</sub> increases and may even benefit from it (Koch et al., 2013). In the current CO<sub>2</sub> levels, seagrasses are dissolved inorganic carbon (DIC) limited as they are inefficient in utilising bicarbonate (Invers et al., 2001).

Hence, under increased CO<sub>2</sub> conditions, it is expected seagrasses will increase utilization of CO<sub>2</sub> (Beer et al. 1996) resulting in increased photosynthesis and consequently growth and productivity (Koch et al., 2013, Russel et al., 2013). Despite these predictions, the real-time studies conducted overtime both ex-situ with high CO<sub>2</sub> and in-situ at natural CO<sub>2</sub> seeps on seagrasses not completely support this expectation. For example, the photosynthetic activity of *Cymodocea nodosa* was stimulated by the low pH at natural CO<sub>2</sub> seeps of Vulcano, with significant increase in chlorophyll-a content of leaves, maximum electron transport rate and compensation irradiance (Apostolaki et al., 2014). Low pH promoted productivity, but was not translated into biomass production, probably due to nutrient limitation, grazing or poor environmental conditions (Apostolaki et al., 2014). Similar results were obtained by Alexandra et al. (2012) for *C. nodosa* grown for five months under high CO<sub>2</sub> conditions. On the other hand, the biomass and net primary production of *Halophila ovalis* and *Cymodocea serrulata* increased near the CO<sub>2</sub> seeps, whereas the abundance of species increased only for *C. serrulata*, suggesting species specific response to elevated CO<sub>2</sub> (Russell et al., 2013). Long term elevated CO<sub>2</sub> experiments on *Zostera marina* for over one year showed greater reproductive outputs, increased below ground biomass and shoot density (Palacios and Zimmerman, 2007), whereas short term experiments resulted in increased photosynthetic rate and shoot productivity

(Zimmerman et al., 1997). In contrast, experiments on *Cymodocea serrulata* have shown no enhancement in productivity at higher CO<sub>2</sub> as they are carbon saturated in current CO<sub>2</sub> concentrations (Schwarz et al., 2000). Recent CO<sub>2</sub> enrichment studies on three tropical seagrass species showed a significant increase in net productivity with increase in CO<sub>2</sub> levels, but a different growth rate between species was noticed due to varying strategies of carbon allocation among species (Ow et al., 2015).

Studies on natural CO<sub>2</sub> seeps suggest that seagrass species can be adapted to survive and live under elevated CO<sub>2</sub> conditions (Hall Spencer et al., 2008, Fabricius et al., 2011 and Takahashi et al., 2016). Much of the natural CO<sub>2</sub> seeps of Europe are concentrated in the Mediterranean Sea (Dando et al., 1999). In the Mediterranean Sea, most known seeps are concentrated in the shallow waters of the active volcanic arcs in Aegean Seas and are usually of the gas hydrothermal type due to the large volume of gas released (Dando et al., 1999). These natural CO<sub>2</sub> seeps provide future oceanic conditions (Hall Spencer et al., 2008; Hall-Spencer and Rodolfo-Metalpa, 2009) and are expected to affect seagrass communities due to changes in the physical and chemical features of seawater and sediments with possibly large effects on functioning features (Vizzini et al., 2010). In these conditions the growth and age structure of the seagrass *Cymodocea nodosa* in these seeps have not been investigated. The seagrass *Cymodocea nodosa* is an endemic species that supports highly complex and biodiverse climax communities in the Mediterranean Sea (Mazzella et al., 1986). Nevertheless, the effects of hydrothermal CO<sub>2</sub> gas release associated to explosive volcanism activity on seagrass productivity has been studied (Vizzini et al., 2010). How these changes will affect higher levels of biological organization, such as seagrass population dynamics (e.g. shoot recruitment rate), is less studied.

The objective of this work is to assess for the first time the long-term responses of the population dynamics and production of seagrass, *C. nodosa* exposed to elevated CO<sub>2</sub> levels. Populations in the vicinity and away from the influence of volcanic seeps were compared. CO<sub>2</sub> seep sites have been used to assess the long-term effects of elevated CO<sub>2</sub> on benthic marine ecosystems and respective underlying mechanisms (Hall-Spencer et al., 2008, Fabricius et al., 2011, Vizzini et al., 2013, Enochs et al., 2015). However, other cofounding factors related to the volcanic seeps, such as the emissions of heavy metals (Dando et al., 1999, Vizzini et al., 2013, Kadar et al., 2013) and sulphide (Dando et al., 1999, Boatta et al., 2013) may influence the plants and population responses to elevated CO<sub>2</sub>. These are very variable among seeps

(Dekov et al., 2004, Varnavas et al., 2005) whereas the major CO<sub>2</sub> composition of emissions is constant. To cope with possible confounding factors of the effects of CO<sub>2</sub> on the population dynamics of *C. nodosa*, we replicated the sampling effort in three seeps, two at the island of Milos in Greece and one at Vulcano island in Italy, to consider only the common responses as effects of elevated CO<sub>2</sub>.

### 3.2. Methods

#### *Study sites*

##### *Milos Islands, Greece*

Paleochori Bay (36.67 N, 24.51 E) and Milos thermal stations (36.70 N, 24.46 E) are part of Milos island (Fig.1A). Extensive submarine venting occurs offshore, from the intertidal to depths of more than 100 m over a 34 km<sup>2</sup> area of seabed (Dando et al.,1999). The released gases are 95% CO<sub>2</sub> with some H<sub>2</sub>S, CH<sub>4</sub> and H<sub>2</sub> (Dando et al., 1999).

##### *Vulcano, Italy*

We sampled Levante Bay (38.4 N, 15.0 E) off Vulcano island (Fig. 1B). Some parts of this bay are well-suited for studies of the effects of increased CO<sub>2</sub> levels (Boatta et al., 2013) despite areas with elevated H<sub>2</sub>S and metals (Vizzini et al., 2013). The main underwater gas seeps are located along southern and western shores of the bay at <1 m depth (Boatta et al., 2013). Total CO<sub>2</sub> output is about 3.6 tonne d<sup>-1</sup> (Inguaggiato et al., 2012), and the underwater gas emissions are 97-98% CO<sub>2</sub> with 2.2% H<sub>2</sub>S close to the seeps, decreasing to less than 0.005% H<sub>2</sub>S towards the north-eastern part of the bay, where most ocean acidification research has been located (Capaccioni et al., 2001; Milazzo et al., 2014). There is a step gradient in carbonate chemistry with pH 5.65 at the main gas seeps increasing to pH 8.1, which is typical for present day Mediterranean surface seawater, at >350 m from the seeps (Boatta et al., 2013). *Cymodocea nodosa* were absent at the main vents.

In each location, two sites were established where well developed populations of *C. nodosa* were present, a high CO<sub>2</sub> site near the seeps and a reference site away from the influence of the vent. Reference and CO<sub>2</sub> seep sites were established at the same depths, under similar hydrodynamics conditions. Overall, *C. nodosa* meadows sampled were at shallow (<5m) depth.

### *Water sampling:*

Water samples were collected at CO<sub>2</sub> seeps and reference sites (n=5) in 100 ml Winkler bottles fixed with 20 µl mercury chloride in the field, stored in the dark and transported to the laboratory for total alkalinity (TA) and dissolved inorganic carbon (DIC) analysis. The pH (using pH meter, Titrimo Methron) and temperature of the water samples were measured in the field immediately after collection. In the laboratory pH and temperature were measured again and 80 ml of water sample was used in Lab Titrimo analyser (Dickson et al. 2007) for the analysis of TA. Temperature, pH and TA data were used to calculate the other carbonate parameters ( $p\text{CO}_2$ ,  $\text{HCO}_3^-$ , aragonite saturation state, etc.) using CO<sub>2</sub>Sys program (Pierrot and Wallace.2006).

### *Seagrass sampling and analysis:*

The meadow density and biomass, the rhizome growth and production, the morphometric characteristics of the plants, the population age structure and derived population dynamics (long-term average recruitment, present recruitment and population growth rates) and the total C and N contents of plants were characterized in CO<sub>2</sub> seeps and reference sites. Reconstruction techniques, an indirect measure of plant growth history and population dynamics changes (Duarte et al., 1994; Fourqurean et al. 2003), were used to evaluate the *C. nodosa* responses to the different environmental conditions caused by the seeps. It was hypothesized that increased CO<sub>2</sub> promote higher plant growth, higher meadow production, and thus higher population dynamics.

Ten *C. nodosa* samples were collected from each site with a 20-cm diameter core (15 cm in Greece) into a depth of about 30 cm in May 2013 in Italy and May 2014 in Greece. The sediment was carefully rinsed off to prevent the modular sets disconnecting from each other and to keep the rhizome mat intact as required for the reconstruction of seagrass dynamics (Duarte et al., 1994). In each sample, the number of both shoots and apicals was counted to estimate the density of shoots and of physically independent individuals. The age of *C. nodosa* shoots was estimated by counting the number of leaf scars on the vertical rhizomes plus the number of leaves in each shoot multiplied by the leaf plastochrome interval (PI). To estimate the PI of each study site, i.e., the time needed to produce a new leaf, the sequence of average internodal length of *C. nodosa* shoots collected with the cores plus additional plants collected by hand was plotted. Then a 30% running average was applied to filter short-term seasonal variability and the difference in the number of vertical leaf scars between two consecutive

length modes was counted. The modes represent annual growth periods and thus the average number of leaf scars produced between modes was averaged to estimate the leaf PI of the population (Short et al. 2001). The PI estimates of *Cymodocea nodosa* were 29.3 for all three locations off Greek and Italy islands.

To estimate the vertical and horizontal rhizome elongation rates the length of both the vertical and horizontal rhizomes between consecutive shoots was measured and the number of both vertical and horizontal internodes between consecutive shoots was counted (see Duarte et al., 1994 for details of method). The number of leaves per shoot were measured from intact shoots in each sample ( $n = 10$ ). The horizontal and vertical rhizome production rates were estimated by multiplying the elongation rates (vertical or horizontal) by density (shoots or apicals), by the specific dry weight of rhizomes (vertical or horizontal) and by the dry weight contents of C (vertical or horizontal). Annual leaf production of each population was calculated as the product of the number of leaves annually produced per shoot, the shoot density, the mean specific dry weight of fully developed leaves and their C content.

The leaves, vertical rhizomes, horizontal rhizomes and roots were separated and dried for 48 h at 60° C for biomass and production estimates. Dried plant material (leaves, vertical rhizomes, horizontal rhizomes and roots) was grounded and analysed for CHN contents in a CHN analyser (EA 1110 Model, Elemental Microanalysis Ltd, Oakhampton, Devon, UK).

The long-term average recruitment ( $R$ ) was estimated from the shoot age structure using the general model:  $N_x = N_0 e^{-Rx}$ , where  $N_x$  is the number of shoots in age class  $x$ ,  $N_0$  is the number of shoots recruited into the population; assuming that mortality and recruitment have had no trend over the lifespan of the oldest shoots in the population, i.e. have remained constant over the lifespan of the oldest shoots, with year to year random variation around some mean value of mortality and recruitment (Fourqurean et al. 2003, Duarte et al. 2005). The recruitment for the current year of sampling ( $R_0$ ) was estimated using the method described by Duarte et al., (1994). The population growth rate ( $r$ ) was estimated as:  $r = R_0 - M$ , where  $M$  is the long-term mortality rate, which equals the long-term recruitment rate ( $R$ ) under the assumptions of near steady state (Fourqurean et al., 2003). Population was considered growing if  $r$  is positive ( $R_0 > R$ ), shrinking if  $r$  is negative ( $R_0 < R$ ), or with the same trajectory pattern if  $R_0$  is not significantly different from  $R$  (Fourqurean et al., 2003).

### *Statistical analysis:*

Significant differences in biomass, density, production and plant morphometry among sites (CO<sub>2</sub> seeps and reference) and locations (Milos, Paleochori and Vulcano) were investigated using two-way ANOVA, after testing for homogeneity of variances and normality of distribution. The Tukey's multiple comparison test was applied to determine significant differences between factor levels. When ANOVA assumptions were not verified, comparison of data sets were performed using the non-parametric test of Kruskal-Wallis and the post hoc Dunn's test. Log transformations of variables were performed where needed.

The species vertical and horizontal rhizome elongation, and the population recruitment rates were obtained considering all replicates in each site. The t-test for the difference between two regression lines was used to compare the vertical rhizome elongation rates as these are equal to the slopes of the linear regression between age and size of rhizomes. Statistical analyses were not performed for the horizontal rhizome elongation rate, because just one value was obtained for each site. The confidence limits of the exponential decay regression model used to estimate the long-term average recruitment rate ( $R$ ) allowed its statistical comparison to the present recruitment rate ( $R_0$ ) as described in Fourqurean et al., (2003). Significant differences of the long-term recruitment rate among sites were tested using one-way ANOVA. Significance levels was considered at  $p < 0.05$  (Sokal and Rohlf, 2012).

### 3.3. Results

Seawater carbonate chemistry values of the three locations Milos, Paleochori and Vulcano are presented in Table 1. At all locations, the  $p\text{CO}_2$  concentration was higher (and pH lower) near the  $\text{CO}_2$  seeps (Table 1). The reference sites lying further away from the  $\text{CO}_2$  seeps (Reference) showed lower concentrations of  $\text{CO}_2$ . DIC and  $\text{CO}_2$  concentrations were higher at Milos and lower at Vulcano.

Significant response of *C. nodosa* to increased  $\text{CO}_2$  was noticed on density and biomass at all three  $\text{CO}_2$  seeps. Both shoot and apex density, i.e. the number of physically independent plants was higher in all  $\text{CO}_2$  seep sites. Both shoot and apex density was higher at Milos (Fig.2). Total biomass, leaf biomass, horizontal rhizome biomass and above ground-below ground biomass ratio of *C. nodosa* varied significantly at all three  $\text{CO}_2$  seeps. Total biomass, leaf biomass and horizontal biomass was higher at seeps of Vulcano, whereas above ground-below ground biomass ratio was higher at seeps of Paleochori (Fig. 5).

Table 1:

Seawater carbonate chemistry measurements at Milos, Paleochori and Vulcano  $\text{CO}_2$  seeps calculated with  $\text{CO}_2\text{Sys}$  programme, using constants from Dickson and Millero, 1987, and pH in NBS scale.

	<b>Milos</b>		
	<b>pH (NBS scale)</b>	<b><math>p\text{CO}_2</math> (<math>\mu\text{atm}</math>)</b>	<b>DIC (<math>\mu\text{ mol kg}^{-1}</math>)</b>
CO <sub>2</sub> seeps	7.5 ± 0.04	2457.9 ± 1.87	3474.03 ± 4.55
Reference	8.2 ± 0.03	405.5 ± 1.65	1407.44 ± 51.25
	<b>Paleochori</b>		
CO <sub>2</sub> seeps	7.9 ± 0.01	884.3 ± 3.07	2738.39 ± 16.49
Reference	8.2 ± 0.01	402.9 ± 1.1	1396.14 ± 48.69
	<b>Vulcano</b>		
CO <sub>2</sub> seeps	7.98 ± 0.08	737 ± 158	2377 ± 37
Reference	8.17 ± 0.05	427 ± 68	2244 ± 40

Table 2.

Responses of *Cymodocea nodosa* biomass (means  $\pm$  SE) at CO<sub>2</sub> seeps off Greek (Milos and Paleochori) and Italy (Vulcano) islands. Tukey's multiple comparison ( $p < 0.05$ ) test results are presented. Different letters indicate significant difference between CO<sub>2</sub> seeps and reference sites.

	<b>CO<sub>2</sub> seeps</b>	<b>Reference</b>
	<b>Milos</b>	
Vertical rhizome biomass (g DW m <sup>-2</sup> )	34.4 $\pm$ 5.3	28.6 $\pm$ 4.2
Root biomass (g DW m <sup>-2</sup> )	41.1 $\pm$ 1.1	31.5 $\pm$ 3.0
	<b>Paleochori</b>	
Vertical rhizome biomass (g DW m <sup>-2</sup> )	28.60 $\pm$ 5.9	14.4 $\pm$ 2.7
Root biomass (g DW m <sup>-2</sup> )	36.0 $\pm$ 2.6	30.3 $\pm$ 1.8
	<b>Vulcano</b>	
Vertical rhizome biomass (g DW m <sup>-2</sup> )	67.6 $\pm$ 10.0	124.4 $\pm$ 17.6
Root biomass (g DW m <sup>-2</sup> )	135.9 $\pm$ 27.4	154.6 $\pm$ 18.7

The vertical rhizome biomass was significantly lower near the seeps of Vulcano but not in the other seeps (Table 2). Significant effects of CO<sub>2</sub> on plant morphology was observed on number of leaves and vertical rhizome. The number of leaves was higher at Milos and the vertical rhizome which was shorter near the seeps at all sites (Fig.3). Density, biomass and morphometry of *C. nodosa* varied significantly between Greek and Italian locations (Milos vs Vulcano, Paleochori vs Vulcano) but not between Greek locations (Milos vs Paleochori), except for number of leaves which was significant between Greek locations and shoot density that did not vary among locations (see Appendix). Significant differences between apex density was found between Milos vs Vulcano and Vulcano vs Paleochori (See Appendix). Apex density was lower at Vulcano (Fig.2) whereas the total and plant part biomass were higher (Table 2). Leaf number (Fig.3) and horizontal rhizome length were lower at Vulcano (see Appendix), whereas vertical rhizome length was higher (Fig.3). Significant differences were found for vertical rhizome length and number of leaves between Milos vs Vulcano and Vulcano vs Paleochori, whereas the only significant difference in morphometry were found for number of leaves at Milos vs Paleochori (see Appendix). Vertical rhizome length in Vulcano was higher than Milos and Paleochori (Fig.3). Significant differences were found for vertical elongation rate and the vertical elongation rate was higher at CO<sub>2</sub> seeps at all locations (see Appendix).

Horizontal elongation rate was higher at CO<sub>2</sub> seep sites at all locations except for Vulcano CO<sub>2</sub> seep site (See Appendix).

Table 3

Age structure and population dynamics values of *Cymodocea nodosa* shoots at CO<sub>2</sub> seeps and reference sites off Greek (Milos and Paleochori) and Italy (Vulcano) islands. Mean  $\pm$  Standard errors are presented for the shoot age. The exponential coefficient  $\pm$  standard errors of the exponential decay regression are presented for the long-term average recruitment rate (R). Different letters indicate significant difference between CO<sub>2</sub> seeps and reference sites, ns= not significant. P values obtained from one way ANOVA for Milos (p=0.116), Paleochori (p=0.013) and Vulcano (p= 0.105)

	<b>Milos</b>	
	<b>CO<sub>2</sub> seeps</b>	<b>Reference</b>
Shoot longevity (years)	3.21	3.53
Long term avg. recruitment (R, year <sup>-1</sup> )	1.03 $\pm$ 0.05 <sup>ns</sup>	0.90 $\pm$ 0.05 <sup>ns</sup>
Present recruitment rate (R <sub>0</sub> , year <sup>-1</sup> )	1.08	0.94
Population growth rate (r, year <sup>-1</sup> )	0.05	0.04
	<b>Paleochori</b>	
	<b>CO<sub>2</sub> seeps</b>	<b>Reference</b>
Shoot longevity (years)	2.41	2.81
Long term avg. recruitment (R, year <sup>-1</sup> )	1.41 $\pm$ 0.08 <sup>a</sup>	0.42 $\pm$ 0.23 <sup>b</sup>
Present recruitment rate (R <sub>0</sub> , year <sup>-1</sup> )	1.54	0.52
Population growth rate (r, year <sup>-1</sup> )	0.07	0.09
	<b>Vulcano</b>	
	<b>CO<sub>2</sub> seeps</b>	<b>Reference</b>
Shoot longevity (years)	7.14	8.27
Long-term average recruitment rate (R, year <sup>-1</sup> )	0.64 $\pm$ 0.11 <sup>ns</sup>	0.37 $\pm$ 0.05 <sup>ns</sup>
Present recruitment rate (R <sub>0</sub> , year <sup>-1</sup> )	0.65	0.36
Population growth rate (r, year <sup>-1</sup> )	0.01	-0.01

Significant effects of elevated CO<sub>2</sub> were also observed on *C. nodosa* production. The vertical and horizontal rhizome productions were higher at all three CO<sub>2</sub> seep sites (Fig.6) The vertical and horizontal rhizome production was higher at Milos (Fig.6). Significant differences were found for vertical rhizome production between Vulcano vs Milos and Milos vs Paleochori sites, whereas for horizontal rhizome production significant differences were found at all three

CO<sub>2</sub> seep sites (see Appendix). Significant effects of elevated CO<sub>2</sub> were observed on the annual leaf production with higher leaf production at CO<sub>2</sub> seeps off all locations (Fig.7). Within the three locations significant differences were found for Milos vs Vulcano and Vulcano vs Paleochori (see Appendix).

The elemental content of carbon and C: N ratios of *C. nodosa* plant compartments (leaves, rhizomes and roots) showed that whenever there are significant differences, these reflect a carbon enrichment of tissues near CO<sub>2</sub> seeps (Fig.4). Significantly higher carbon content was observed near the seeps in the leaves and roots at Milos, leaves at Paleochori and leaves and rhizomes at Vulcano. The only significant difference found in the tissues N content was in the rhizomes of Paleochori, where it was lower near the seeps (Fig.4). Significant differences in C: N ratios were observed in roots off Milos, rhizomes off Paleochori and roots and rhizomes off Vulcano (Fig.4). Significant differences of C content of leaves among locations was found at Milos vs Vulcano and Vulcano vs Paleochori, whereas for C content of roots significant differences were found at Milos vs Paleochori and Milos vs Vulcano (see Appendix). Significant differences of N content among locations was found in leaves at Vulcano vs Paleochori and Milos vs Vulcano, whereas for N content of roots significant differences among locations was found at Milos vs Paleochori and Vulcano vs Paleochori (see Appendix). For C: N content significant differences was found in the leaves and rhizomes at Milos vs Paleochori and Vulcano vs Paleochori (see Appendix).

The age structure of *C. nodosa* was significantly affected at CO<sub>2</sub> seeps, as both the average shoot age (Fig.8) and shoot longevity, were lower near the seeps at all locations (Table 3). The lifespan of the oldest shoots was between seven to eight years at Vulcano, whereas at Paleochori and Milos it varied between three to four years (Fig. 9). The present and long term recruitments were always higher near the seeps, even though the differences were not statistically significant, except for Paleochori (Table 3). These results suggest higher dynamics of populations near the CO<sub>2</sub> seeps as the turnover of plants are higher near the seeps plants die earlier and recruitment is higher. Supporting this, the age frequency distribution of *C. nodosa* showed a significant higher number of younger plants (1 and 2 years older) at the CO<sub>2</sub> seeps, whereas the number of older plants (more than 3 years) were higher near the reference sites at all three locations (Fig.9).




Fig. 1. Study sites with location of CO<sub>2</sub> seeps and Reference sites, A) Milos and Paleochori off Greece and B) Vulcano island off Italy




Fig.2. Density (Apex and Shoot) of *Cymodocea nodosa* at the CO<sub>2</sub> seeps off the Greek (Milos and Paleochori) and Italy (Vulcano) islands. Error bars represent standard errors. Significant differences between CO<sub>2</sub> seeps and reference sites are indicated by different letters.




Fig.3. Morphometry (Number of Leaves and Vertical rhizome length) of *Cymodocea nodosa* at CO<sub>2</sub> seeps and reference sites off Greek (Milos and Paleochori) and Italy (Vulcano) islands. Error bars represent standard errors. Significant differences between CO<sub>2</sub> seeps and reference sites are indicated by different letters.




Fig. 4. Carbon, nitrogen elemental contents and its ratio for Milos (a, b, c), Paleochori (d, e, f) and Vulcano (g, h, i) in leaves, rhizomes and roots of *Cymodocea nodosa* plants at CO<sub>2</sub> seeps and reference sites off Greece and Italy respectively, (mean ± SE, n = 5). Error bars represent standard errors. Significant differences between CO<sub>2</sub>seeps and reference sites are indicated by different letters.







Fig. 5. Biomass of *Cymodocea nodosa* at the CO<sub>2</sub> seeps and reference sites off Greek (Milos and Paleochori) and Italy (Vulcano) islands. Error bars represent standard errors. Significant differences between CO<sub>2</sub> seeps and reference sites are indicated by different letters. (Ag: Bg; Above ground biomass: Below ground biomass)



a



b



c

Fig.6. Vertical and horizontal rhizome productions of *Cymodocea nodosa* at the CO<sub>2</sub> seeps and reference sites off Greek (Milos and Paleochori) and Italy (Vulcano) islands. Error bars represent standard errors. Significant differences between CO<sub>2</sub> seeps and reference sites are indicated by different letters




Fig.7. Annual leaf Production of *Cymodocea nodosa* at the CO<sub>2</sub> seeps and reference sites off Greek (Milos and Paleochori) and Italy (Vulcano) islands. Error bars represent standard errors. Significant differences between CO<sub>2</sub> seeps and reference sites are indicated by different letters.




Fig.8. Shoot age of *Cymodocea nodosa* at the CO<sub>2</sub> seeps and reference sites off Greek (Milos and Paleochori) and Italy (Vulcano) islands. Error bars represent standard errors. Significant differences between CO<sub>2</sub> seeps and reference sites are indicated by different letters.




Fig. 9. Age frequency distribution of *Cymodocea nodosa* population along the CO<sub>2</sub> seeps and reference sites off Milos (a, b), Paleochori (c, d) and Vulcano (e, f) respectively. The long-term average recruitment rate (R) was estimated from the exponential decay regression line fitted to age frequency distribution.

### 3.4. Discussion:

The shallow CO<sub>2</sub> volcanic seeps of the Greek (Milos and Paleochori) and Italy (Vulcano) islands have had elevated  $p\text{CO}_2$  level for many years, providing insights about the effects of elevated CO<sub>2</sub> on seagrass communities. Despite the variation in  $p\text{CO}_2$  at the seeps, the sites represented current (reference) and future 2100 (elevated CO<sub>2</sub>) scenarios, enabling us to assess the long-term effects of elevated CO<sub>2</sub> on seagrass communities. Increase in CO<sub>2</sub> concentrations from the seeps, not only increase the dissolved CO<sub>2</sub> concentrations in the surrounding waters, but also increased the relative portion of dissolved CO<sub>2</sub> to HCO<sub>3</sub><sup>-</sup> (Short and Neckles, 1999) with a positive effect on seagrasses, as observed for the vertical and horizontal rhizome productivity, apex and shoot density and biomass of *C. nodosa* found at Milos, Paleochori and Vulcano CO<sub>2</sub> seeps.

*C. nodosa* communities showed an increased plant growth and production, and higher population dynamics characterized by higher shoot recruitment and younger shoots under increased CO<sub>2</sub> at Milos, Paleochori and Vulcano sites. Plants developing near the CO<sub>2</sub>-seeps off Milos, Paleochori and Vulcano express an increase in fitness confirming that seagrasses are able to thrive in predicted scenarios of global CO<sub>2</sub>/OA changes (Hall-Spencer et al. 2008; Russell et al. 2013). Increased growth and production with proximity to CO<sub>2</sub>-seeps suggests increase in CO<sub>2</sub> concentrations will favour the utilization of inorganic carbon sources for seagrass as plants were C limited (Invers et al., 2001; Apostolaki et al., 2014)

The significant spatial variability in *C. nodosa* may be related to high level of patchiness of the seagrass meadow which was evident during sampling. Apex density of *C. nodosa* were higher at all three CO<sub>2</sub> seeps compared to reference sites, but the higher density was not transferred to significant patch expansion of seagrass meadows close to the seeps, even though *C. nodosa* can translocate itself to make meadows (Kraemer and Mazzella, 1999). Lack of *C. nodosa* meadows near the CO<sub>2</sub> seeps can also be related to the associated geochemical features (e.g. toxic levels of trace elements) of individual CO<sub>2</sub> seeps affecting the migration of meadow spatially. Increase in density at all three CO<sub>2</sub> seeps and biomass (only at Greek CO<sub>2</sub> seep sites) for *C. nodosa* from our studies coincides with a range of seagrass species, for instance density and biomass increased with increase in CO<sub>2</sub> levels, for *P. oceanica* (Hall-Spencer et al., 2008), *C. rotundata* and *C. serrulata* (Fabricius et al., 2011b; Russell et al., 2013) and *Zostera marina* (Palacios and Zimmerman, 2007), whereas decrease in biomass for *C. nodosa* in our studies in

Vulcano CO<sub>2</sub> seeps coincides with the results of Russell et al., (2013) for *Halophila ovalis* implying species-specific response towards increased CO<sub>2</sub> levels in future oceans. Higher density at all three CO<sub>2</sub> seeps and higher horizontal rhizome biomass of *C. nodosa* at Vulcano among three CO<sub>2</sub> seeps, suggest, *C. nodosa* at these seeps are adjusted to their carbon utilization capacity to the dissolved inorganic carbon (both CO<sub>2</sub> (aq) and HCO<sub>3</sub><sup>-</sup>) concentrations of the surrounding water, similar results were obtained for *T. testudinum* in field and laboratory conditions (Durako.,1993). The low pH conditions were followed by increase in density (shoot and apex) of *C. nodosa* at all three CO<sub>2</sub> seeps, compared to the reference sites, coinciding with results observed for *Cymodocea rotundata* and *Cymodocea serrulata* at Papua New Guinea CO<sub>2</sub> seeps (Takahashi et al., 2016) and differing from results of *C. nodosa* at CO<sub>2</sub> seeps of Vulcano (Eugenia et al., 2014), whereas decrease of total biomass in Vulcano agrees to the results of *C. nodosa* at CO<sub>2</sub> seeps of Vulcano (Eugenia et al., 2014). The above ground biomass was above 20% for Greek and Italy CO<sub>2</sub> seep sites, which agrees with findings for *C. nodosa* at Ischia seeps (Gianluigi et al., 2002), suggesting lower below ground biomass (lower than 80%) of seagrass meadows in elevated CO<sub>2</sub> levels. Our results of lower below ground biomass didn't match with results from other seagrass (*Zostera marina*) in CO<sub>2</sub> enriched conditions (Welsh et al. 1997; Palacios and Zimmerman, 2007).

Carbon content in *C.nodosa* leaves increased at the CO<sub>2</sub> seep sites for Paleochori and Milos sites, and remained invariable for Vulcano sites, which agrees with experimental results from *T. Testudinum* and *T. Hemprichii* that showed that carbon content remained invariable in CO<sub>2</sub> enriched conditions (Campbell and Fourqurean et al., 2013; Jiang et al., 2010). The N content of the *C. nodosa* leaves (1.67 % DW ± 0.36) were low at Paleochori CO<sub>2</sub> seep and were below the threshold (1.8% DW) for N limitation in seagrass (Duarte. 1990) suggesting that nitrogen deficiency may have limited overall plant growth, whereas N content of *C.nodosa* leaves at Milos (1.85% DW ± 0.05) were not along the threshold level and may have added to higher growth and density of plants at the Milos CO<sub>2</sub> seeps. In case of *C. nodosa* leaves at Vulcano even with higher N content, growth and density remained lower, similar results were obtained for *C. nodosa* in experimental CO<sub>2</sub> and nutrient enrichment conditions (Khan et al., 2016). Simultaneously lower growth can be attributed to the epiphytic microalgae on seagrass leaves, which gets dominant during high N content, compete with the seagrass in production and eventually shade the seagrass leaves (Valiela et al., 1997). Lower C: N ratios in leaves of the plants at Milos, Paleochori and Vulcano (Fig.4) suggests that plants near the seeps utilize

the available nitrogen rapidly relative to their carbon content (Duarte, 1990). As nutrient availability increases seagrass tissues, must have been become progressively enriched in nitrogen relative to the carbon content implying decreasing C: N ratios, as found in rhizomes of all three locations and roots of Vulcano CO<sub>2</sub> seeps, whereas the opposite trends were observed in the roots of *C. nodosa* at Greek CO<sub>2</sub> seeps (Fig.4).

Additionally, the Milos, Paleochori and Vulcano seeps, apart from CO<sub>2</sub> supplies, may also be enriched volcanic inputs such as trace elements, which above certain levels can be toxic (Vizzini et al. 2013) along with low pH and depleted oxygen levels can deteriorate the chemical properties of nearby water and sediment and in turn affect the seagrass productivity negatively. It is also likely that responses of seagrasses to elevated levels of CO<sub>2</sub> depends on other factors, such as interactions with temperature, nutrients and light of that ecosystem (Palacios and Zimmerman, 2007).

The low rhizome elongation (ca. 4.27 – 10.6 cm year<sup>-1</sup>) for *C. nodosa* at Milos, Paleochori and Vulcano (ca.7.91 cm year<sup>-1</sup>) CO<sub>2</sub> seeps suggested a reduced recruitment (Duarte and Sand-Jensen, 1990) compared to other populations elsewhere (40 cm year<sup>-1</sup>; Marbà and Duarte, 1998). Increased growth (vertical and horizontal elongation) and production (vertical rhizome, horizontal rhizome and leaf) with proximity to CO<sub>2</sub> seeps than reference sites at all three locations suggests plants were C limited in current conditions in the world oceans (Zimmerman et al., 1997; Invers et al., 2001). Our results comply with what has been observed for other seagrass species at natural CO<sub>2</sub>-seeps elsewhere, for instance *Cymodocea serrulata* and *Halophila ovalis* in Papua New Guinea (Russell et al. 2013) and for *Posidonia oceanica* in Ischia, Italy (Hall-Spencer et al., 2008). Similar results were also obtained under experimental CO<sub>2</sub> enrichment for *Zostera marina* (Palacios and Zimmerman, 2007) and *Zostera noltii* (Alexandre et al. 2012).

Our results represent an early assessment of seagrass *C. nodosa* response under long term CO<sub>2</sub> enrichment using population reconstruction techniques as a tool. This is first of its kind to represent seagrass growth, production at the CO<sub>2</sub> seeps of Milos, Paleochori and Vulcano off Greece and Italy. Comparison to similar studies of *C. nodosa* at other seeps suggests that seagrass response to naturally acidified conditions is not so straight forward as the response is species-specific and depends on the biogeochemical characteristics of the site. Clearly, further

research of seagrass growth and production using reconstruction techniques in CO<sub>2</sub> seeps in world oceans is necessary, keeping in mind the ecosystem services seagrass provides, their capacity to act as global carbon sink and their major role in mitigation of the elevated levels of CO<sub>2</sub> in future oceans.

### 3.5. References

- Alexandre, A, Silva, J., Buapet, P., Bjork, R., and Santos, R., 2012. Effects of CO<sub>2</sub> enrichment on photosynthesis, growth and nitrogen metabolism of the seagrass *Zostera noltii*. *Eco. and Evol.*10: 2620-2630
- Brierley, S. A., Kingsford, J.M., 2009. Impacts of climate change on marine organisms and ecosystems. *Curr. Bio.* 14: 602-604.
- Beer, S., and Koch, E., 1996. Photosynthesis of marine macro algae and seagrasses in globally changing CO<sub>2</sub> environments. *Mar. Eco. Prog. Ser.* 141:199–204
- Boatta, F., D'Alessandro, W., Gagliano, A.L., Liotta, M., et al., 2013. Geochemical survey of Levante bay, Vulcano island (Italy), a natural laboratory for the study of ocean acidification. *Mar. Pollut. Bull.* 73: 485-494
- Cabaço, S., Santos, R., Duarte, C.M., 2008. The impact of sediment burial and erosion seagrasses: A review. *Estu. Coast. and Shelf Sci.*79: 354-366.
- Campbell, J.E., Fourqurean, J.W., 2013. Effects of in situ CO<sub>2</sub> enrichment on the structural and chemical characteristics of the seagrass *Thalassia testudinum*. *Mar. Bio.* 160: 1465-1475
- Caldeira, K., and Wickett, M.E., 2003. Oceanography: anthropogenic carbon and ocean pH. *Nature.* 425:365.
- Capaccioni, B., Tassi, F., Vaselli, O., 2001. Organic and inorganic geochemistry of low temperature gas discharges at the Baia di Levante beach, Vulcano Island, Italy. *J. Volcanol. Geoth. Res.* 108: 173–185.
- Cunha, A.H., Duarte, C.M., 2005. Population age structure and rhizome growth of *Cymodocea nodosa* in the Ria Formosa (southern Portugal). *Mar. Bio.* 146: 841-847.
- den Hartog, C., 1970. *The Seagrasses of the World*. North Holland Publication., Amsterdam, 275pp.
- Dando, P. R., Stuben, D., & Varnavas, S. P., 1999. Hydrothermalism in the Mediterranean Sea. *Prog. in Oceano.* 44: 333–367.
- Dando, P., Aliani, S., Arab, H., Bianchi, C., et al., 2000. Hydrothermal studies in the Aegean Sea. *Phys. Chem. Earth B. Hydrol. Oceans Atmos.* 25: 1-8
- Dickson, A.G., Wolf-Gladrow, D. A., Zeebe, R.E., Klaas, C., et al., 2007. Total alkalinity: The explicit conservative expression and its application to biogeochemical processes. *Mar. Chem.* 106: 287-300.

- Duarte, C.M., Marbà, N., Agawin, N., Cebrián, J., et al., 1994. Reconstruction of seagrass dynamics: age determinations and associated tools for the seagrass ecologist. *Mar. Eco. Prog. Ser.* 107: 195-209.
- Duarte, C.M., Sand-Jensen, K., 1990. Seagrass colonization: biomass development and shoot demography in *Cymodocea nodosa* patches. *Mar. Eco. Prog. Ser.* 67: 97-103.
- Duarte, M. C., 1990. Seagrass nutrient content. *Mar. Eco. Prog. Ser.* 67: 201-207.
- Dekov, V.M., Savelli, C., 2004. Hydrothermal activity in the SE Tyrrhenian Sea: an overview of 30 years of research. *Mar. Geol.* 204: 161-185.
- Enochs, I.C., Manzello, D.P., Donham, E.M., Kolodziej, G., et al., 2015. Shift from coral to macroalgae dominance on a volcanically acidified reef. *Nat. Clim. Change.* 5: 1083–1088.
- Eugenia, T.A., Vizzini, S., Hendriks, I.E., Olsen, Y.S., 2014. Seagrass ecosystem response to long term high CO<sub>2</sub> in a Mediterranean volcanic vent. *Mar. Env. Res.* 99: 9-15.
- Fabricius, K.E., Langdon, C., Uthicke, S., Humphrey, C., et al., 2011b. Losers and winners in coral reefs acclimatized to elevated carbon dioxide concentrations. *Nat. Clim. Change.* 1: 165-169
- Feely, R. A., Sabine, C. L., Lee, K., Berelson, W., et al., 2004. Impact of anthropogenic CO<sub>2</sub> on the CaCO<sub>3</sub> system in the oceans. *Science* 305:362–366
- Fourqurean, W.J., Duarte, C.M., Marba. N., 2003. Elucidating seagrass population dynamics: Theory, constraints and practice. *Limnol. Oceano.* 48: 2070-2074.
- Gianluigi, C., Buia, M.C., Mazzella, L., 2002. Structure and growth dynamics of *Cymodocea nodosa meadows*. *Sci. Mar.* 66: 365-373.
- Hall-Spencer, J.M., Rodolfo-Metalpa, R., Martin, S., Ransome, E., et al., 2008. Volcanic carbon dioxide seeps show ecosystem effects of ocean acidification. *Nature.* 454: 96-99.
- Hall-Spencer, J.M., Rodolfo-Metalpa, R.A.B., 2009. Using scientific diving to investigate the long-term effects of ocean acidification at CO<sub>2</sub> seeps. *Online Geosci.* 22: 72-76
- Hemminga, M.A., Duarte, C.M., 2000. *Seagrass Ecology*. Cambridge University Press, Cambridge. 289 pp.
- Hendriks, I. E., Duarte, C.M., Alvarez, M., 2010. Vulnerability of marine biodiversity to ocean acidification: a meta-analysis. *Estu. Coastal and Shelf Sci.* 86: 157–164.
- International Panel for Climate Change (IPCC). 2014. *Fifth Assessment Report on Climate change: Impacts, Adaptation and Vulnerability*.
- Invers, O., Zimmerman, R.C., Alberte, R.S., Perez, M., et al., 2001. Inorganic carbon sources for seagrass photosynthesis: an experimental evaluation of bicarbonate use in species inhabiting temperate waters. *J. of Exp. Mar. Bio. Eco.* 265: 203-217
- Inguaggiato, S., Mazot, A., Diliberto, I.S., Inguaggiato, C., et al., 2012. Total CO<sub>2</sub> output from Vulcano Island (Aeolian Islands, Italy). *Geochem. Geophys. Geosyst.* 13.Q02012
- Jiang, Z.J., Huang, X.P., Zhang, J.P., 2010. Effects of CO<sub>2</sub> enrichment on photosynthesis, growth, and biochemical composition of seagrass *Thalassia hemprichii* (Ehrenb.) Aschers. *J. of Int. Plant Bio.* 52: 904-913.

- 
- Kádár, E., Costa, V., Segonzac, M., 2007. Trophic influences of metal accumulation in natural pollution laboratories at deep-sea hydrothermal seeps of the Mid- Atlantic Ridge. *Sci. Tot. Environ.* 373: 464-472.
- Khan, F., Alexandre, A., Ullah, H., Santos, R., 2016. Effects of elevated CO<sub>2</sub> and nutrients on the community metabolism of a *Cymodocea nodosa* bed. *Turk. J. of Bot.* 40: 250-257
- Koch, M., Bowes, G., Ross, C., Zhang, X., 2013. Climate change and ocean acidification effects on seagrasses and marine macro algae. *Glo. Change Bio.* 19: 103-132
- Kroeker, K. J., Kordas, R. L., Crim, R.N., Singh, G. G., 2010. Meta-analysis reveals negative yet variable effects of ocean acidification on marine organisms. *Eco. Let.* 13: 1419–1434
- Kraemer, G.P., Mazzella, L., 1999. Nitrogen acquisition, storage, and use by the co-occurring Mediterranean seagrasses *Cymodocea nodosa* and *Zostera noltii*. *Mar. Eco. Prog. Ser.* 183: 95-103.
- Marbà, N., Cebrián, J., Enríquez, S., Duarte, C.M., 1994. Migration of large-scale subaqueous bed forms measured with seagrasses (*Cymodocea nodosa*) as tracers. *Lim. and Oceano.* 39: 126-133.
- Marbà, N., Duarte, C.M., 1994. Growth response of the seagrass *Cymodocea nodosa* to experimental burial and erosion. *Mar. Eco. Prog. Ser.* 107: 307-311.
- Marbà, N., Duarte, C.M., 1995. Coupling of seagrass (*Cymodocea nodosa*) patch dynamics to subaqueous dune migration. *J. of Eco.* 83: 1-9.
- Marbà, N., Duarte, C.M., 1998. Rhizome elongation and seagrass clonal growth. *Mar. Eco. Prog. Ser.* 174: 269-280.
- Martin, S., Rodolfo -Metalpa, R., Ransome E., Rowley S., et al., 2008. Effects of naturally acidified seawater on seagrass calcareous epibionts. *Bio. Letters.* 4: 689-692
- Mazzella, L. and Alberte, R.S., 1986. Light adaptation and the role of autotrophic epiphytes in primary production of the temperate seagrass, *Zostera marina* (L). *J. of Exp. Mar. Bio. Eco.* 100: 165-180.
- Meehl, G.A., Stocker, T.F., Collins, W.D., Friedlingstein, P., et al., 2007. The Physical Science Basis Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge University Press. 747-845.
- Milazzo, M., Rodolfo-Metalpa, R., Chan, V.B.S., Fine, M. et al., 2014. Ocean acidification impairs vermetid reef recruitment. *Scienti. Rep.* 4 (4189).
- Olesen, B., Enríquez, S., Duarte, C.M., Sand-Jensen, K., 2002. Depth-acclimation of photosynthesis, morphology and demography of *Posidonia oceanica* and *Cymodocea nodosa* in the Spanish Mediterranean Sea. *Mar. Eco. Prog. Ser.* 236: 89-97.
- Ow. Y.X., Collier. C.J., Uthicke. S., 2015. Response of three tropical seagrass species to CO<sub>2</sub> enrichment. *Mar. Bio.* 162: 1005-1017.
- Palacios, S.L., Zimmerman, R.C., 2007. Response of eelgrass *Zostera marina* to CO<sub>2</sub> enrichment: possible impacts of climate change and potential for remediation of coastal habitats. *Mar. Eco. Prog. Ser.* 344: 1-13
-

- Perez, M., Duarte, C.M., Romero, J., Sand-Jensen, K., et al., 1994. Growth plasticity in *Cymodocea nodosa* stands: the importance of nutrient supply. *Aq. Bot.* 47: 249-264.
- Pierrot, D. E. L., and Wallace, D. W. R., 2006. MS Excel Program Developed for CO<sub>2</sub> System Calculations. ORNL/CDIAC-105a. Carbon Dioxide Information Analysis Centre, Oak Ridge National Laboratory, US Department of Energy, Oak Ridge, TN.
- Russell, B.D., Connell, S.D., Uthicke, S., Muehlehner, N., et al., 2013. Future seagrass beds: can increased productivity lead to increased carbon storage? *Mar. Pollut. Bull.* 73: 463-469
- Schwarz, A.M., Bjork, M., Buluda, T., Mtolera, H., et al., 2000. Photosynthetic utilisation of carbon and light by two tropical seagrass species as measured in situ. *Mar. Biol.* 137: 755-761
- Short, F.T., Coles, R.C., 2001. *Global Seagrass Research Methods*. Elsevier Science, B.V., Amsterdam, 473 pp.
- Sokal, R. R. and Rohlf, F.J., 2012. *Biometry: the principles and practice of statistics in biological research*. 4th edition. W. H. Freeman and Co.: New York. 937 pp.
- Suggett, D.J., Hall-Spencer, J.M., R Metalpa, R., Boatman, R.G., et al., 2012. Sea anemones may thrive in a high CO<sub>2</sub> world. *Global Change Bio.* 18: 3015-3025.
- Takahashi. M., Noonan. S. H. C., Fabricius. K. E., Collier. C. J., 2016. The effects of long-term in situ CO<sub>2</sub> enrichment on tropical seagrass communities at volcanic seeps. *ICES J. of Mar. Sci.* 73: 876-886.
- Valiela, I., McClelland, J., Hauxwell, J., Behr, P.J., et al., 1997. Macroalgal blooms in shallow estuaries: References and Ecophysiological Ecosystem Consequences. *Limnol. and Oceano.* 42: 1105-1118
- Varnavas, S.P., Cronan, D.S., 2005. Submarine hydrothermal activity off Santorini and Milos in the Central Hellenic Volcanic Arc: a synthesis. *Chem. Geol.* 224: 40-54.
- Vizzini, S., Tomasello, A., Maida, G.D., Pirrotta, M., et al., 2010. Effects of shallow hydrothermal seeps on the  $\delta^{13}\text{C}$  and growth performance in the seagrass *Posidonia oceanica*. *J. of Eco.* 98: 1284-1291.
- Vizzini, S., Valentina, C., Cecilia, T., Paola, G. et al., 2013. Trophic Transfer of Trace Elements in an Isotopically Constructed Food Chain from a Semi-Enclosed Marine Coastal Area (Stagnone di Marsala, Sicily, Mediterranean). *Arch. Environ. Contam. Toxicol.* 65: 642-653
- Welsh, D., Bourgues, S., de Wit, R., Auby, I., 1997. Effect of plant photosynthesis, carbon sources and ammonium availability on nitrogen fixation rates in the rhizosphere of *Zostera noltii*. *Aq. Micro. Eco.* 12:285-290
- Zimmerman. R. C., Kohrs. D.G., Steller. D.L., Alberte. R.S., 1997. Impacts of CO<sub>2</sub> enrichment on Productivity and Light requirements of Eelgrass. *Plant Physio.* 115: 599-607.

## 3.6. Appendix

## Annex-A

Response of *Cymodocea nodosa* morphometry (mean  $\pm$  S.E) at the volcanic CO<sub>2</sub> seeps and Reference sites off Greece and Italy. Kruskal- Wallis P values are presented. Different letters indicate significant difference between CO<sub>2</sub> seeps and reference sites.

	CO <sub>2</sub> seeps	Reference
	<b>Milos</b>	
Horizontal rhizome length (cm)	4.2 $\pm$ 0.52	3.4 $\pm$ 0.51
	<b>Paleochori</b>	
Horizontal rhizome length (cm)	4.8 $\pm$ 0.39	5.2 $\pm$ 0.52
	<b>Vulcano</b>	
Horizontal rhizome length (cm)	1.6 $\pm$ 0.1	1.8 $\pm$ 0.2

## Annex-B

Two-way ANOVA results of *C. nodosa* morphometry, Sample: 2 variables (CO<sub>2</sub> seeps, Reference) and Site: 3 levels (Milos (M), Paleochori (P) and Vulcano (V)) off Greece and Italy. Post hoc Tukey's test ( $p < 0.05$ ) indicate significant differences among sites. Tukey's test with no values are represented by (-).

Variable	Source of Variation	p value	Tukey's test (p values)		
			M vs P	M vs V	V vs P
No. of leaves	Sample	0.019			
	Site	<0.001	0.002	<0.001	<0.001
	Sample x site	<0.001			
Horizontal rhizome length	Sample	0.118			
	Site	<0.001	-	<0.001	<0.001
	Sample x site	-			
Vertical rhizome length	Sample	<0.001			
	Site	<0.001	1.000	<0.001	<0.001
	Sample x site	0.063			

## Annex-C

Two-way ANOVA results of *C. nodosa* density and biomass, (Sample: 2 variables (CO<sub>2</sub> seeps, Reference) and Site: 3 levels (Milos (M), Paleochori (P) and Vulcano (V)) off Greece and Italy. Post hoc Tukey's test ( $p < 0.05$ ) indicate significant differences among sites. Tukey's test with no values were represented by (-).

Variable	Source of Variation	p value	Tukey's test (p values)		
			M vs P	M vs V	V vs P
Shoot density	Sample	0.020			
	Site	0.929	-	-	-
	Sample x site	0.262			
Apex density	Sample	<0.001			
	Site	0.008	0.183	0.005	0.309
	Sample x site	0.452			
Total biomass	Sample	0.038			
	Site	<0.001	0.185	<0.001	<0.001
	Sample x site	0.019			
Leaf biomass	Sample	0.045			
	Site	<0.001	0.969	<0.001	<0.001
	Sample x site	0.062			
Vertical rhizome biomass	Sample	0.272			
	Site	<0.001	0.071	<0.001	<0.001
	Sample x site	0.136			
Horizontal rhizome biomass	Sample	0.040			
	Site	<0.001	0.461	<0.001	<0.001
	Sample x site	<0.001			
Root biomass	Sample	0.657			
	Site	<0.001	0.974	<0.001	<0.001
	Sample x site	0.950			
Above: below ground biomass	Sample	<0.001			
	Site	<0.001	1.000	<0.001	<0.001
	Sample x site	<0.001			


## Annex-D

Two-way ANOVA results of *C. nodosa* carbon (C), nitrogen (N) and C: N ratios, (Sample: 2 variables (CO<sub>2</sub> seeps, Reference) and Site: 3 levels (Milos (M), Paleochori (P) and Vulcano (V)) off Greece and Italy. Post hoc Tukey's test ( $p < 0.05$ ) indicate significant differences among sites. Tukey's test with no values were represented by (-).

Variable	Source of Variation	p value	Tukey's test (p values)		
			M vs P	M vs V	V vs P
Leaves (%C)	Sample	0.023			
	Site (CO <sub>2</sub> seeps)	<0.001	0.996	<0.001	<0.001
	Site (Reference)	<0.001	<0.001	<0.001	<0.001
	Sample x site	0.013			
Rhizomes (%C)	Sample	0.545			
	Site	0.196	-	-	-
	Sample x site	0.147			
Roots (%C)	Sample	0.176			
	Site (CO <sub>2</sub> seeps)	<0.001	<0.001	<0.001	0.235
	Site (Reference)	<0.001	0.300	<0.001	<0.001
	Sample x site	0.022			
Leaves (%N)	Sample	0.234			
	Site	<0.001	0.244	0.009	<0.001
	Sample x site	0.903			
Rhizomes (%N)	Sample	0.186			
	Site (CO <sub>2</sub> seeps)	0.042	0.043	0.289	0.003
	Site (Reference)	0.042	0.045	0.158	0.947
	Sample x site	0.003			
Roots (%N)	Sample	0.317			
	Site	0.897	-	-	-
	Sample x site	0.685			
Leaves (C: N)	Sample	0.608			
	Site	<0.001	0.003	0.711	0.002
	Sample x site	0.735			
Rhizomes (C: N)	Sample	0.059			
	Site (CO <sub>2</sub> seeps)	0.023	0.031	0.241	0.002
	Site (Reference)	0.023	0.073	0.143	0.999
	Sample x site	0.004			
Roots (C: N)	Sample	0.195			
	Site	0.343	-	-	-
	Sample x site	0.933			

Annex-E

Elongation rate of *C. nodosa* at CO<sub>2</sub> seeps and Reference sites off Greece and Italy. Error bars represent standard errors of vertical rhizome elongation rate. Different letters indicate significant differences between CO<sub>2</sub> seeps and reference sites.




## Annex-F

Two-way ANOVA results of *C. nodosa* rhizome and leaf production, (Sample: 2 variables (CO<sub>2</sub> seeps, Reference) and Site: 3 levels (Milos (M), Paleochori (P) and Vulcano (V)) off Greece and Italy. Post hoc Tukey's test ( $p < 0.05$ ) indicate significant differences among sites. Tukey's test with no values were represented by (-).

Variable	Source of Variation	p value	Tukey's test (p values)		
			M vs P	M vs V	V vs P
Vertical rhizome production	Sample	0.002			
	Site	<0.001	0.032	<0.001	0.305
	Sample x site	0.096			
Horizontal rhizome production	Sample	<0.001			
	Site	<0.001	<0.001	<0.001	0.004
	Sample x site	-			
Annual leaf production	Sample	0.012			
	Site	<0.001	0.997	<0.001	<0.001
	Sample x site	0.583			
Vertical rhizome production rate	Sample	0.002			
	Site	<0.001	0.020	<0.001	0.328
	Sample x site	0.112			
Horizontal rhizome production rate	Sample	<0.001			
	Site	<0.001	<0.001	<0.001	0.006
	Sample x site	-			



**Elevated trace elements in sediments and seagrass at six  
Mediterranean volcanic seeps**



Mishra, A.K., Santos, R., Hall -Spencer, J.M.

(To be submitted)



### **Elevated trace elements in sediments and seagrass at six Mediterranean volcanic seeps**

#### **Abstract:**

Volcanic CO<sub>2</sub> seeps are now being used as natural analogues for ocean acidification studies although these areas can be affected by trace element input and may alter ecosystem responses to gradient in carbonate chemistry. Here Fe and a range of trace elements (Cd, Co, Cu, Hg, Mn, Pb, Ni and Zn) were analysed from sediments and from the roots, rhizomes and leaves of seagrass at six CO<sub>2</sub> seeps and reference sites off Greece and Italy. There were higher metal levels in sediment and seagrasses at all CO<sub>2</sub> seeps than reference sites. Sediment Quality Guideline Quotient, a commonly used pollution index, indicated that some of the metals (Cd, Cu, Hg, Ni) were in high enough concentrations to have adverse biological effects, such as Cu at Ischia site and Hg at Vulcano. Higher accumulation of elements from sediments in roots and leaves at CO<sub>2</sub> seeps were found from Bio Sediment Accumulation Factor index. There were higher levels of Cu, Fe and Zn in leaves and rhizomes for *P. oceanica* and higher levels of Cd, Cu, Fe and Zn in *C. nodosa* compartments at CO<sub>2</sub> seeps. Fe was found with positive correlation within sediment-roots and sediment-rhizomes, whereas Cd and Pb were found with positive correlation in compartments of *C. nodosa*. In *P. oceanica* positive correlation were only observed for Cd within sediment-roots and plant compartments. Low pH and ocean acidification increased the concentration of elements at CO<sub>2</sub> seeps than reference sites. Thus, caution is needed, when using volcanic seep systems as analogue for the effects of rising CO<sub>2</sub>, as metals can reach levels that are toxic to seagrass, masking any potential benefits of increased levels of carbon dioxide for seagrass productivity.

Keywords: ocean acidification, metal toxicity, coastal habitats, diving science, CO<sub>2</sub> seeps



### 4.1. Introduction

Earth's climate is undergoing significant changes due to anthropogenic CO<sub>2</sub> emissions (IPCC, 2014). Around 30% of these CO<sub>2</sub> emissions dissolve into the surface ocean causing the pH to fall and altering seawater carbonate chemistry in a process called ocean acidification (Caldeira and Wickett, 2003). These chemical changes have raised concerns about the knock-on effects on marine organisms (Doney et al., 2009; Fabricius et al., 2014). Ocean acidification is expected to reduce marine biodiversity and alter trophic interactions (Kroeker et al., 2013; Sunday et al., 2016) which will impact a range of ecosystem services (Lemasson et al., 2017).

There are concerns that changes in ocean carbonate chemistry will also exacerbate the harmful effects of metal pollution which is a widespread problem in coastal ecosystems (Ivanina et al., 2015; Lewis et al., 2016), particularly in the Mediterranean (Soliman et al., 2015). Ocean acidification is expected to increase the bioavailability and toxicity of elements like Cd, Cu, Ni, Zn by altering their speciation (Zeng et al., 2015). However, the ecological risk posed by the effects of ocean acidification on metal contaminated sediments is difficult to assess, due to the complex behaviour of trace elements in marine environment (Millero et al., 2009).

Trace elements occur naturally in very low concentrations in the environment and can be either essential (e.g. Co, Cu, Mn) or non-essential (e.g. Cd, Cs, Sr) to living organisms (Alloway, 1995). Trace elements such as arsenic, copper, lead and mercury can be harmful to biota in coastal waters (Stumm Morgan, 1995). The toxicity of these elements depends on their form, so arsenic, for example is toxic when found in its metalloid form, whereas mercury and lead are toxic as free ions and copper is toxic when reduced to Cu(I) and is widely used in antifouling paints for this reason (Tchounwou et al., 2014).

Submarine hydrothermal activity is of ecotoxicological interest as this can cause natural gradients in both ocean acidity and trace elements, offering opportunities to assess their combined effects (Monia Renzi et al., 2011; Kadar et al., 2012; Vizzini et al., 2013). Some trace elements become more toxic if seawater pH falls from 8.1 to 7.8, which is a concern since this is the shift expected in mean surface seawater pH this century. Lead and zinc both become more toxic over this range (Millero et al., 2009; Dong et al., 2016). Trace metal toxicity depends on concentration and speciation of the element as well as interactions at receptor sites that determine uptake by organisms (Batley et al., 2004). Increased seawater acidity as well as lower oxygen levels helps release of metals that are bound to sediments (Atkinson et al., 2007).

Metal toxicity increases at submarine hydrothermal vents with copper, for example, vent fluids lower the pH which releases non-toxic Cu (II) but this is then reduced to the toxic form Cu (I) as hydrothermal fluids are low in oxygen.

Decreases in seawater pH due to increases in  $p\text{CO}_2$  levels can therefore increase the toxic concentration of certain metals in sediments (Roberts et al., 2013) and the water column (Millero et al., 2009). Ocean acidification is expected to increase the free ion concentration of Cu by 115% (Pascal et al., 2010; Richards et al., 2011), whereas the free ion concentration of other metals such as Cd may decrease or be unaffected (Lacoue-Labarthe et al., 2012; Pascal et al., 2010). Overall greater metal toxicity is predicted under ocean acidification (Lacoue-Labarthe et al., 2009; 2012; Pascal et al., 2010). Most studies on the bioavailability of trace metals at elevated  $\text{CO}_2$  have been carried out in simplified laboratory conditions (Besar et al., 2008; Richir et al., 2013; Bravo et al., 2016) so there is limited understanding of the interplay between trace elements and ocean acidification in the field.

The behaviour, speciation and bioavailability of many trace elements in seawater is strongly dependent on seawater chemistry, with Cd, Co, Cu, Hg, Ni, Pb and Zn all known to be sensitive to speciation changes within the surface seawater pH range projected due to anthropogenic  $\text{CO}_2$  emissions (Byrne et al., 1998; Richards et al., 2011). Most studies of the combined effects of ocean acidification and metal toxicities have been focused on marine fauna in laboratory settings (e.g. mussels and invertebrates) (Lewis et al., 2014; Shi et al., 2016). Lewis et al., (2014) observed toxic effects of Cu on sperm DNA damage of *Arenicola marina*, whereas Shi et al., (2016) observed damage of epithelia in *Mytilus edulis* due to Cd toxicity. There are opportunities to investigate the combined effects of metal toxicity on habitat forming primary producers since macroalgae and seagrasses can be abundant at  $\text{CO}_2$  seeps (Apostolaki et al., 2014; Vizzini et al., 2010, 2014).

Shallow water volcanic seeps are widely used as natural analogues for future coastal ecosystems as they can have areas of seabed where entire communities of marine organisms are exposed to the shifts in carbonate chemistry that are expected due to continued anthropogenic  $\text{CO}_2$  emissions (Hall-Spencer et al., 2008; Enochs et al., 2015; Connell et al., 2017). At such seeps, there are often elevated levels of hydrogen sulphide ( $\text{H}_2\text{S}$ ) and trace elements (Kadar et al., 2012; Boatta et al., 2013; Vizzini et al., 2013). Thus, care is needed when using  $\text{CO}_2$  seeps as analogues for ocean acidification due to confounding factors that may be harmful to marine biota (Barry et al., 2010). While relationships among organisms,

environmental factors and trace elements have received much attention at deep sea hydrothermal vents (Kadar et al., 2007; Cravo et al., 2007), those of shallow marine CO<sub>2</sub> seeps are still little understood. Observations at multiple shallow CO<sub>2</sub> seeps show contrasting effects on seagrasses. These plants form lush stands at some seeps, e.g. off Ischia in the Mediterranean and at several sites around Papua New Guinea (Hall-Spencer et al., 2008; Russell et al., 2013). Yet at other CO<sub>2</sub> seeps seagrasses do poorly e.g. off Panarea and Vulcano in Italy, where H<sub>2</sub>S and elevated trace elements are thought to be toxic to the plants (Vizzini et al., 2010, 2013).

Seagrass are important coastal habitats due to their high productivity and biodiversity (Thom, 2001). They provide various ecosystem services, such as food and nurseries for fish, turtles and mammals (Coles et al., 2007), they can be major carbon sinks and can sequester contaminants such as excess nutrients and metals (Orth et al., 2006; Fourqurean et al., 2012). Seagrass productivity is predicted to increase as CO<sub>2</sub> levels continue to rise if temperature increases do not become too stressful (Koch et al., 2013; Brodie et al., 2014). Seagrasses accumulate trace elements and so are used as bioindicator in coastal ecosystems (Catsiki and Panayotidis, 1993). The plants take in trace elements via the roots, rhizomes or the leaves and can translocate them between these tissue compartments (Ralph et al., 2006). The rate of uptake varies between essential and non-essential trace elements and between tissues and this introduces these substances into the food web via grazing and decomposition (Lewis and Devereux, 2009). For instance, *Cymodocea serrulata* transfers trace elements to detrital feeders (Klumpp et al., 1989).

Studies in laboratory mesocosms using high CO<sub>2</sub> have shown toxic effects of elevated concentrations of Cu, Pb and Zn on the physiology of the seagrasses *Zostera capricorni* (Ambo-Rappe et al., 2007) and *Halophila ovalis* (Ambo-Rappe et al., 2011). Although seagrass meadows are found at many CO<sub>2</sub> seeps around Greece and Italy (Hall-Spencer et al., 2008; Vizzini et al., 2010; Apostolaki et al., 2014) little is known about the influence of CO<sub>2</sub> seeps on seagrass trace metal contents. Vizzini et al., (2013) reported on Fe and trace element (As, Cd, Co, Cr, Cu, Hg, Mn, Mo, Ni, Pb, V and Zn) concentrations at Vulcano CO<sub>2</sub> seeps, where they found a wide variability in trace element concentrations in sediment and seagrass and reported on the possible bioavailability of harmful trace elements in areas with lower pH.

Here we expand on the work undertaken by Vizzini et al., (2013) to quantify the concentrations of Fe and trace elements, in sediments and seagrass at multiple seep sites around the Mediterranean. Our aim was to find out whether increased levels of trace elements near

seeps correlate with increases in trace elements in seagrass roots, rhizomes and leaves. We also test the hypothesis that low pH will increase bioavailability of certain elements and we consider whether natural CO<sub>2</sub> seeps have potential for the study of ocean acidification on metal toxicity.

### 4.2. Methods

#### *Study sites*

We surveyed six sites in the Mediterranean Sea, all of which had seagrasses (*Posidonia oceanica* or *Cymodocea nodosa*) growing on sandy bottom in high salinity, high alkalinity low nutrient conditions (Table 1). At each location, a high CO<sub>2</sub> site and a reference site was sampled between May-July, 2014. The annual temperature range was around 18-22° C for all six sites and the CO<sub>2</sub> seeps were at 1-10 m depth with a tidal range of 0.30-0.50 m.

#### *Vulcano, Italy*

We sampled Levante Bay (38.4 N, 15.0 E) off Vulcano island (Fig. 1A). Some parts of this bay are well-suited for studies of the effects of increased CO<sub>2</sub> levels (Boatta et al., 2013) despite areas with elevated H<sub>2</sub>S and metals (Vizzini et al., 2013). The main underwater gas seeps are located along southern and western shores of the bay at <1 m depth (Boatta et al., 2013). Total CO<sub>2</sub> output is about 3.6 tonne d<sup>-1</sup> (Inguaggiato et al., 2012), and the underwater gas emissions are 97-98% CO<sub>2</sub> with 2.2% H<sub>2</sub>S close to the seeps, decreasing to less than 0.005% H<sub>2</sub>S towards the north-eastern part of the bay, where most ocean acidification research has been located (Capaccioni et al., 2001; Milazzo et al., 2014). There is a step gradient in carbonate chemistry with pH 5.65 at the main gas seeps increasing to pH 8.1, which is typical for present day Mediterranean surface seawater, at >350 m from the seeps (Boatta et al., 2013). *Cymodocea nodosa* were absent at the main vents, but were collected on the periphery of the CO<sub>2</sub> seeps at 1 m depth.

#### *Ischia, Italy*

Off Ischia (Fig.1A) South of Castello Aragonese (40°43'50.4"N; 13°57'48.2"E) CO<sub>2</sub> bubbles up from the seabed in shallow water. The gas is 90–95% CO<sub>2</sub>, 3–6% N<sub>2</sub>, 0.6–0.8% O<sub>2</sub>, 0.2–0.8% CH<sub>4</sub> and 0.08–0.1% air and at ambient temperature without sulphur (Tedesco, 1996). There is a pH gradient from 6.57 to 8.17 along a 300m stretch of rocky shore with a very shallow (0.5 m depth) *Posidonia oceanica* meadow (Hall-Spencer et al., 2008) where we took our samples (Fig.2a).

### *Panarea, Italy*

Panarea island (38°38'12.2"N; 15°06'42.5"E) is to the east of the Aeolian Archipelago in the Southern Tyrrhenian Sea (Fig.1A). It is twelve miles south of the active Stromboli volcano, and emerges from a larger caldera of about 460km<sup>2</sup> that rises one thousand metres above the surrounding seabed (Gabianelli et al., 1990). On the main island and on the surrounding seafloor, tectonic faults have many gas seeps (Gabianelli et al., 1990; Voltattorni et al., 2009). These volcanic seeps have been used in several studies of the effects of CO<sub>2</sub> on the marine environment as well as field laboratories for the validation of monitoring and sampling techniques (Caramanna et al., 2011; Gofreddo et al., 2017). Here *Posidonia oceanica* was sampled at 5m depth.

### *Milos Islands, Greece*

Paleochori Bay (36.67 N, 24.51 E) and Milos thermal stations (36.70 N, 24.46 E) are part of Milos island (Fig.1B). Extensive submarine venting occurs offshore, from the intertidal to depths of more than 100 m over a 34 km<sup>2</sup> area of seabed (Dando et al.,1999). The released gases are 95% CO<sub>2</sub> with some H<sub>2</sub>S, CH<sub>4</sub> and H<sub>2</sub> (Dando et al., 1999). *Cymodocea nodosa* meadows were sampled at 4m depth at both stations (Fig.2b)

### *Methana, Greece*

The Methana peninsula (37.638428 N; 23.359730 E) is the westernmost volcanic system of the northern Aegean Volcanic Arc (Fig.1B), derived from the subduction of the African tectonic plate beneath the Eurasian plate. We sampled the area described by Baggini et al., (2014) near Agios Nikolaos village on the NE part of the peninsula. The gases were 90% CO<sub>2</sub>, with small amounts of nitrogen, carbon monoxide and methane (D'Alessandro *et al.*, 2008). Here we sampled *Posidonia oceanica* meadows at 8-10 m depth.

### *Water sampling*

Water samples (n=5) were collected at each CO<sub>2</sub> seep and Reference site in 100 ml Winkler bottles fixed with 20 µl mercuric chloride in the field, stored in dark cool- boxes and transported to the laboratory for total alkalinity (TA) analysis. The pH<sub>NBS</sub> (using pH meter, Titrimo Methron) and temperature of the water samples were measured in the field immediately after collection. In the laboratory 80 ml water samples were analysed for TA using a Lab Titrimo

analyser following methods given by Dickson et al., (2007). Temperature,  $\text{pH}_{\text{NBS}}$  and TA data were used to calculate  $p\text{CO}_2$  using  $\text{CO}_2\text{SYS}$  program following methods given by Pierrot and Wallace, (2006).

### *Sediment & seagrass sampling*

Sediment samples were collected from  $\text{CO}_2$  seep and Reference sites by SCUBA diving using a 10-cm long syringe with the tip cut off to collect the upper 5 cm of sand. The sediment samples were in Teflon tubes in dark boxes then analysed for the grain size following dry sieving at Half Phi intervals (Blott and Pye, 2001).

Samples of *Cymodocea nodosa* (from Vulcano, Milos and Paleochori islands) and *Posidonia oceanica* (from Ischia, Panarea and Methana) were collected by SCUBA diving at up to 10-meter depth at the  $\text{CO}_2$  seep and Reference sites. The seagrasses were rinsed to remove sediments, then air-dried and stored in polybags until analyses. *Cymodocea nodosa* and *Posidonia oceanica* leaves were scraped to remove epiphytes. Sediment samples and seagrass leaves, roots and rhizomes were oven dried at  $40^\circ\text{C}$  and powdered in a mortar.

### *Analytical Methods*

Total trace element concentrations were determined by Aqua Regia Soluble Total method. Dried sediment (0.25 g) was put into digestion tubes (Tecator type). Cold and concentrated acids in the order: 4.5 mL Hydrochloric acid (HCl): 1.5 mL Nitric acid ( $\text{HNO}_3$ ) was added to the tubes. The digestion tubes were left to pre-digest, for one hour then heated for 2 hours at  $95 - 100^\circ\text{C}$ . After cooling, the digest was filtered quantitatively into a volumetric flask and diluted using 2%  $\text{HNO}_3$  (25 ml volume).

For dried seagrass (leaves, rhizomes and roots) samples, 0.25g of sample was dissolved in 6mL of  $\text{HNO}_3$  and the volume made up to 25ml. Similarly, blanks and standards (Laboratory of the Government Chemist (LGC Reference Materials, UK) used for sediments and plants were prepared using the same method. Analysis of Cd, Co, Cu, Fe, Hg, Mn, Pb and Zn was performed using an ICP-MS (Thermo Scientific, iCAP 7000 Series) and an ICP-AES (Thermo Scientific, X Series-2) in duplicate.

*Data Analysis*

To assess the sediment quality of all six locations we used Sediment Quality Guidelines Quotient (SQG-Q, Long and MacDonald, 1998). Among the environmental quality indices in the literature, this was chosen for its simplicity, comparability and robustness as reported by Caeiro et al., (2005). The SQG-Q consists of two values: a threshold effects level (TEL) and a probable effect level (PEL) (MacDonald et al., 1996). TEL represents concentrations below which adverse biological effects are unlikely.

The SQG-Q was calculated as follows:

$$\text{SQG-Q} = (\sum_{i=1}^n \text{PEL-Q}_i) / n$$

Where  $\text{PEL-Q}_i = \text{contaminant}/\text{PEL}$ . The  $\text{PEL-Q}_i$  represents the probable effect level quotient (PEL-Q) of the  $i$  contaminant. Using the SQG-Q index, we put sediments into three categories as established by MacDonald et al. (2000).  $\text{SQG-Q} \leq 0.1$ - low potential for adverse biological effects;  $0.1 < \text{SQG-Q} < 1$ - moderate potential for adverse biological effects;  $\text{SQG-Q} \geq 1$ - high potential for adverse biological effects. Our data were compared with a biological effect database for sediments (BEDS) developed by Mac Donald et al. (1996), which comprises of 350 reports on the analysis of toxicity and/or benthic community composition of marine and estuarine sediments.

To understand bio-accumulation of elements, we calculated the Bio Sediment Accumulation Factor (BSAF), which is defined as the ratio between metal concentration in the organism and that in the sediment (Lau et al., 1998; Szefer et al., 1999), given by:

$$\text{BSAF} = M_p / M_s$$

Where  $M_p$  is the concentration of the element in the seagrass and  $M_s$  is the concentration of the element in the sediment (Fergusson. 1990). BSAF is an important factor in understanding how the increase or decrease of trace element concentrations affects the fate of trace elements in the plant compartments. Roots (the usual point of uptake) and leaves (the usual point of release) of trace elements (Weis & Weis, 2004) were used for BSAF analysis.

*Statistics*

A three-way ANOVA was used to analyse the variability in trace element concentration among plant compartments (leaves, rhizomes, roots), sediments and sites (reference, CO<sub>2</sub> seeps) between three locations (Ischia, Panarea and Methana) for *Posidonia oceanica* and Milos, Paleochori and Vulcano for *Cymodocea nodosa*. All data was pre-checked for normality and homogeneity of variances, when variances were not homogenous, data was  $\ln(x+1)$  transformed. After transformation, some data was still not showing homogeneity of variances. In these cases, we analysed the data untransformed, since ANOVA is considered robust to lack of homogeneity of variances with balanced designs and considerable large amount of treatments (Underwood, 1997). In some cases, ANOVA main effect was difficult to interpret due to the presence of statistically significant interactions, but in mixed effect ANOVA's the test of the fixed main effect is potentially interpretable even in the presence of interaction (Quinn and Keough.2002). Holm-Sidak test was performed for a *posteriori* comparisons among levels to check significant main effects in ANOVA. Pearson's correlation co-efficient was applied to identify correlations between trace element concentration in sediment and seagrass compartments, after testing for normality of distribution on raw or log transformed data. When normality was not achieved, non-parametric Spearman's rank correlation coefficient was applied. All statistical tests were conducted with a significance level of  $\alpha = 0.05$  and data were reported as mean  $\pm$  standard error (SE).

**4.3. Results**

As expected  $p\text{CO}_2$  concentration were highest (and pH lowest) at each of the seeps whereas reference sites had normal CO<sub>2</sub> and pH levels (Table 1). Salinity, temperature and total alkalinity were not affected by the seeps.

Grain size analysis showed that 99% of the particles sampled at all locations were sand in the very coarse, coarse, medium and fine fractions. Sediment samples from the seeps usually had higher levels of iron and several trace elements than at reference sites, but most were in such low concentrations that they were unlikely to be toxic (Table 2). Exceptions were Hg at Vulcano, Cu at Ischia as well as Ni and Cd at Paleochori (Table 2). These metals were at high enough concentrations in sediments at the seeps to likely have adverse biological effects on biota such as seagrasses. Fe, Co and Mn are excluded from these results as they have no SQG-

## Chapter 4

Q values (Mac Donald et al., 1996) to compare with our results. Considering the possible moderate or adverse biological impacts of trace elements on seagrass rest of the results and discussion are based on the elements presented in Table 2 and iron as it is needed in much of the metabolic activities in seagrass.

Table 1: Seawater salinity, temperature, total alkalinity, pH and  $p\text{CO}_2$  values (mean  $\pm$  SE, n=5) at six Mediterranean  $\text{CO}_2$  seeps.

Site	Salinity (psu)	Temp.(°C)	pH <sub>NBS</sub>	TA ( $\mu\text{mol Kg SW}^{-1}$ )	$p\text{CO}_2$
<b>Vulcano</b>					
Reference	35.8	21.6	8.17 $\pm$ 0.05	2439	427 $\pm$ 6.8
$\text{CO}_2$ seep	35.8	22.4	7.98 $\pm$ 0.08	2432	2928 $\pm$ 15.8
<b>Ischia</b>					
Reference	35.6	17.7	8.19 $\pm$ 0.06	2596	428 $\pm$ 2.3
$\text{CO}_2$ seep	35.7	17.8	7.78 $\pm$ 0.05	2589	1653 $\pm$ 10.2
<b>Panarea</b>					
Reference	36.0	20.5	8.18 $\pm$ 0.05	2507	420 $\pm$ 4.6
$\text{CO}_2$ seep	36.0	22.3	7.47 $\pm$ 0.04	2500	3370 $\pm$ 2.3
<b>Milos</b>					
Reference	36.7	22.6	8.2 $\pm$ 0.03	2715	405.5 $\pm$ 1.6
$\text{CO}_2$ seep	36.7	23.5	7.5 $\pm$ 0.04	2704	2457.9 $\pm$ 1.8
<b>Paleochori</b>					
Reference	36.0	22.6	8.2 $\pm$ 0.01	2711	402.9 $\pm$ 1.1
$\text{CO}_2$ seep	36.0	22.8	7.9 $\pm$ 0.01	2706	1884.3 $\pm$ 3.0
<b>Methana</b>					
Reference	36.8	22.8	8.2 $\pm$ 0.01	2715	460 $\pm$ 6.9
$\text{CO}_2$ seep	36.8	23.0	7.8 $\pm$ 0.02	2704	1980 $\pm$ 4.4

Table 2. Sediment Quality Guidelines-quotient (SQG-Q) of sediment calculated with Probable Effects Level for Reference and CO<sub>2</sub> seep sites in Greece and Italy. SQG-Q <0.1 (low effect), <0.1 SQG-Q>1 (moderate effect), SQG-Q>1 adverse biological effects. Numbers and letters in bold indicate possible adverse effects of trace elements.

Location	Element	SQG-Q		Effects	
		Reference	CO <sub>2</sub> seeps	Reference	CO <sub>2</sub> seeps
Vulcano	Cu	0.08	0.33	Low	Moderate
	Hg	0.32	<b>1.18</b>	Moderate	<b>Adverse</b>
	Ni	0.13	0.21	Moderate	Moderate
	Zn	0.09	0.13	Low	Moderate
Ischia	Cu	0.93	<b>1.06</b>	Moderate	<b>Adverse</b>
	Hg	0.64	0.86	Moderate	Moderate
	Pb	0.11	0.13	Moderate	Moderate
	Zn	0.12	0.10	Moderate	Moderate
Panarea	Cd	0.10	0.16	Low	Moderate
	Cu	0.06	0.11	Low	Moderate
	Hg	0.79	0.84	Moderate	Moderate
	Ni	0.03	0.18	Low	Moderate
	Pb	0.09	0.57	Low	Moderate
	Zn	0.05	0.12	Low	Moderate
Milos	Cd	0.21	0.21	Moderate	Moderate
	Ni	0.31	0.41	Moderate	Moderate
Paleochori	Cd	0.04	<b>1.84</b>	Low	<b>Adverse</b>
	Ni	0.71	<b>1.01</b>	Moderate	<b>Adverse</b>
Methana	Ni	0.11	0.16	Moderate	Moderate
	Pb	0.05	0.42	Low	Moderate

Table 3. Three-way ANOVA differences in Fe and trace element levels between Sample:2 variables (CO<sub>2</sub> seeps, Reference)) Site: 3 levels (Methana (M), Panarea(P) and Ischia (V)) and compartments :4 levels (Sediments (Sd), Rhizomes (Rh), Roots (R), Leaves (L)). Holm-Sidak significant test (p<0.05) is presented for locations, sediment and *P. oceanica* compartments. Numbers (in bold) indicate differences that were not significant.

Element	Variation	p value	Holm-Sidak p values								
			Location			Sediment vs Compartment			Compartments		
			M vs P	M vs V	V vs P	Sd vs R	Sd vs Rh	Sd vs L	R vs Rh	Rh vs L	R vs L
Fe	Sample	<0.001	<0.001	<0.001	<0.001						
	Site	<0.001									
	Compt.	<0.001				<0.001	<0.001	<0.001	<0.001	<0.001	<0.001
Cd	Sample	<0.001	<0.001	<0.001	<0.001						
	Site	<0.001									
	Compt.	<0.001				<0.001	<0.001	<0.001	<0.001	<0.001	<0.001
Cu	Sample	<0.001	<0.001	<0.001	<0.001						
	Site	<0.001									
	Compt.	<0.001				<0.001	<0.001	<b>0.314</b>	<0.001	<0.001	<0.001
Ni	Sample	<0.001	<0.001	<0.001	<0.001						
	Site	<0.001									
	Compt.	<0.001				<0.001	<0.001	<0.001	<0.001	<0.001	<0.001
Pb	Sample	<0.001	<0.001	<0.001	<0.001						
	Site	<0.001									
	Compt.	<0.001				<0.001	<0.001	<0.001	<0.001	<0.001	<b>0.652</b>
Zn	Sample	<0.001	<0.001	<0.001	<0.001						
	Site	<0.001									
	Compt.	<0.001				<b>0.222</b>	<0.001	<0.001	<0.001	<0.001	<0.001

Table 4. Three-way ANOVA differences in Fe and trace element levels between Sample:2 variables (CO<sub>2</sub> seeps, Reference) Site: 3 levels (Milos (M), Paleochori (P) and Vulcano (V)) and compartments: 4 levels (Sediments (Sd), Rhizomes (Rh), Roots (R), Leaves (L)). Holm-Sidak significant test (p<0.05) was significant is presented for locations, sediment versus plant compartments and *C. nodosa* compartments. Numbers (in bold) indicate differences that were not significant

Element	Variation	p value	Holm-Sidak p values			Sediment vs Compartment			Compartments		
			Location			Sd vs R	Sd vs Rh	Sd vs L	R vs Rh	Rh vs L	R vs L
			M vs P	M vs V	V vs P						
Fe	Sample	<0.001	<0.001	<0.001	<0.001						
	Site	<0.001									
	Compt.	<0.001				<0.001	<0.001	<0.001	<0.001	<0.001	<0.001
Cd	Sample	<0.001	<0.001	<0.001	<0.001						
	Site	<0.001									
	Compt.	<0.001				<0.001	<0.001	<0.001	<0.001	<b>0.787</b>	<0.001
Cu	Sample	<0.001	<b>0.626</b>	<0.001	<0.001						
	Site	<0.001									
	Compt.	<0.001				<0.001	<b>0.621</b>	<0.001	<0.001	<0.001	<0.001
Ni	Sample	<0.001	<0.001	<b>0.853</b>	<0.001						
	Site	<0.001									
	Compt.	<0.001				<0.001	<0.001	<0.001	<0.001	<0.001	<0.001
Zn	Sample	<0.001	<0.001	<0.001	<0.001						
	Site	<0.001									
	Compt.	<0.001				<0.001	<0.001	<0.001	<0.001	<b>0.910</b>	<0.001




Fig.1. Study sites in Italy a) and b) Greece, showing reference and CO<sub>2</sub> seep sites which were all sampled between May to July 2014.

**a****b**

Fig. 2. a) *Posidonia oceanica* and b) *Cymodocea nodosa* meadows at CO<sub>2</sub> seeps off Ischia (Italy) and Paleochori (Greece)

Fe and trace element levels within sediments at all six CO<sub>2</sub> seeps were higher at seeps than reference sites, except at Methana (Figs. 3 and 4). Wide variation in concentration levels of Fe and trace elements were observed at all CO<sub>2</sub> seeps. Paleochori seep sediments were more significantly enriched with Cd than rest of the seeps. Sediments off Ischia had the highest Cu concentration among all seep locations, but this higher concentration was not reflected in seagrass compartments, so this copper may not be bioavailable. Mercury was only found in seep sediments off Italy, with highest level observed at Vulcano. Sediments at Paleochori seeps had a significantly higher concentration of Ni than at reference site, whereas all other CO<sub>2</sub>

seeps had lower levels. Zinc concentration in sediments were similar at Vulcano, Milos, Ischia and Panarea seeps, but were low at Panarea. Levels of Pb were high and significant in the sediments of Ischia CO<sub>2</sub> seeps than all other seeps, but this was not reflected in the seagrass compartments (Figs.3 and 4).

Fe was significantly different within with the roots, rhizomes and leaves in *P. oceanica* at all three CO<sub>2</sub> seeps except in leaves at Ischia and Panarea (Fig.3). Fe was observed with highest concentration followed by Cu and Zn in *P. oceanica* compartments (Fig.3). Trace element concentration were high and significant within roots (Cd, Cu, Hg and Zn), rhizomes (Cd, Cu, Hg, Ni and Zn) and leaves (Co, Cu, Hg and Zn) of *P. oceanica* at Ischia than Panarea and Methana respectively (Fig.3). Variation in concentration of Fe and trace elements within location (Ischia vs Panarea, Ischia vs Methana and Methana vs Panarea) were significant at all three locations (Table 3). The concentration of elements between sediments and those found in plant compartments were significantly different at all three locations except for Cu (sediment-leaves) and Zn (sediment-roots). Within *P. oceanica* compartments all elements analysed were significantly different except for Pb (roots-leaves) at all three seeps (Table 3). The variation in concentration of elements were higher in roots followed by leaves and rhizomes in *P. oceanica* at the CO<sub>2</sub> seeps off Ischia, Panarea and Methana (Fig.3).

Iron levels were significantly different within the roots, rhizomes and leaves of *C. nodosa* at Vulcano, in rhizomes at Paleochori and in roots at Milos CO<sub>2</sub> seeps (Fig. 4). Trace elements concentration were high and significant within the roots (Cd, Hg, Ni and Zn), rhizomes (Hg and Zn) and leaves (Cd, Cu, Hg, Ni and Zn) of *C. nodosa* at Vulcano than at Milos and Paleochori (Fig.4). Variation in Fe and trace elements concentration within locations (Milos vs Paleochori, Paleochori vs Vulcano, Vulcano vs Milos) was significant except for Cu at Milos vs Paleochori, Ni at Vulcano vs Milos (Table 4). Element concentrations found in sediment and that in *C. nodosa* compartments were different significantly except for Cu (sediment vs rhizomes) at all three locations (Table 4). Within the plant compartments (roots vs rhizomes, rhizomes vs leaves, roots vs leaves) Fe and trace elements concentration were significantly different except Zn (rhizomes vs leaves) and Cd (rhizomes vs leaves) at all three locations (Table 4). For *C. nodosa*, variation in concentration of elements were high and significant in roots followed by rhizomes and leaves off Milos, Paleochori and Vulcano CO<sub>2</sub> seeps (Fig.4). Iron had the highest concentration followed by Cu and Ni in *C. nodosa* compartments (Fig. 4).




Fig. 3. continued




Fig.3. continued




Fig. 3. Element concentrations (mean  $\pm$  SE,  $n=5$ ) for Fe, Cd, Co, Cu, Hg, Mn, Ni, Pb and Zn for *Posidonia oceanica* in sediments and within plant compartments at reference and CO<sub>2</sub> seep sites off Italy and Greece. Error bars represent standard error. Different letters indicate significant difference between reference and CO<sub>2</sub> seeps site for each location.




Fig. 4. continued




Fig. 4. continued







Fig. 4. Element concentration (mean  $\pm$  SE, n=5) for Fe, Cd, Co, Cu, Hg, Mn, Ni, Pb and Zn for *Cymodocea nodosa* in sediments and within plant compartments at reference and CO<sub>2</sub> seeps off Italy and Greece. Error bars represent standard error. Different letters indicate significant difference between reference and CO<sub>2</sub> seep sites for each location.




a. Vulcano




d. Ischia




b. Milos



e. Panarea



c. Paleochori



f. Methana

Fig. 5. Mean values of Bio-Sediment Accumulation Factor of trace elements for *Cymodocea nodosa* at reference and CO<sub>2</sub> seeps at Vulcano (a), Milos (b) and Paleochori (c) and for *Posidonia oceanica* at Ischia (d), Panarea (e) and Methana (f) at Italy and Greece respectively. Reference sites are represented as Ref. and CO<sub>2</sub> seeps as seeps for each location.

Bio Sediment Accumulation Factor index indicated that both *C. nodosa* and *P. oceanica* roots and leaves accumulated higher trace element concentrations than those found in sediments at CO<sub>2</sub> seeps, though different element was accumulated by each seagrass. For *P. oceanica* mean BSAF>1 was observed in roots and leaves at Panarea and Methana and in leaves at Ischia (Fig.5). In *P. oceanica* Cd, Cu, Hg, Ni, Pb and Zn were observed with BSAF>1 in the roots or leaves at all three CO<sub>2</sub> seeps. For *C. nodosa* BSAF>1 were observed for roots at Milos and Paleochori and for leaves at Vulcano. Trace elements (Cd, Cu, Ni and Zn) were observed with BSAF >1 in the roots and leaves of *C. nodosa* at all three CO<sub>2</sub> seeps (Fig.5). Most elements (4/9) with BSAF>1 were found at CO<sub>2</sub> seeps off Vulcano and Panarea for *C. nodosa* and *P. oceanica* respectively (Fig.5).

Correlations between metal content in sediments and those recorded in *P. oceanica* compartments were significant and positive for Zn and Ni in rhizomes at Ischia and Panarea seeps respectively (Table 5). Correlations were negative for Ni and Cd in rhizomes at Methana and Panarea seeps respectively (Table 5). The correlation for element content between any two organs of *P. oceanica* were positive and significant for Cd and negative for Cu at Ischia CO<sub>2</sub> seeps (Table 5).

Correlations of Fe content in sediment and those found in rhizomes of *C. nodosa* at Vulcano were significant and positive, whereas significant and negative correlation were observed for Zn content found between sediment and rhizomes and leaves (Table 6). For any two plant organs Cd was found with positive co-relation at Vulcano, where significant and negative correlation was observed for Cu and Hg at Vulcano and Fe and Cu at Milos (Table 6).

Correlation between elements and plant compartments of both seagrass indicate an absence of common trend for these elements between the roots, rhizomes and leaves of seagrasses at CO<sub>2</sub> seeps.

## Chapter 4

Table 5. Correlation between trace elements in sediments versus *P. oceanica* roots, rhizomes and leaves and plant compartments at Mediterranean CO<sub>2</sub> seeps. The correlation co-efficient (r) and significance level (p) are presented. Numbers in bold indicate significant correlation, only trace elements with significant correlations are shown

Location	Element	Sediment-roots		Sediment-rhizomes		Sediment-Leaves	
		r	p	r	p	r	p
Ischia	Zn	-0.234	0.704	0.870	<b>0.048</b>	0.321	0.598
Panarea	Cd	0.841	<b>0.014</b>	-0.910	<b>0.032</b>	-0.064	0.918
	Ni	-0.358	0.554	0.884	<b>0.046</b>	-0.787	0.114
Seagrass compartments							
Location	Element	Roots-Rhizomes		Roots-Leaves		Rhizomes-Leaves	
		r	p	r	p	r	p
Ischia	Cd	0.300	0.683	0.975	<b>0.016</b>	0.359	0.517
	Cu	-0.273	0.657	0.877	<b>0.049</b>	-0.577	0.308

Table 6. Correlation between trace elements in sediment versus *C. nodosa* roots, rhizomes and leaves and between plant compartments at Mediterranean CO<sub>2</sub> seeps. The correlation co-efficient (r) and significance (p) level are presented. Numbers in bold indicate significant correlations, only trace elements with significant co-relation are shown

Location	Element	Sediment-roots		Sediment-rhizomes		Sediment-Leaves	
		r	p	r	p	r	p
Vulcano	Fe	0.437	0.462	0.992	<b>0.000</b>	-0.836	0.078
	Zn	-0.795	0.108	-0.966	<b>0.007</b>	-0.906	<b>0.034</b>
Seagrass Compartments							
Location	Element	Roots-Rhizomes		Rhizomes-Leaves		Roots-Leaves	
		r	p	r	p	r	p
Vulcano	Cd	1.000	<b>0.016</b>	0.158	0.783	0.158	0.783
	Cu	-0.986	<b>0.002</b>	0.534	0.354	-0.620	0.265
	Hg	-0.135	0.783	0.216	0.683	-0.947	<b>0.016</b>
Milos	Fe	-0.975	<b>0.016</b>	-0.300	0.683	0.205	0.683
	Cu	-0.872	0.083	-1.000	<b>0.016</b>	0.872	0.083

### 4.4. Discussion

The six volcanic CO<sub>2</sub> seeps that we surveyed in Greece and Italy, all had higher trace element concentrations in sediments than at reference sites. This is to be expected, as volcanic seeps emit fluids often rich in CO<sub>2</sub>, H<sub>2</sub>S and precipitates of trace metals that accumulate in sediments (Aiuppa et al., 2000; Sternbeck et al., 2001). Levels of metals at volcanic seeps can be harmful to marine life (Vizzini et al., 2013), so most work using seeps as analogues for ocean acidification has been carried out on hard substrata away from harmful levels of H<sub>2</sub>S and metals (Fabricius et al., 2014; Milazzo et al., 2016). As seagrass colonise sediments we aimed to find out why they thrive at certain CO<sub>2</sub> seeps but not at others, as this may relate to the effects of pH on the bioavailability of metals that can be toxic.

In shallow marine CO<sub>2</sub> seeps off Greece (Milos, Paleochori and Methana) and Italy (Vulcano, Ischia and Panarea) redox and pH conditions are altered by the emission of CO<sub>2</sub> and H<sub>2</sub>S (Sedwick et al., 1996; Boatta et al., 2013). This alters adsorption/desorption phenomena of elements in sediments and water column (Prokop et al., 2003). Our surveys showed clear enrichment of Fe and all trace element (Cd, Cu, Hg, Ni, Pb and Zn) levels in sediments, which we expected as volcanic seeps are usually enriched with trace elements (Monia-Renzi et al., 2011). Our Italian CO<sub>2</sub> seep sediment samples had higher trace element concentrations than those in Greece. The concentrations that we found in the sediments at the Vulcano seeps were in the same range as those found by Vizzini et al., (2013). At Panarea, Pb concentrations were lower than those recorded by Monia-Renzi et al., (2011), which is probably because metal concentrations are heterogeneous around seeps and depend on highly variable redox and pH conditions. Mercury was found at all three seep locations off Italy, but was not detected at the Greek seeps confirming that not all seeps emit mercury (Stoffers et al., 1999).

Comparing trace element concentrations of sediments from the six CO<sub>2</sub> vent locations of Greece and Italy with SQGs (Long et al., 1998; MacDonald et al., 2000), the obtained values were higher than TEL and ERL values and remained lower or equal to PEL and ERM values. Based on SQGs, element such as Cd at Paleochori, Hg at Vulcano and Cu at Ischia were found at levels that are likely to have adverse biological impacts, whereas at Methana and Panarea none of the trace elements were at high enough levels to be likely to have adverse biological impacts. Our results have similar values for Ni from Vulcano as found by Vizzini et al., (2013) from SQGs indices. The moderate or adverse biological impacts of the sediments depends on the concentration of element in the sediments and the metal speciation form they are present.

Higher number of elements with adverse impacts at the CO<sub>2</sub> seeps than reference sites observed in our results indicates that metal speciation is affected by low pH resulting in increased availability along with the metal enrichment at CO<sub>2</sub> seeps.

To understand if sediments enriched with trace elements were also available for uptake, the upper 5 cm of the sediments were sampled where the root/rhizome mat system of both *C. nodosa* and *P. oceanica* are present that takes active part in metal uptake and thus keeps this layer of sediment oxygenated for various redox reactions (Kristensen, 2000). Secondly trace metals that are easily available for uptake are dominant in this upper layer (Pekey et al., 2004; Atkinson et al., 2007) increasing the bioavailability of these elements both in sediment pore water and water column (Atkinson et al., 2007). The uptake of various elements in both seagrass compartments from sediments were reflected from bio-sediment accumulation factor index. BSAF which indicates higher/lower element concentration in plant compartments than observed in sediments, showed BSAF>1 for different trace element (Cd, Cu, Ni, Pb and Zn) for *P. oceanica* and for *C. nodosa* (Cd, Cu and Ni) within sediment-roots and sediment-leaves at all six CO<sub>2</sub> seeps. Higher variation in accumulation were observed in *P. oceanica* than *C. nodosa* indicating that *P. oceanica* have greater capability than *C. nodosa* for accumulation of these trace elements (Lafabrie et al., 2007; Bonanno et al., 2016 and references there in) from sediments at the CO<sub>2</sub> seeps.

Considering the capability of both seagrasses to accumulate elements from CO<sub>2</sub> seep sediments, the principal uptake rate of elements in both seagrasses were the sediment-root and leaf-roots pathways. The extent to which these uptake methods are active or passive depends on the physiological regulation of the seagrass and the element concentration in the surrounding to which seagrass are exposed to (Lafabrie et al., 2007; Bonanno et al., 2016 and references there in) which was observed in our studies, higher Zn concentration in sediments were reflected in the leaves of *C. nodosa* at Vulcano CO<sub>2</sub> seeps. The high accumulation patterns of trace elements in the roots and leaves of *P. oceanica* and *C. nodosa* suggests a mixed tolerance strategy, based on root accumulation and losing leaves. Root accumulation are common in rooted species, both terrestrial and aquatic, where bulk of the trace elements were stored for protection against the adverse effects on photosynthetic process (Bonanno & Di Martino, 2016 and references therein). The other mechanism involves accumulation of toxic elements in rhizomes, which doesn't participate actively in photosynthetic processes, similar pattern of accumulation was observed for Ni in *C. nodosa* rhizomes at Milos and *P. oceanica* rhizomes

at Panarea seeps (Figs. 3 and 4). Malea and Haritnoids, (1999) and Bonanno & Di Martino, (2016) observed this possible tolerance mechanism in *C. nodosa*, whereas Di Leo et al., (2013) and Richir & Gobert, (2016) observed it in *P. oceanica*. They pointed out that active mobilization of metals like Cd, Ni and Pb from roots to shoots in the seagrass, facilitated metal loss in both seagrasses due to the high turnover rate of leaves. Uptake kinetics and passive absorption properties of leaves also differ from roots and element absorbed can be internally redistributed through active or passive transport mechanisms (Bonanno & Di Martino, 2016; Richir & Gobert, 2016).

Though the element concentration were higher in the sediments at CO<sub>2</sub> seeps, all elements are not preferably accumulated from sediments through the sediment-root pathways suggesting element specific uptake by seagrass. We found there was no positive correlation for Cd in the sediment-roots, sediment-rhizomes and sediment-leaves for *P. oceanica* (Table 5) except for *P. oceanica* at Panarea CO<sub>2</sub> seeps, indicating that Cd is not transferred from sediments into the *P. oceanica* compartments (Lafabrie et al., 2007; Di Leo et al., 2013). However, positive correlation for Cd in leaves-roots for Ischia (Table 5) and roots-rhizomes for Vulcano, indicate the preferable route for Cd transfer in *P. oceanica* and *C. nodosa* is by water column through leaves and then are translocated to roots (Lafabrie et al., 2007; Di Leo et al., 2013). This indicates efficient distribution pattern of Cd at CO<sub>2</sub> seeps from roots-rhizomes-leaves as found by Malea et al., (1994) in *P. oceanica* off Greece. The negative correlation for Cu, Ni and Zn found in the *P. oceanica* compartments (Table 5) indicate that though these elements are present in higher concentrations in the sediment but they do not get transferred between the root-rhizome-leaves pathway. However, these elements are essential elements and required for *P. oceanica* in biochemical pathways indicating absorption from the water column at the CO<sub>2</sub> seeps (Di Leo et al., 2013). This explains higher concentration of these trace elements in the leaves of *P. oceanica* (Fig.2) and involvement of *P. oceanica* in cycling of these elements (Lafabrie et al., 2007; Di Leo et al., 2013).

Significant positive correlation was observed in *C. nodosa* for Fe in sediment-rhizomes (Table 6). This indicates mobility of Fe from sediment to the *C. nodosa* compartments as observed previously for Fe by Malea et al., (1999). Positive or negative correlation between *Cymodocea nodosa* compartments indicate that, *C. nodosa* have effective active uptake kinetics to translocate trace elements to entire plant system. This was observed by Malea et al., (1999) for Cd, Cu and Pb and by Ahmad et al., (2015) for Cd and Cu in *C. serrulata*. Mobility of Zn

within *C. nodosa* compartments have a positive correlation, which agrees with the findings from Malea et al., (1999) for Zn. Hg was found with negative correlation in *C. nodosa* compartments at Vulcano, indicating Hg being toxic is not allowed for translocation within the seagrass compartments, similar results were found for *C. nodosa* by Sanchiz et al., (2001).

Correlation data between seagrass compartments are different because element accumulation patterns in seagrass are governed by multitudes of factors (Llagostera et al., 2011) as compared to land plants where metal uptake happens by two phases, one from sediments to roots and secondly from roots to other plant parts (Kvesitadze et al., 2009). So, in our studies at CO<sub>2</sub> seeps increased levels of elements in seagrass roots, rhizomes and leaves indicate the capacity of seagrass to absorb element simultaneously from sediments and water, as they are always submerged. This also indicates seagrass at these CO<sub>2</sub> seeps have adapted to the variation in element concentrations found in sediments at these CO<sub>2</sub> seeps with an effective mechanism of survival. Based on element concentrations, *P. oceanica* and *C. nodosa* can regulate their gene expression for metal toxicity which was observed for *P. oceanica* by Lauritano et al., (2015) and *C. nodosa* by Olive et al., (2017).

At CO<sub>2</sub> seeps the sediments receive a higher input of elements from the emission of seep fluids and thus there is a possibility of increase of element concentration in the water column too. Generally dissolved metal concentrations in overlying waters are low due to low solubility of metals in seawater and absorption on sediments. But this is going to change in near future due to ocean acidification and low pH as metal speciation is influenced by pH variation and low pH favours release of metals from pore waters (Simpson et al., 2004; Atkinson et al., 2007) that are generally not bioavailable. Once emitted from the seeps metals are absorbed on sediments where biological and chemical process results in redox stratification of metal bound particles with depth (Eggleton et al., 2004; Basallote et al., 2014), until resuspension of these particles happens due to physical processes and bioturbation. Resuspension of these metal bound particles with oxygenated overlying waters results in metal speciation in the dissolved phase (Simpson et al., 2002), making the metals bioavailable in pore waters (Simpson and Batley, 2003). Once released from pore waters into water column, these metals are bioavailable to aquatic organisms till precipitation of these metals are initiated by the fine fraction (<63 micron) of the sediments suspended in water column (Zoumis et al., 2001; Fan et al., 2002).

Concentration of Cu, Ni, Cd, Pb, Zn, increases with low pH as these metals are released from sediments at low pH (Atkinson et al., 2007; Basallote et al., 2014). Low pH also favours the release of elements like Hg from the sediment surface layers bound to sulphides and free sulphides are oxidised in the sediment upper layer by oxygen provided by seagrass mat system to release Hg. The chemical form in which the metal is present (bound to organic or inorganic compounds) in the sediment is a key issue determining its bioavailability. Metals like Cu which can be associated with minerals in low pH conditions are not likely to be released into water column, since they are trapped within the crystal structure of minerals, remaining unavailable to aquatic system (De Orte et al., 2014), whereas acidification of seawater near the CO<sub>2</sub> seeps tends to release the metals that are less strongly associated with sediments increasing their potential bioavailability (Riba et al., 2004). Thus, acidification due to low pH can increase the concentration of dissolved metals, which could affect the toxicity of the sediment to the exposed organism i.e., by increasing metal (Cu, Cd and Zn) bioavailability (Basallote et al., 2014).

In our research, all the six CO<sub>2</sub> seeps had low pH (7.4-7.9) conditions which increased the availability of Fe and Cd, Cu, Ni, Pb and Zn in their free ion forms and hence more availability and toxicity depending on the organisms (Roberts et al., 2013). However, Fe and trace elements (Cd, Cu Hg, Ni, Pb and Zn) toxicity have never been studied on *P. oceanica*, but a few studies have been conducted on *Cymodocea serrulata* (Prange and Dennison, 2000), *Halophila ovalis* and *H. spinulosa* (Prange and Dennison, 2000; Ambo-Rappe et al., 2011). Many of the concentrations found in our research for Cu and Pb are significantly higher than the toxic range previously reported for various seagrasses, affecting their photosynthesis as well as root and leaf structures (Ambo -Rapee et al., 2011; Prange and Dennison, 2000). In our studies toxicity tests were not performed, but the levels of trace elements that we found suggest possible toxic effects on seagrasses at CO<sub>2</sub> seeps (Richir, 2016). This certainly helps explain why seagrasses grow well and can take advantage of elevated CO<sub>2</sub> levels at some seeps but not at others.

To sum up: ocean acidification will affect trace metal chemistry, speciation and their bio-availability (Millero et al., 2009; Zeng et al., 2015) and may impact seagrass ecosystem in areas with contaminated sediments.

Natural CO<sub>2</sub> seeps are now widely used in global change studies, but our studies show CO<sub>2</sub> seeps off Italy and Greece have moderate to high metal levels that seagrasses tend to

accumulate. Organisms that eat seagrass may accumulate and transfer trace elements resulting in toxic effects (Bouchon et al., 2016). Therefore, while considering CO<sub>2</sub> seeps as field laboratories for global change studies, the effect of trace element contamination should be kept in mind (Vizzini et al., 2013).

We have shown that Mediterranean CO<sub>2</sub> seep sites consistently have elevated levels of trace elements in sediments and that these can be used to study the interactions between ocean acidification, metal bioavailability and accumulation within seagrasses. We affirm warnings by Bary et al., (2011), Vizzini et al., (2013), Lauritano et al., (2015) and Olive et al., (2017) that care is needed when using volcanic CO<sub>2</sub> seeps as analogues for the effects of ocean acidification as there can be areas of H<sub>2</sub>S emissions and toxic levels of metals that may be confounding factors. In some cases, such as Ischia, we found that high levels of trace elements such as copper were not bioavailable to seagrass. In other cases, such as Vulcano, we found that elevated levels of Zn were bioavailable and became concentrated in seagrass roots and rhizomes. Our findings are relevant to those agencies responsible for monitoring the effects of trace metal contamination in the marine environment, since increased CO<sub>2</sub> levels can increase metal availability and toxicity.

---

#### 4.5. References

- Ahmad, F., Azman, S., Savid, M.I.M., Baloo, L., 2015. Tropical seagrass as bioindicator of metal accumulation. *Sain. Malaysia*. 203-210
- Aiuppa, A., Dongarrà, G., Capasso, G., Allard, P., 2000. Trace elements in the thermal ground waters of Vulcano Island (Sicily). *J. of Volcanol. Geotherm. Res.* 98:189- 207
- Ambo Rappe, R., Lajus, D.L., Schreider, M.J., 2007. Translational fluctuating asymmetry and leaf dimension in seagrass, *Zostera capricorni* Aschers in a gradient of heavy metals. *Environ. Bioindic.*, 2: 99-116.
- Ambo-Rappe, R., Lajus, D.L., Schreider, M.J., 2011. Heavy metal impact on growth and leaf asymmetry of seagrass *Halophila ovalis*. *Jour. Of. Envir. Chem. and Ecotox.* 6:145-149
- Apostolaki E.T., Vizzini, S., Hendriks, I.E., Olsen, Y.S., 2014. Seagrass ecosystem response to long-term high CO<sub>2</sub> in a Mediterranean volcanic vent. *Mar. Environ. Res.* 99: 9–15
- Atkinson, C.A., Jolley, D.F., Simpson, S.L., 2007. Effect of overlying water pH, dissolved oxygen, salinity and sediment disturbances on metal release and sequestration from metal contaminated marine sediments. *Chemosphere.* 9: 1428-1437
- Baggini, C., Salomidi, M., Voutsinas, E., Bray, L., et al., 2014. Seasonality affects Macroalgal Community Response to Increases in pCO<sub>2</sub>. *PLOS one.* 9: 1-13
- Barry, J.P., Hall-Spencer, J.M., Tyrrell, T., 2010. In situ perturbation experiments: natural venting site, spatial/temporal gradients in ocean pH, manipulative in situ pCO<sub>2</sub> perturbations. In: Riebesell, U., Fabry, V.J., Hansson, L., Gattuso, J.-P. (Eds.), *Guide to Best Practices for Ocean Acidification Research and Data Reporting*. Publications Office of the European Union, Luxembourg, pp. 123-136.
- Basallote, M.D., De Orte, M.D., Angel Del Valls, T., Riba, I., 2014. Studying the effect of CO<sub>2</sub> induced acidification on sediment toxicity using Acute Amphipod Toxicity Test. *Environ. Sci. and Tech.* 48: 8864-8872.
- Batley, G.E., Apte, S.C., Stauber, J.L., 2004. Speciation and bioavailability of trace metals in water: Progress since 1982. *Australia. J. of Chem.* 57: 903- 919
- Besar, S.N.T., Shazili, N.A.M., Abdullah, S.A., Mamat, A.S., 2008. Experimental and field study on accumulation of heavy metals. *J. of Sustain. And Manag.* 3: 41-73
- Blott, S.J. and Pye, K., 2001. GRADISTAT: a grain size distribution and statistics package for the analysis of unconsolidated sediments. *Earth Sur. Pro. and Landfo.* 26: 1237-1248
- Boatta, F., D'Alessandro, W., Gagliano, A.L., Liotta, et al., 2013. Geochemical survey of Levante bay, Vulcano island (Italy), a natural laboratory for the study of ocean acidification. *Mar. Pollut. Bull.* 73: 485-494
- Bonanno, G., Di Martino, V., 2016. Seagrass *Cymodocea nodosa* as a trace element biomonitor: Bioaccumulation patterns and biomonitoring uses. *J. of Geochem. Explor.* 169: 43-49.

- Bouchon, C., Lemoine, S., Dromard, C., Bouchon-Navarao, C., 2016. Level of contamination by metallic trace elements and organic molecules in the seagrass beds of Guadeloupe Island. *Environ. Sc. Pollut. Res.* 72: 23-61
- Bravo, I., Focaracci, F., Cerfolli, F., Papetti, P., 2016. Relationships between trace elements in *Posidonia oceanica* shoots and in sediment fractions along Latium coasts (northwestern Mediterranean Sea). *Env. Mont. Assess.* 3:157
- Brodie, J., Williamson, C., Smale, D.A., Kamenos, N.A., et al., 2014. The future of the northeast Atlantic benthic flora in a high CO<sub>2</sub> world. *Ecol. Evol.* 13: 2787-2798
- Byrne et al. 1988. The influence of temperature and pH on trace metal speciation in seawater. *Mar. Chem.* 25: 163–181.
- Caeiro, S., Costa, M. H., Ramos, T. B., 2005. Assessing Heavy Metal Contamination in Sado Estuary Sediment: An Index Analysis Approach. *Ecol. Indi.* 5: 151–169
- Caldeira, K., Wickett, M.E. 2003. Oceanography: anthropogenic carbon and ocean pH. *Nature.* 425: 365
- Capaccioni, B., Tassi, F., Vaselli, O., 2001. Organic and inorganic geochemistry of low temperature gas discharges at the Baia di Levante beach, Vulcano Island, Italy. *J. Volcanol. Geoth. Res.* 108: 173–185
- Caramanna, G., Voltattorni, N., Maroto-Valer, M.M., 2011. Is Panarea Island (Italy) a valid and cost-effective natural laboratory for the development of detection and monitoring techniques for submarine CO<sub>2</sub> seepage? *Greenhouse Gases: Sci. Technol.* 1: 200–210
- Catsiki, V. A. and Panayotidis, P., 1993. Copper, chromium and nickel in tissues of the Mediterranean seagrasses *Posidonia oceanica* and *Cymodocea nodosa* from Greek coastal areas. *Chemosphere.* 26:963–978.
- Connell, S.D., Doubleday, Z.A., Hamlyn, S.B., Foster, N.R., et al., 2017. How ocean acidification can benefit calcifiers. *Cur. Bio.* 3:95-96.
- Coles, R. G., McKenzie, L. J., Rasheed, M. A., Mellors, J. E., et al., 2007. Status and Trends of Seagrass Habitats in the Great Barrier Reef World Heritage Area. Report to the Marine and Tropical Sciences Research Facility. Reef and Rainforest Research Centre Limited, Cairns (122 pp.).
- Cravo, A., Foster, P., Almeida, C., Company, R., et al., 2007. Metals in the shell of *Bathymodiolus azoricus* from a hydrothermal vent site on the Mid-Atlantic Ridge. *Environ. Int.* 33: 609- 615.
- Dando, P. R., Stuben, D. & Varnavas, S. P., 1999. Hydrothermalism in the Mediterranean Sea. *Prog. Oceanogr.* 44:333–367
- D'Alessandro, W., Brusca, L. Kyriakopoulos, K., Michas, G. et al., 2008. Methana, the westernmost active volcanic system of the South Aegean Arc (Greece): Insights from fluids geochemistry. *Jour. Of Volcano. And Geoth. Res.* 178: 818-828

- De Orte, M. R., Sarmiento, A. M., Basallote, M. D., Rodríguez- Romero, A., et al., 2014. Effects on the mobility of metals from acidification caused by possible CO<sub>2</sub> leakage from sub-seabed geological formations. *Sci. Total Environ.* 470: 356–363
- Dickson, A.G., Sabine, C.L., Christian, J.R., (Eds), 2007. Guide to best practices for ocean CO<sub>2</sub> measurements. PICES Special Publication, 3: 1-191
- DI Leo, A., Annicchiarico, N., Cardellicchio, L., 2013. Trace metal distribution in *Posidonia oceanica* and sediments from Taranto Gulf (Ionian Sea, Southern Italy). *Med. Mar. Sci.* 14: 204-213
- Doney, S.C., Fabry, V.J., Feely, R.A., Kleypas, J.A., 2009. Ocean acidification: the other CO<sub>2</sub> problem. *Annu. Rev. Mar. Sci.* 1: 169–192
- Dong, Y., Rosenbaum, R.K., Hauschild, M.Z., 2016. Assessment of metal toxicity in marine ecosystems; comparative toxicity potentials for nine cationic metals in coastal water. *Env. Sci. and Tech.* 50: 269-278.
- Eggleton, J., Thomas, K.V., 2004. A review of factors affecting the release and bioavailability of contaminants during sediment disturbance events. *Environ. Int.* 30: 973–980
- Enochs, I.C., Manzello, D.P., Donham, E.M., Kolodziej, G., et al. 2015. Shift from coral to macroalgae dominance on a volcanically acidified reef. *Nat. Cli. Ch.* 5:1083-88.
- Fabricius, K.E., De'ath, G., Noonan, S., Uthicke, S., 2014. Ecological effects of ocean acidification and habitat complexity on reef-associated macroinvertebrate communities. *Proc. R. Soc. B Biol. Sci.* 281 (1775)
- Fan, W., Wang, W-X., Chen, J., Li, X., et al., 2002. Cu, Ni and Pb speciation in surface sediments from a contaminated bay of northern China. *Mar. Pollut. Bull.* 44:816–32.
- Fergusson, J.E., 1990. The heavy elements' chemistry, Environmental impacts and Health effects. Pergamon Press, Oxford, UK.
- Fourqurean, J.W., Duarte, C.M., Kennedy, H., Marbà, N., et al., 2012. Seagrass ecosystems as a globally significant carbon stock. *Nat. Geosci.* 5: 505–509
- Gabianelli, G., Gillot, P.Y., Lanzafame, G., Romagnoli, C. et al., 1990. Tectonic and volcanic evolution of Panarea (Aeolian Islands, Italy). *Mar. Geo.* 92: 313–326
- Goffredo, S., Prada, F., Caroselli, E., Capaccioni, B., et al., 2017. Biomineralization control related to population density under ocean acidification. *Nat. Clim. Cha.* 4:593-597
- Guilizzoni, P., 1991. The role of heavy metals and toxic materials in the physiological ecology of submersed macrophyte. *Aqua. Bot.* 41:87–109.
- Hall-Spencer, J.M., Rodolfo-Metalpa, R., Martin, S., Ransome, E. et al., 2008. Volcanic carbon dioxide seeps show ecosystem effects of ocean acidification. *Nature* 454:96-99
- IPCC. 2014. Fifth Assessment Report on Climate change: Impacts, Adaptation and Vulnerability

- Inguaggiato, S., Mazot, A., Diliberto, I.S., Inguaggiato, C., Madonia, P., Rouwet, D. et al., 2012. Total CO<sub>2</sub> output from Vulcano Island (Aeolian Islands, Italy). *Geochem. Geophys. Geosyst.* 13: Q02012
- Ivanina, A.V., Sokolova, I.M., 2015. Interactive effects of metal pollution and ocean acidification on physiology of marine organisms. *Cur. Zool.* 64: 653-668.
- Kadar, E., Costa, V., Segonzac, M., 2007. Trophic influences of metal accumulation in natural pollution laboratories at deep-sea hydrothermal seeps of the Mid- Atlantic Ridge. *Sci. Tot. Environ.* 373: 464-472
- Kadar, E., Fisher, A., Stolpe, B., Harrison, R.M. et al., 2012. Metallic nanoparticle enrichment at low temperature, shallow CO<sub>2</sub> seeps in Southern Italy. *Mar. Chem.* 140: 24-32
- Klumpp, D.W., Howard, R.K., Pollard, D.A., 1989. Trophodynamics and nutritional ecology of seagrass communities. Elsevier, Amsterdam, pp-394-437
- Koch, M., Bowes, G., Ross, C., Zhang, X., 2013. Climate change and ocean acidification effects on seagrasses and marine macroalgae. *Glob. Change Biol.* 19:103-132.
- Kristensen, E., 2000. Organic matter diagenesis at the oxic/anoxic interface in coastal marine sediments, with emphasis on the role of burrowing animals. *Hydrobiologia.* 426:1-24.
- Kroeker, K.J., Micheli, F., Gambi, M.C., Martz, T.R., 2011. Divergent ecosystem responses within a benthic marine community to ocean acidification. *PNAS* 108 :14515-14520.
- Kroeker, K. J., Kordas, R.L., Crim, R., Hendriks, I.E. et al., 2013. Impacts of ocean acidification on marine organisms: quantifying sensitivities and interaction with warming. *Glob. Change Biol.* 19:1884-1896
- Kvesitadze, E., Sadunishvili, T. & Kvesitadze, G., 2009. Mechanisms of organic contaminants uptake and degradation in plants. *World Acad. Sci. Eng. Tech.* 55: 458-468
- Lacoue-Labarthe, T., Martin, S., Oberhansli, F., Teyssie, J.L. et al., 2009. Effects of increased pCO<sub>2</sub> and temperature on trace element (Ag, Cd and Zn) bioaccumulation in the eggs of the common cuttlefish, *Sepia officinalis*. *Biogeosciences*, 6: 2561-2573.
- Lacoue-Labarthe, T., Martin, S., Oberhansli, F., Teyssie, J.L. et al., 2012. Temperature and pCO<sub>2</sub> effect on the bioaccumulation of radionuclides and trace elements in the eggs of the common cuttlefish, *Sepia officinalis*. *J. of Exp. Mar. Bio. and Eco.* 413:45-49
- Lafabrie, C., Pergent, G., Kantin, R., Pergent-Martini, C. et al., 2007. Trace metals assessment in water, sediment, mussel and seagrass species - Validation of the use of *Posidonia oceanica* as a metal bio monitor. *Chemosphere* 68: 2033-2039.
- Lau, S., Mohamed, M., Tan Chi Yen, A., Suut, S., 1998. Accumulation of heavy metals in freshwater molluscs. *Sci. Total Environ.* 214:113-121.
- Lauritano, C., Ruocco, M., Dattolo, E., Buia, M.C., et al., 2015. Response of key stress-related genes of the seagrass *Posidonia oceanica* in the vicinity of submarine volcanic vents. *Biogeo. Discus.* 12: 4947-4971

- Lemasson, A.J., Fletcher, S., Hall-Spencer, J.M., Knights, A.M., 2017. Linking the biological impacts of ocean acidification on oysters to changes in ecosystem services: A review. *J. of Expt. Mar. Bio. and Ecol.* 492:49-62
- Lewis, M.A., Devereux, R., 2009. Non-nutrient anthropogenic chemicals in seagrass ecosystems: fate and effects. *Environ. Toxicol. Chem.* 28: 644- 661
- Lewis, C. Ellis, R.P., Mangan, S., Campbell, A.L., 2014. Ocean acidification increases copper toxicity to the early life history stages of the Polychaete *Arenicola marina* in artificial seawater. *Envir. Sci. & Tech.* 48:9745-9753
- Lewis, C., Ellis, R.P., Vernon, E., Elliot, K., et al., 2016. Ocean acidification increases copper toxicity differentially in two key marine invertebrates with distinct acid-base responses. *Nature.* 6: 1-10.
- Llagostera, I., Pérez, M. & Romero, J. 2011. Trace metal content in the seagrass *Cymodocea nodosa*: Differential accumulation in plant organs. *Aqua. Bot.* 95: 124-128
- Long, E. R., and D. D. MacDonald., 1998. Recommended uses of empirically derived, sediment quality guidelines for marine and estuarine ecosystems. *Hum. and Ecol. Risk Assess.* 4; 1019-1039.
- MacDonald, D. D., Carr, R. S., Calder, F. D., Long, E. R, et al., 1996. Development and evaluation of sediment quality guidelines for Florida coastal waters. *Ecotoxicology.* 5:253-278
- MacDonald, D.D., Lindskoog, R.A., Smorong, D.E., Greening, H., et al., 2000. Development of an Ecosystem-based Framework for Assessing and Managing Sediment Quality Conditions in Tampa Bay, Florida. Tampa Bay Estuary Pro- gram, Florida, USA.
- Malea, P., Haritonidis, S., Kevrekidis, T., 1994. Seasonal and local variations of metal concentrations in the seagrass *Posidonia oceanica* (L.) Delile in the Antikyra Gulf, Greece. *Sci. of the Total Env.* 153: 225- 235.
- Malea, O. and S. Haritonidis., 1999. *Cymodocea nodosa* (Ucria) Aschers. as a Bioindicator of Metals in Thermaikos Gulf, Greece, during Monthly Samplings. *Bot. Mar.* 42: 419-430
- Milazzo, M., Rodolfo-Metalpa, R., Chan, V.B.S., Fine, M. et al., 2014. Ocean acidification impairs vermetid reef recruitment. *Scienti. Rep.* 4 (4189).
- Milazzo, M., Cattano, C., Alonzo, S.H., Foggo, A. et al., 2016. Ocean acidification affects fish spawning but not paternity at CO<sub>2</sub> seeps. *Royal Society.* 283 (1835).
- Millero, F.J., Woosley, R., DiTrollo, B., Waters, J., 2009. Effect of ocean acidification on the speciation of metals in seawater. *Oceanography.* 22:72–85.
- Monia-Renzi, Teresa, R., C, G., Guido, P., Italianod, S. E et al., 2011. Temporal trends and matrix-dependent behaviors of trace elements closed to a geothermal hot-spot source (Aeolian Archipelago, Italy). *Proced. Earth and Planetary Sci.* 4:10 – 28
- Olive, I., Silva, J., Lauritano, C., Costa, M.M., et al., 2017. Short term responses of seagrasses exposed to CO<sub>2</sub> in volcanic vents. *Scientific Rep.*7:42278

- Orth, R.J., Carruthers, T.J.B., Dennison, W.C., Duarte, C.M. et al., 2006. A Global Crisis for Seagrass Ecosystems. *BioScience* 56: 987–996.
- Pascal, P.Y., Fleeger, J.W., Galvez, F., Carman, K.R., 2010. The toxicological interaction between ocean acidity and metals in coastal meiobenthic copepods. *Mar. Pollut. Bull.* 60: 2201–2208
- Pekey, H., Karakas, D., Ayberk, S., Tolun, L., et al., 2004. Ecological risk assessment using trace elements from surface sediments of Izmit Bay (North-eastern Marmara Sea) Turkey. *Mar. Pollut. Bull.* 48: 946-953
- Prange, J, A, Dennison, W.C., 2000. Physiological responses of five seagrass species to trace metals. *Mar. Pollut. Bull.*, 41: 327-336
- Prokop, Z., Vangeheluwe, M.L., Van Sprang, P.A., Janssen, C.R., et al., 2003. Mobility and toxicity in sandy sediments deposited on land. *Ecotox. Environ. Saf.* 34: 65-73
- Quinn, G. P. and Keough, M. J.: *Experimental design and data analysis for biologists*, Cambridge University Press, Cambridge, 2002.
- Ralph, P.J., David, T., Kenneth, M., Stephanie, S. et al., 2006. Human impacts on seagrasses: eutrophication, sedimentation, and contamination. In: Larkum, A.W.D., Orth, R.J., Duarte, C.M. (Eds.), *Seagrasses: Biology, Ecology and Conservation*. Springer, Dordrecht, The Netherlands, pp. 567-593.
- Riba, I., Delvalls, T., Á.; Forja, J. M., Gómez-Parra, A., 2004. The influence of pH and salinity on the toxicity of heavy metals in sediment to the estuarine clam *Ruditapes philippinarum*. *Environ. Toxicol. Chem.* 23: 1100–1107
- Richards, R., Chaloupka, M., Sano, M., Tomlinson, R., 2011. Modelling the effects of “coastal” acidification on copper speciation. *Ecol. Model.* 222: 3559–3567
- Richir, J., Gobert, S., 2013. The effect of size, weight, body compartment, sex and reproductive status on the bioaccumulation of 19 trace elements in rope-grown *Mytilus galloprovincialis*. *Ecol. Ind.* 36: 33–47
- Richir, J., Gobert, S., 2014. A reassessment of the use of *Posidonia oceanica* and *Mytilus galloprovincialis* to bio monitor the coastal pollution of trace elements: New tools and tips. *Mar. Pollut. Bull.* 89: 390-406
- Richir, J., Gobert, S., 2016. Trace elements in Marine Environment; Occurrence, Threats and Monitoring with Special Focus on the Coastal Mediterranean. *Env. & Anal. Toxicol.* 6:1-19
- Roberts, D. A., Birchenough, S.R., Lewis, C., Sanders, M.B. et al., 2013. Ocean acidification increases the toxicity of contaminated sediments. *Glo. Change Bio.* 19: 340-351
- Ruocco, M., Musacchia, F., Olive, I., Costa, M.M., et al., 2017. Genomewide transcriptional reprogramming in the seagrass *Cymodocea nodosa* under experimental ocean acidification. *Mol. Ecol.* 1-19

- Russell, B.D., Connell, S.D., Uthicke, S., Muehllehner, N. et al., 2013. Future seagrass beds: can increased productivity lead to increased carbon storage? *Mar. Pollut. Bull.* 73:463-469
- Sanchiz, C., Garcı́a-Carrascosa, A.M., Pastor, A., 2001. Relationships between sediment physico-chemical characteristics and heavy metal bioaccumulation in Mediterranean soft-bottom macrophyte. *Aquat. Bot.* 69: 63–73
- Sedwick PN, Stuben D., 1996. Chemistry of shallow submarine warm springs in an arc-volcanic setting: Volcano Island, Aeolian Archipelago, Italy. *Mar. Chem.* 53: 147-161.
- Shi, W., Zhao, X., Han, Y., Che, Z., et al., 2016. Ocean acidification increases cadmium accumulation in marine bivalves: a potential threat to seafood safety. *Sci. Rep.* 6:20197
- Simpson, S.L., Rochford, L., Birch, G.F., 2002. Geochemical influences on metal partitioning in contaminated estuarine sediments. *Mar. Freshw. Res.* 53: 9–17
- Simpson, S.L., Batley, G.E., 2003. Disturbances to metal partitioning during toxicity testing iron(II)-rich estuarine pore waters and whole- sediments. *Environ. Toxicol. Chem.* 22: 424–432
- Simpson, S.L., Angel, B.M., Jolley, D.F., 2004. Metal equilibration in laboratory-contaminated (spiked) sediments used for the development of whole-sediment toxicity tests. *Chemosphere* 54: 597–609
- Soliman, F.N., Nasr, M.S., Okbah, M.A., 2015. Potential ecological risk of heavy metals in sediments from the Mediterranean coast, Egypt. *J. of Env. Heal. Sci. and Eng.* 10: 1-12
- Sternbeck J, Östlund P., 2001. Metals in sediments from the Stockholm region: Geographical pollution patterns and time trends. *Water Air Soil Pollut. Focus.* 1:151-165
- Stoffers, P., Hannington, M., Wright, I., Herzig, P. et al., 1999. Elemental mercury at submarine hydrothermal seeps in the Bay of Plenty, Taupo volcanic zone, New Zealand. *Geol. Soci. of America.* 10: 931-934
- Stumm, W., and Morgan, J.J., 1995. *Aquatic chemistry: chemical equilibria and rate in natural waters*, 3rd ed.
- Sunday, J.M., Fabricius, K.E., Kroeker, K.J., Anderson, K.M. et al., 2016. Ocean acidification can mediate biodiversity shifts by changing biogenic habitat. *Nat. Clim. Change.* 7: 81-85
- Szefer, P., Ali, A.A., Ba-Haroon, A.A., Rajeh, A.A., et al., 1999. Distribution and relationships of selected trace metals in molluscs and associated sediments from the Gulf of Aden, Yemen. *Environ. Pollut.* 106: 299–314
- Tanhua, T., Hainbucher, D., Schroeder, K., Cardin, V., et al., 2013. The Mediterranean Sea system: A review and an introduction to the special issue. *Ocean Sci.*5:789-803.
- Tchounwou, P.B., Yedjou, C.G., Patlolla, A.K., Sutton, D.J., 2014. Heavy metals toxicity and the Environment. *Mol., Clinical and Env. Toxicol.* 101:133-164
- Tedesco, D., 1996. Chemical and isotopic investigation of fumarolic gases from Ischia Island (Southern Italy): evidence of magmatic and crustal contribution. *J. Vulcanol. Geother. Res.* 74:233–242

Thom, R. M., 2001. Carbon Sinks in Nearshore Marine Vegetated Ecosystems. Pacific Northwest Laboratory, Marine Science Laboratory.

Underwood, A. J. 1997. Experiments in Ecology: Their Logical Design and Interpretation Using Analysis of Variance, Cambridge University Press, Cambridge.

Voltattorni, N., Sciarra, A., Caramanna, G., Cinti, D. et al., 2009. Gas geochemistry of natural analogues for the studies of geological CO<sub>2</sub> sequestration. *Applied Geochem.* 24:1339–1346

Vizzini, S, Tomasello, A., Maida, G. Di, Pirrotta, M. et al., 2010. Effect of explosive shallow hydrothermal seeps on <sup>13</sup>C and growth performance in the seagrass *Posidonia oceanica*. *J. of Ecol.* 98:1284–1291

Vizzini, S, Di Leonardo, R., Costa, V., Tramati, C.D. et al., 2013. Trace element bias in the use of CO<sub>2</sub> seeps as analogues for low pH environments: Implications for contamination level in acidified oceans. *Estuarine, coast. and Shelf Sci.* 134:19-30

Weis, J. S., Weis, P., 2004. Metal uptake, transport and release by wetland plants: implications for phytoremediation and restoration. *Environ. Int.*30:685–700

Zeng, X., Chen, X., 2015. The positive relation between ocean acidification and pollution. *Mar. Poll. Bull.* 91:14-21.

Zoumis, T., Schmidt, A., Grigorova, L., Calmano, W., 2001. Contaminants in sediments: remobilisation and demobilisation. *Sci. Total Environ.* 266: 195–202

## General Discussion





---

**General Discussion**

**5.1. Main findings and implications for seagrass ecosystem**

Seagrasses are one of the dominant coastal ecosystems characterized by high primary production rates, nutrient concentrations and biodiversity (Thom, 2001). These habitats are important for the ocean's biogeochemical cycles, and they play a crucial role in providing ecosystem services (Gattuso et al., 1998; Erwin et al., 2009). Information on response of seagrass community and individual plants to high CO<sub>2</sub> scenarios are important to predict the state of seagrass community production and survival in future conditions and therefore needed to frame possible adaptive measures to keep in flow the ecosystem services they provide. This thesis contributed to assess seagrass responses to future high CO<sub>2</sub> levels and the main findings are as follows

- Short term CO<sub>2</sub> enrichment incubations on intertidal *Z. noltii* community indicated high CO<sub>2</sub> (750 ppm) levels projected for the end of this century will increase the net community production and decrease the community respiration during emersion periods. Light requirements of plants were reduced under CO<sub>2</sub> enrichment conditions (Chapter 2)
- At Mediterranean CO<sub>2</sub> seeps of Greece and Italy, *C. nodosa* population showed positive response to long term exposure of elevated CO<sub>2</sub> through higher growth, density, biomass and morphometry. Higher number of younger plants (1 and 2 years older) were found at the CO<sub>2</sub> seeps, whereas the number of older plants (more than 3 years) were lower near the CO<sub>2</sub> seeps at all three locations (Chapter 3).
- Fe and trace element (Co, Cu, Hg, Ni, Mn, Pb and Zn) concentrations were higher under high CO<sub>2</sub> and low pH conditions in the sediments and in *C. nodosa* and *P. oceanica* roots, rhizomes and leaves. Bioavailability of trace elements increased with increase in CO<sub>2</sub> concentration and low pH. Higher concentration in sediments resulted in higher accumulation of trace elements in the seagrass compartments. Trace element concentrations in sediments and plant compartments were found at levels that can be toxic to seagrass at CO<sub>2</sub> seeps of Greece and Italy. CO<sub>2</sub> seeps of Italy (Vulcano, Ischia and Panarea) were observed with higher concentration of elements than CO<sub>2</sub> seeps of Greece (Milos, Paleochori and Methana) (Chapter 4).

Thus, seagrass ecosystem and individual plants can get benefitted from the increased CO<sub>2</sub> levels in the atmosphere, whereas increase in bioavailability of elements and possible toxic

effects due to ocean acidification and low pH can hinder this positive influence of high CO<sub>2</sub> in future scenarios.

Our results indicate positive response of seagrass to elevated levels (future projections of 750 ppm) of CO<sub>2</sub>. This positive response of seagrass to higher CO<sub>2</sub> concentrations was noticed on both the short-term experiments on *Z. noltii* community and at the CO<sub>2</sub> seeps of Greece and Italy, where the seagrass (*C. nodosa*) have been exposed to long term elevated CO<sub>2</sub> levels (750 ppm and higher). Positive response of *Z. noltii* through increased community production in (Chapter 2) indicates that the plants are carbon limited in the current carbon dioxide concentration in atmosphere and gradually as CO<sub>2</sub> concentration increases the plants will adapt and increase their carbon utilization capacity, which can result in higher density, biomass and higher production rates as observed for *C. nodosa* at CO<sub>2</sub> seeps exposed to long term elevated CO<sub>2</sub> levels (Chapter 3). Similar response of other seagrass (*Z. marina* and *T. hemprichii*) species have been observed with a positive response to high CO<sub>2</sub> levels (Zimmerman et al., 1997; Jiang et al., 2010).

The positive response of seagrass to increased CO<sub>2</sub> and to overcome the current carbon limitation in future oceans, a lot will depend on how seagrass to utilize the dissolved inorganic carbon (DIC) available to them in high CO<sub>2</sub> oceans. The amount of DIC (1% CO<sub>2</sub> (aq), 90% HCO<sub>3</sub><sup>-</sup> and 9% CO<sub>3</sub><sup>2-</sup>) in the current oceans is going to get higher (Campbell et al., 2013) with an overall increase of CO<sub>2</sub> (>250%) compared to bicarbonate (24%, HCO<sub>3</sub><sup>-</sup>) under the projected pH decrease by 2100 (Koch et al., 2013)

Seagrasses will require to utilize the abundant HCO<sub>3</sub><sup>-</sup> ions to increase their photosynthetic production efficiency by extracellular dehydration of into CO<sub>2(aq)</sub> via membrane bound carbonic anhydrase enzymes (Beer and Rehnberg, 1997; Bjork et al., 1997) or by proton extrusion into an unstirred boundary layer adjacent to the leaf surface facilitating carbonic anhydrase activity (Uku et al., 2005). Extracellular carbonic anhydrase dehydrates HCO<sub>3</sub><sup>-</sup> to CO<sub>2</sub>, allowing CO<sub>2</sub> to diffuse into the cell. In general, active utilization of H<sup>+</sup> ions at the leaf boundary layer for HCO<sub>3</sub><sup>-</sup> utilization is energetically costly compared to passive CO<sub>2</sub> uptake and could be limited by low light levels (Burnell et al., 2014), but the seagrass ecosystems in shallow habitats with higher light penetration can favour the use of HCO<sub>3</sub><sup>-</sup> as substrate.

Positive response of *Z. noltii* production to CO<sub>2</sub> enrichment (Chapter 2) and increase in density and biomass of *C. nodosa* in the CO<sub>2</sub> seeps of Greece and Italy (Chapter 3) suggests that seagrasses here have adapted to the utilization of DIC (CO<sub>2</sub> and HCO<sub>3</sub><sup>-</sup>) as *Z. noltii* is

intertidal receiving high light intensities during emersion and the *C. nodosa* meadows at the CO<sub>2</sub> seeps of Greece were in shallow waters receiving higher light. Both seagrasses can prefer the use of CO<sub>2</sub> when available in higher concentration than HCO<sub>3</sub><sup>-</sup>, as utilizing CO<sub>2</sub> requires less energy than HCO<sub>3</sub><sup>-</sup>. Though the degree of adaptation to utilize HCO<sub>3</sub><sup>-</sup> is species specific, the availability of light also plays a major role in seagrass utilizing HCO<sub>3</sub><sup>-</sup>, observed in seagrass species, such as *C. serrulata* (Uku et al., 2005; Ow et al., 2016), *C. rotundata* (Schwarz et al., 2000) and *Z. marina* (Invers et al., 2001) which showed increase production due to increased DIC and light, whereas *Halophila uninervis* productivity was not affected by increased DIC but only by light (Ow et al., 2016). This implies in future oceans, increased DIC concentrations and increased production (Invers et al., 2002; Hall-Spencer et al., 2008).

The adaptation to use DIC will also depend on the available concentration of DIC in ecosystem and habitat of the seagrass. The seagrasses which will remain submerged all the time will have better capacity to adapt and utilize HCO<sub>3</sub><sup>-</sup>, than intertidal species which are both submerged and air exposed (Campbell et al., 2013, Ow et al., 2016). This differences can favour one seagrass in a mixed meadow to improve and increase its carbon balance, thus increasing its competitive ability to utilize high CO<sub>2</sub>, while other species can remain less beneficial. Similar, results have been reported for *Thalassia testudinum* and *Halophila wrightii* (Campbell et al., 2013). Increase in DIC concentration as observed at the CO<sub>2</sub> seeps of Greece and Italy (Chapter 3&4) will favour the increase in photosynthetic capacity of seagrass by providing more carbon for fixation and simultaneously lowering photorespiration rates (Long et al., 2004), whereas increase in light intensities will also increase the maximum photosynthetic capacity by boosting production and reducing intermediates for the carbon reduction cycle (Hall et al., 1994). As both factors have independent mode of actions, their effect on seagrass productivity will be synergistic and beneficial to the seagrasses.

DIC enrichment can also reduce the light saturation point for seagrasses and increase the production capacity of seagrasses, as observed for *Z. noltii* in our research (Chapter 1), similar results were observed for *C. serrulata* (Ow et al., 2016) and *T. hemprichii* (Jiang et al., 2010) and *Z. marina* (Zimmerman et al., 1997). This suggests seagrass can increase their light saturated photosynthetic capacity under increased CO<sub>2</sub> concentration to reduce the daily photoperiod required for positive carbon balance (Zimmerman et al., 1997, Ow et al., 2016) and increase their migration into greater depths.

Under future ocean acidification scenarios, seagrasses can benefit from the increased DIC concentration and CO<sub>2</sub> levels (Hall-Spencer et al., 2008; Fabricius et al., 2011). While short term and long term studies on seagrass have documented varying degrees of DIC limitation in seagrass, physiological processes that could regulate responses to increased DIC are less explored for seagrass. Other than DIC limitation, other factors that will regulate seagrass growth and production will be nutrient availability (Stitt and Krapp, 1999) and water temperature (Touchette and Burkholder, 2000), which are also going to increase in the future oceans due to anthropogenic pollution and climate change. Saying that, seagrass leaves will also have to compete with the leaf epiphytes (mostly algae) for the same resources of increased DIC concentrations. This competition with epiphytic algae will determine the composition, biomass and turnover rates of seagrass leaves (Campbell et al., 2013). Finally, light availability, due to anthropogenic pollution, sedimentation and epiphytic algal growth can negate the positive effects of high CO<sub>2</sub> on seagrass productivity in future oceans (Burnell et al., 2014).

Increase in CO<sub>2</sub> concentration will favour the increase in seagrass production, but productions of seagrass ecosystem will be under considerable stress from various factors such as metal contamination. Metal levels is going to increase in the world oceans due to anthropogenic impacts from coastal developments and agricultural runoff (Ivanina et al., 2015; Zeng et al., 2015). Ocean acidification and low pH will alter the speciation and availability of elements in seawater and sediments resulting in potential toxicity to the marine biota (Millero et al., 2009; Roberts et al., 2013). Higher availability of element levels in the sediments can result in possible contamination of marine sediments and higher accumulation in the seagrass compartments as observed in our results (Chapter 4).

Generally marine sediments have significant higher magnitude of metal concentrations than in water column making the sediments important reservoir of these metals (Clark, 2002). These metal levels are going to increase from anthropogenic pollution and a combination of ocean acidification and low pH in the water column (Roberts et al., 2013). Presence of metal levels in higher concentration may inhibit the primary production in marine ecosystems and decrease the efficiency of marine carbon sink (Zeng et al., 2015). Seagrasses have metal binding proteins that can help them regulate metal toxicity on their metabolic activity and physiology under normal conditions, but under ocean acidification and low pH conditions, it have been observed in *P. oceanica* (Lauritano et al., 2015) and *C. nodosa* (Olive et al., 2017) downregulation of these metal binding proteins occurs in seagrass. This downregulation of

metal binding proteins suggests, seagrasses may not be able to tolerate the metal toxicity levels that is going to occur by the end of this century. This also indicates that metal levels we found in our results at Ischia and Panarea CO<sub>2</sub> seeps in *P. oceanica* compartments and at Vulcano in *C. nodosa* compartments (Chapter 4) are already at higher levels affecting their metabolic activity and physiology. Low longevity of *C. nodosa* at CO<sub>2</sub> seeps off Greece and Italy than reference sites (Chapter 3) may be an effect of metal toxicity too due to similar levels of pH and metal concentration in sediments at CO<sub>2</sub> seeps off Italy and Greece (Chapters 3&4).

However, accumulation patterns of metals in seagrass will determine their concentration in the roots, rhizomes and leaves of seagrass. This suggests presence of metals in higher concentration in water or sediments may not necessarily have toxic effects on seagrass as they have an exclusive storage capacity of metals in rhizomes and exclusion mechanism by leaves.

Similarly, environmental factors other than carbon dioxide will determine favourable conditions for seagrass ecosystem. These factors include the natural biophysical parameters that regulate the physiological activity and morphology of seagrasses (such as temperature, salinity, waves, currents, depth, substrate, day length, light, nutrients, water currents, wave action, epiphytes and diseases), the availability of seeds and vegetative fragments and the anthropogenic inputs that impact plant resources (such as excess nutrients and sediment loading). Combinations of these parameters will permit, encourage or prevent seagrass meadows thriving (Cullen-Unsworth et al., 2014). Climate change is a significant long-term threat to seagrass. Managing seagrasses for future resilience to climate change is about understanding current stressors and how they may change and about knowledge of temperature and ocean chemistry including developing greater knowledge of distribution limits, understanding ecosystem recovery and defining clear physical thresholds (Cullen-Unsworth et al., 2014).

### **5.2. Summary and Main Conclusion**

Global changes in terms of elevated CO<sub>2</sub> levels and low pH have the potential to change the seawater carbonate chemistry and influence the marine ecosystem (Hall-Spencer et al., 2008; Porzio et al., 2011). This thesis contributes to the growing knowledge of effects of high CO<sub>2</sub> concentrations on seagrass ecosystem in both short term and long term enrichment. It is clear from the findings that will increase in CO<sub>2</sub> concentrations will have a positive impact on the seagrass productivity (Alexandre et al. 2012; Russell et al. 2013; Apostolaki et al., 2014) in

both short and long term and will also increase seagrass community productivity and decrease current carbon limitation of seagrass (Zimmerman et al., 1997; Invers et al., 2001). This increase in production will also be affected by various physical factors (light, temperature) and the surrounding environment where the coastal seagrass population are found (Khan et al., 2016; Takahashi et al., 2016). This thesis observes that response of seagrass to high CO<sub>2</sub> concentrations will be species-specific and won't be straight forward and impact of trace metal contamination can over shadow the positive effect of CO<sub>2</sub> enrichment on the seagrass population and community production (Vizzini et al., 2013; Apostolaki et al., 2014).

### **Main Conclusions:**

1. Short term CO<sub>2</sub> enrichment on *Z. noltii* community and sediment community indicated increased community production under CO<sub>2</sub> enrichment than natural conditions.
2. Effect of light and seasons had a clear impact on the net community production of *Z. noltii* and sediment community
3. Short term incubation methods can be used as a reliable to understand intertidal community production using CO<sub>2</sub> enrichment.
4. *C. nodosa* growth, density, biomass, morphometry showed an increased fitness under long term elevated CO<sub>2</sub> concentrations at high CO<sub>2</sub> seeps than control sites of Greece and Italy
5. *C. nodosa* population showed increased recruitment rate and production but longevity of plants decreased under elevated CO<sub>2</sub> concentrations.
6. Reconstruction techniques provide a successful tool to interpret and predict seagrass population dynamics under future high CO<sub>2</sub> scenarios.
7. Trace elements concentration increased under low pH conditions with continuous input from the CO<sub>2</sub> seeps into the sediments
8. Higher concentration in sediments was reflected in the trace element concentration in the seagrass *Cymodocea nodosa* and *Posidonia oceanica* roots, rhizomes and leaves with higher concentrations than control sites.
9. Trace element concentrations in the sediments were higher enough to have moderate or adverse biological impacts on the seagrass ecosystem
10. All the six Mediterranean CO<sub>2</sub> seeps of Greece and Italy was found with higher level of metal contamination than the control sites

11. Seagrass can overcome their current carbon limitation and increase their production and carbon metabolism under future high CO<sub>2</sub> projections and can act as a source of global carbon sink.

### 5.3. References

Alexandre, A., Silva, J., Buapet, P., Bjork, M., Santos, R., 2012. Effects of CO<sub>2</sub> enrichment on photosynthesis, growth and nitrogen metabolism of the seagrass *Zostera noltii*. *Ecol. and Evol.* 2:2626-2635

Apostolaki E.T., Vizzini, S., Hendriks, I.E., Olsen, Y.S., 2014. Seagrass ecosystem response to long-term high CO<sub>2</sub> in a Mediterranean volcanic vent. *Mar. Env. Res.* 99: 9–15

Beer, S and Rehnberg, J., 1997. The acquisition of inorganic carbon by the seagrass *Zostera marina*. *Aquat. Bot.* 56; 277-283.

Bjork, M., Weil, A., Semesi, S., Beer, S., 1997. Photosynthetic utilisation of inorganic carbon by seagrass from Zanzibar, East Africa. *Mar. Biol.* 129; 363-366.

Burnell, O.W., Connell, S.D., Irving, A.D., Watling, J.R., et al., 2014. Contemporary reliance on bicarbonate acquisition predicts increased growth of seagrass *Amphibolis antarctica* in a high-CO<sub>2</sub> world. *Conserv. Physiol.* 2: 1-11

Campbell, J.E., Fourqurean, J.W., 2013. Mechanisms of bicarbonate use influence the photosynthetic carbon dioxide sensitivity of tropical seagrasses. *Lim. and Oceano.*3: 839-848.

Campbell, J.E., Fourqurean, J.W. 2013. Effects of in situ CO<sub>2</sub> enrichment on the structural and chemical characteristics of the seagrass *Thalassia testudinum*. *Mar. Biol.* 160: 1465-1475

Clark R, 2002. *Marine Pollution*. Oxford: Clarendon Press

Cullen-Unsworth, L. C., Nordlund, L. M., Paddock, J., Baker, S., et al., 2014. Seagrass meadows globally as a coupled social–ecological system: Implications for human wellbeing, *Mar. Pollut. Bull.* 83:387–397.

Erwin, K. L., 2009. *Wetlands and Global Climate Change: The Role of Wetland Restoration in a Changing World.* *Wetlands Ecol. and Manage.* 17: 71-84.

Fabricius, K.E., Langdon, C., Uthicke, S., Humphrey, C., et al., 2011. Losers and winners in coral reefs acclimatized to elevated carbon dioxide concentrations. *Nat. Clim. Change.*1: 165-9

Gattuso, J.-P., Frankignoulle, M., Bourge, I., Romaine, S., et al., 1998. Effect of calcium carbonate saturation of seawater on coral calcification. *Glob. Planet. Change.* 18: 37-46

Hall-Spencer, J.M., Rodolfo-Metalpa, R., Martin, S., Ransome, E., et al., 2008. Volcanic carbon dioxide vents show ecosystem effects of ocean acidification. *Nature.* 454: 96–99.

- Hall, D.O., Rao, K.K., 1994. Photosynthesis, 6<sup>th</sup> ed. Great Britain: Cambridge University Press. 198 p
- Invers, O., Zimmerman, R.C., Alberte, R.S., Perez, M., et al., 2001. Inorganic carbon sources for seagrass photosynthesis: an experimental evaluation of bicarbonate use in species inhabiting temperate waters. *J. of Exp. Mar. Bio. Ecol.* 265: 203-217
- Invers, O., Tomas, F., Perez, M., Romero, J., 2002. Potential effect of increased global CO<sub>2</sub> availability on the depth distribution of the seagrass *Posidonia oceanica* (L.) Delile: A tentative assessment using a carbon balance model. *Bull. Mar. Sci.* 71: 1191–1198.
- Ivanina, A.V., Sokolova, I.M., 2015. Interactive effects of metal pollution and ocean acidification on physiology of marine organisms. *Cur. Zoo.* 64: 653-668.
- Jiang, Z., Huang, X., Zhang, J., 2010. Effects of CO<sub>2</sub> enrichment on photosynthesis, growth, and biochemical composition of seagrass *Thalassia hemprichii* (Ehrenb.) Aschers. *J. Integr. Plant. Biol.* 52: 904-913.
- Khan, F., Alexandre, A., Ullah, H., Santos, R., 2016. Effects of elevated CO<sub>2</sub> and nutrients on the community metabolism of a *Cymodocea nodosa* bed. *Turkish J. of Bot.* 40; 250-257
- Koch, M., Bowes, G., Ross, C., Zhang, X., 2013. Climate change and ocean acidification effects on seagrasses and marine macroalgae. *Glob. Change Biol.* 18: 2792-2803.
- Lauritano, C., Ruocco, M., Dattolo, E., Buia, M.C., et al., 2015. Response of key stress-related genes of the seagrass *Posidonia oceanica* in the vicinity of submarine volcanic vents. *Biogeo. Discuss.* 12: 4947-4971
- Long, S.P., Ainsworth, E.A., Rogers, A., Ort, D.R., 2004. Rising Atmospheric Carbon Dioxide: Plants FACE the future. *Annu. Rev. Plant. Biol.* 55: 591-628
- Millero, F.J., Woosley, R., DiTrollo, B., Waters, J., 2009. Effect of ocean acidification on the speciation of metals in seawater. *Oceanography.* 22:72–85.
- Olive, I., Silva, J., Lauritano, C., Costa, M.M., et al., 2017. Short term responses of seagrasses exposed to CO<sub>2</sub> in volcanic vents. *Scientific Rep.* 7:42278
- Ow, Y.X., Uthicke, S., Collier, C.J., 2016. Light levels affect carbon utilisation in tropical seagrass under ocean acidification. *PLoS ONE.* 3: 1-18
- Porzio, L., M. C. Buia, and J. M. Hall-Spencer. 2011. Effects of ocean acidification on macroalgal communities. *J. Exp. Mar. Biol. Ecol.* 400:278–287
- Roberts, D. A., Birchenough, S.R., Lewis, C., Sanders, M.B. et al., 2013. Ocean acidification increases the toxicity of contaminated sediments. *Glob. Change Bio.* 19:340-351
- Schwarz, A. M., Bjork, M., Buluda, T., Mtolera, H., et al., 2000. Photosynthetic utilisation of carbon and light by two tropical seagrass species as measured in situ. *Mar. Biol.* 137: 755-761
- Stitt, M., Krapp, A., 1999. The interaction between elevated carbon dioxide and nitrogen nutrition; the physiological and molecular background. *Plant cell Environ.* 22: 583-621
- Thom, R. M., 2001. Carbon Sinks in Nearshore Marine Vegetated Ecosystems. Pacific Northwest Laboratory, Marine Science Laboratory

Takahashi, M., Noonan, S.H.C., Fabricus, K.E., Collier, C.J., 2016. The effects of long term in situ CO<sub>2</sub> enrichment on tropical seagrass communities at volcanic vents. *ICES J. of Mar. Sc.* 73: 876-886.

Touchette, B.W., Burkholder, J.M., 2000. Overview of the physiological ecology of carbon metabolism in seagrasses. *J. Exp. Mar. Bio. Ecol.* 250:169-205

Uku, J., Beer, S., Bjork, M., 2005. Buffer sensitivity of photosynthetic carbon utilisation in eight tropical seagrasses. *Mar. Biol.* 147: 1085-1090

Vizzini, S, Di Leonardo, R., Costa, V., Tramati, C.D. et al., 2013. Trace element bias in the use of CO<sub>2</sub> seeps as analogues for low pH environments: Implications for contamination level in acidified oceans. *Estu. Coas. and Shelf Sc.* 134:19-30.

Zimmerman, R.C., Kohrs, D.G., Steller, D.L., Alberte, R.S., 1997. Impacts of CO<sub>2</sub> enrichment of productivity and light requirements of eelgrass. *Plant Physiol.* 115:599-607

Zeng, X., Chen, X., 2015. The positive relation between ocean acidification and pollution. *Mar. Poll. Bull.* 91:14-21.

