



ORIGINAL ARTICLE

A report on a survey among Portuguese Association of Interventional Cardiology associates regarding ionizing radiation protection practices in national interventional cath-labs



Hugo Costa^{a,*}, Hugo Vinhas^a, Rita Calé^b, Ernesto Pereira^b, Joana Santos^c, Graciano Paulo^c, Elisabete Jorge^d, Bruno Brochado^e, Bruno Melica^f, Rui Baptista Gonçalves^g, Eduardo Infante de Oliveira^{h,i,j}, on behalf of APIC^k

^a Cardiology Department, Centro Hospitalar Universitário do Algarve, Portugal

^b Cardiology Department, Hospital Garcia de Orta, Portugal

^c Escola Superior de Tecnologia da Saúde de Coimbra, Portugal

^d Centro Hospitalar Universitário de Coimbra, Portugal

^e Centro Hospitalar Universitário do Porto, Portugal

^f Centro Hospitalar de Vila Nova de Gaia, Portugal

^g Department of Public Health, University of Liverpool, United Kingdom

^h Cardiology Department, Centro Hospitalar de Lisboa Ocidental, Portugal

ⁱ Cardiology Department, Lisbon Lusiadas Hospital, Portugal

^j Institute of Physiology, Faculty of Medicine, University of Lisbon, Portugal

^k Associação Portuguesa de Cardiologia de Intervenção (APIC), Portugal

Received 30 April 2023; accepted 26 July 2023

Available online 10 November 2023

KEYWORDS

Ionizing radiation;
Dosimeter;
Intervention
cardiology;
Safety methods;
Protective equipment

Abstract

Introduction and objectives: Concerns surrounding the consequences of ionizing radiation (IR) have increased in interventional cardiology (IC). Despite this, the ever-growing complexity of diseases as well as procedures can lead to greater exposure to radiation. The aim of this survey, led by Portuguese Association of Interventional Cardiology (APIC), was to evaluate the level of awareness and current practices on IR protection among its members.

Methods: An online survey was emailed to all APIC members, between August and November 2021. The questionnaire consisted of 50 questions focusing on knowledge and measures of IR protection in the catheterization laboratory. Results were analyzed using descriptive statistics.

Results: From a response rate of 46.9%, the study obtained a total sample of 159 responses (156 selected for analysis). Most survey respondents (66.0%) were unaware of the radiation exposure

* Corresponding author.

E-mail address: hugoalexcosta@sapo.pt (H. Costa).

PALAVRAS-CHAVE

Radiação ionizante;
Dosímetro;
Cardiologia de
intervenção;
Técnicas de
minimização;
Equipamentos de
proteção;
Limites de dose

category, and only 60.4% reported systematically using a dosimeter. A large majority (90.4%) employed techniques to minimize exposure to radiation. All participants used personal protective equipment, despite eyewear protection only being used frequently by 49.2% of main operators. Ceiling suspended shields and table protectors were often used. Only two-thirds were familiar with the legally established limit on radiation doses for workers or the dose that should trigger patient follow-up. Most of the survey respondents had a non-certified training in IR procedures and only 32.0% had attended their yearly occupational health consultation.

Conclusions: Safety methods and protective equipment are largely adopted among interventional cardiologists, who have shown some IR awareness. Despite this, there is room for improvement, especially concerning the use of eyewear protection, monitoring, and certification.

© 2023 Sociedade Portuguesa de Cardiologia. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Resultados do inquérito realizado entre os associados da APIC sobre as práticas utilizadas na proteção da radiação ionizante nos laboratórios nacionais de cardiologia de intervenção

Resumo

Introdução e objetivos: A preocupação com as consequências da radiação ionizante tem vindo a crescer na cardiologia de intervenção. O objetivo da Associação Portuguesa de Cardiologia Interventiva (APIC) foi avaliar o nível de sensibilização e práticas atuais na proteção contra a radiação ionizante.

Métodos: Um inquérito voluntário foi enviado por e-mail aos membros da APIC, entre Agosto-Novembro de 2021, que consistiu em 50 questões relativas ao conhecimento e medidas de proteção contra a radiação. Posteriormente foi efetuada a respetiva análise descritiva.

Resultados: Obtiveram-se 159 respostas (46,9% do inquérito enviado) - 156 foram consideradas adequadas para análise. A maioria (66,0%) desconhecia a categoria de exposição à radiação correspondente e apenas 60,4% utilizaram o dosímetro de forma sistemática. 90,4% dos inquiridos utilizaram técnicas para minimizar a exposição à radiação. Praticamente todos os participantes utilizaram equipamento de proteção individual, mas a proteção ocular apenas ocorreu de forma frequente em 49,2% dos operadores principais. Escudos de teto e protetores de mesa foram os materiais disponíveis mais utilizados. Apenas dois terços conheciam a dose limite legalmente estabelecida para os trabalhadores ou a dose que deve desencadear um acompanhamento do paciente. A maioria dos inquiridos tinha uma formação não certificada em radiação ionizante e apenas 32,0% tiveram consulta anual de medicina do trabalho.

Conclusões: A consciencialização e as práticas adotadas para proteção contra radiação ionizante têm vindo a aumentar entre os profissionais de cardiologia de intervenção. Ainda assim, existe espaço para melhorar, nomeadamente no uso da proteção ocular, necessidade de monitorização e certificação.

© 2023 Sociedade Portuguesa de Cardiologia. Publicado por Elsevier España, S.L.U. Este é um artigo Open Access sob uma licença CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Interventional cardiology (IC) has witnessed impressive evolution over the last 20 years, both due to improvements in knowledge of coronary heart disease, but also linked to newer and more modern techniques and equipment. Exposure to ionizing radiation (IR), projected through fluoroscopy/cinefluorography is a consequence of the improvements in minimally invasive techniques in cardiac procedures.¹

Initially, these procedures had mainly diagnostic purposes, exposing patients to a minimum radiation time. The

goal over several years was to improve imaging quality, and thereby increase radiation intensity.² In the last 10 years, the concern regarding the consequences of IR has increased, as the incidence of adverse reactions and stochastic effects rose, leading to the publication of international guidelines on radiological protection during medical procedures.³⁻⁶

Modern equipment enables better image quality with less exposure to IR, by maximizing equipment technology features. Notwithstanding this, alongside technological innovation, procedures became more complex and mostly therapeutic; for example, coronary chronic total occlusions (CTO) with consequent greater exposure to IR. Thus, the

focus on the complexity of the disease and intervention may overshadow the required measures that must be instituted in order to reduce exposure.^{2,7}

The concern over patient exposure to IR is notorious among interventional cardiologists. Radiation doses in clinical practice have become a target of great interest. Unlike patients, the catheterization laboratory (Cath-Lab) team is constantly exposed to radiation, but surprisingly only a few Cath-Labs report their IR data.^{7–10}

Interventional cardiologists should work towards implementing novel methods for IR protection, ensuring controlled exposure in order to reduce the deleterious effects in the short- and the long term, for both patients and professionals.^{11,12}

This survey aimed to evaluate the awareness of IC health professionals regarding IR, its deleterious effects on patients and Cath-Lab team, as well as to understand the current practices concerning ionizing radiation protection.

Objectives

The aim of this survey, developed by the Portuguese Association of Interventional Cardiology (APIC), was to evaluate the level of awareness and current practices regarding IR protection.

Methods

Survey

A voluntary and anonymous questionnaire (Supplement S1) was emailed to all registered APIC members between August and November 2021. The questionnaire consisted of 50 questions, including demographics as well as four main topics: knowledge surrounding IR; techniques and methods used to minimize exposure to IR; equipment used to protect against IR; awareness/prevention on the effects of IR. Inclusion criteria were being an active member of APIC, and currently employed in an IC role. For the purpose of this study, the IC team was composed of the following members: main operator – interventional cardiologist who performs the procedures as the first operator; assistant operator – interventional cardiologist/interventional cardiology fellow/cardiologist fellow or cardiorespiratory technician/nurse who is an intervention assistant on the intervention table; room assistant – cardiorespiratory or radiology technician/nurse whose function is to assist the team that is on the intervention table (for example as a circulating nurse or scrub assistant).

Data analysis

Data were cleaned and a descriptive analysis was subsequently performed. Categorical variables are presented as frequencies and percentages, and continuous variables as means and standard deviations, or medians and interquartile ranges for variables with skewed distribution or a significant Shapiro–Wilk test. Statistical analysis was performed with SPSS Inc., Chicago, IL, USA, version 21.0.




Figure 1 Type of individual monitoring used by main operators (%) (53 responses).

Results

The questionnaire was emailed to a total of 329 APIC members (186 were effective associates – medical specialists, 85 aggregate associates – residents, and 58 affiliates – non-physicians). All active members have a registered email in APIC's mailing list. A total of 159 completed surveys were returned (46.9%). Two questionnaires were excluded as they did not consent to publishing their responses, whereas another was excluded as they did not currently work in a Cath-Lab. The final sample was thus composed of 156 participants. At the time, 85 IC subspecialists were registered in the Association of Medical Doctors of Portugal (Ordem dos Médicos – <https://ordemdosmedicos.pt/estatisticas-nacionais/#1505070622822-9e31b1b1-6d1a>).

Demographics

Most of the survey respondents were male (81 of 156, 51.9%), aged between 30 and 49 years (62.2%). Physicians were the most represented class, accounting for 67 of 156 (42.9%), from which 61 (39.1%) were main operators (only six doctors were considered assistant operators). Radiation exposure reflected by a higher time experience in Cath-Lab for over 20 years was present in 49 (31.4%) participants. Most responses were from professionals based in the Greater Lisbon area (63 of 156, 40.4%), although all regions were represented (Table 1).

Knowledge-based questions

These questions aimed to assess the level of knowledge regarding IR, as well as understanding the different forms of monitoring used by the sample (Supplement S1, Table 1).

Most survey respondents were unaware of what type of occupational radiation exposure category they belonged to (103 of 156, 66.0%), with older participants mostly in category A (27 of 96, 28.7%). The vast majority of participants were monitored dosimetrically (150 of 156, 96.2%), despite only 60.4% (93 of 154) participants reporting a systematic use of the dosimeter. Full body monitoring was the most used approach by all participants (131 of 143, 91.6%). Almost a third of main operators also used a ring for radiation monitoring (15 of 53, 28.3%) (Figure 1). Dosimeter reading was

Table 1 Demographic characteristics, occupation, years of radiation exposure and distribution by region of the survey respondents.

	Age		Total (n=156)
	<40 years (n=60, 38.5%)	≥40 years (n=96, 61.5%)	
<i>Gender</i>			
Male, n (%)	29 (48.3)	52 (54.2)	81 (51.9)
Female, n (%)	31 (51.7)	44 (45.8)	75 (48.1)
<i>Occupation</i>			
Doctor, n (%)	27 (45.0)	40 (41.7)	67 (42.9)
Nurse, n (%)	19 (31.7)	17 (17.7)	36 (23.1)
Cardiorespiratory technician, n (%)	5 (8.3)	11 (11.5)	16 (10.3)
Radiographer, n (%)	9 (15.0)	28 (29.2)	37 (23.7)
<i>Operator role</i>			
Main operator, n (%)	21 (35.0)	40 (41.7)	61 (39.1)
Assistant operator, n (%)	22 (33.6)	17 (17.7)	39 (25.0)
Room assistant, n (%)	17 (28.3)	39 (40.6)	56 (35.9)
<i>Professional experience in Cath-Lab</i>			
<5 years, n (%)	26 (43.3)	5 (5.2)	31 (19.9)
5–10 years, n (%)	23 (38.3)	9 (9.4)	32 (20.5)
10–20 years, n (%)	11 (18.3)	33 (34.4)	44 (28.2)
>20 years, n (%)	0 (0.0)	49 (51.0)	49 (31.4)
<i>Workplace location</i>			
Northern region, n (%)	9 (15.0)	20 (20.8)	29 (18.6)
Central region, n (%)	16 (26.7)	22 (22.9)	38 (24.4)
Greater Lisbon region, n (%)	21 (35.0)	42 (43.8)	63 (40.4)
Southern region, n (%)	11 (18.3)	8 (8.3)	19 (12.2)
Madeira and Azores, n (%)	3 (5.0)	4 (4.2)	7 (4.5)

performed monthly in 100 (64.9%) survey respondents. Exposure in more than one entity (54 of 156, 34.6%), as well as communication between them (13 of 82, 30.5%), was not a usual practice.


Most participants had access to dose recordings (117 of 155, 75.5%) and had not been previously called due to exceeding occupational dose limits (131 of 155, 84.5%). The majority knew the legally defined dose limits (100 of 155, 64.5%), as well as the need for mandatory notification in case of pregnancy (123 of 156, 78.8%), but were not aware of the trigger levels that determine the need for exposed patient follow-up (102 of 156, 65.4%).

Having received training in radiation protection was reported by 104 of 156 (66.7%) survey respondents, but this was mainly informal training. Academic (formal) training was more commonly reported by younger participants (14 of 33, 42.4%) (Figure 2).

Safety


Questions focusing on safety methods aimed to assess the use of techniques in clinical practice that minimize IR exposure (Supplement S1, Table 2).

Most participants used techniques to minimize exposure to IR (141 of 156, 90.4%). Optimization of the interventional table and detector position were reported to be always used by 36 of 58 (62.1%) main operators. Fluoroscopy optimiza-


**Figure 2** Type of training in radiation protection (%) (103 responses).

tion and collimation were also frequently adopted by main operators (always or often in 68.4% and 65.5%, respectively). Distance to the radiation source was less frequently adopted by main operators (Figure 3) (details in Supplement S1, Table 2.1). IC operators mostly used low-dose protocols (total sample: 120 of 156, 76.9%; main operators: 48 of 61, 78.7%), favored a pulse rate of 7.5 pulse/second (p/s) during fluoroscopy (total sample: 71 of 128, 55.5%; main operators: 31 of 56, 55.4%) and a frame rate of 15 frame/second (f/s) in cinefluorography (total sample: 51 of 130, 39.2%; main operators: 28 of 56, 50.0%).


Optimization table / Detector position



Optimization Fluoroscopy



Optimization Magnitude/ Collimation



Optimization radiation source/ Operator distance




Figure 3 Techniques used by main operators to minimize exposure to ionizing radiation and frequency (%) (58 responses).

Almost a quarter reported having finished a procedure by reaching a radiation dose greater than tolerated levels (total sample: 42 of 156, 26.9%; main operators: 17 of 61, 27.9%), mainly when performing a CTO.

When not involved in direct patient contact, 96.5% (110 of 114) of survey respondents would follow specific room positioning within the Cath-Lab, with most favoring the bottom of the patient's table (81 of 112, 72.3%).

Personal protective equipment

One section of the questionnaire also aimed to ascertain the access to and availability of radiological protection and its use within the Cath-Lab (Supplement S1, Table 3).

All participants reported using personal radiation protection equipment, mainly thyroid protectors and aprons (96.2%). Head and eye protections (32 of 123, 26.0%) and arms/hands and legs protections (5 of 88, 5.7%) were used less often (Figure 4). Despite largely available, eye protection with glasses/visors was seldom used (glasses: always and often in 50 of 127, 39.4%; visors: always and often in

6 of 91, 6.6%; glasses and visors: always and often in 56 of 156, 35.9%), and this was also reported when evaluating the main operators only (glasses: always and often in 26 of 58, 44.8%; visors: always and often in 4 of 45, 8.8%; glasses and visors: always and often in 30 of 61, 49.2%) (Supplement S1, Table 3.1). Protective gloves were frequently not used by survey respondents (only 3 of 44 main operators used, 6.8%). Beanies, gloves, sleeves, and boots were the radiation protection equipment least available by the Cath-Labs (Supplement S1, Figure 1).

Most Cath-Labs provided additional radiation protections (Figure 5) including suspended ceiling shield, table with pelvic protection and portable barrier, frequently used (Supplement S1, Figure 2).

Awareness and prevention of the consequences of radiation

The survey also assessed the awareness of participants regarding IR exposure, as well as determining the regularity of occupational health reviews (Supplement S1, Table 4).




Figure 4 Types of radiological protection equipment used and frequency (%).




Figure 5 Types of accessories provided by the Cath-Labs to complement radiation protection (%) (150 responses).

Most survey respondents had attended at least one occupational health consultation (104 of 156, 66.7%), but only 39 of 122 (32.0%) reported a yearly attendance. Lens evaluation during occupational health review was reported by 28 of 134 (20.6%) of participants, and only one participant had been referred for an ophthalmology review. General investigations such as thyroid function tests, chest radiograph and electrocardiogram were usually included in the occupational health review (Table 2).

Table 2 Exams requested in the occupational medicine consultation (%) (119 responses).

	Total n=156
<i>General analysis, n (%)</i>	
119 responses	117 (98.3)
<i>Thyroid function, n (%)</i>	
119 responses	53 (44.5)
<i>Chest X-ray, n (%)</i>	
119 responses	46 (38.7)
<i>Electrocardiogram, n (%)</i>	
119 responses	64 (53.8)
<i>Ophthalmology, n (%)</i>	
119 responses	1 (0.8)
<i>Thyroid ultrasound, n (%)</i>	
119 responses	5 (4.2)
<i>Other, n (%)</i>	
119 responses	4 (3.4)

Other: bone densitometry, respiratory function tests.




Figure 6 Quantification of concern about the radiation the patient (a) and the operator (b) are exposed to, compared according to age (1-low to 5-high) (146 responses).

The majority of survey respondents had concerns regarding the exposure to radiation by operators (150 of 156, 96.2%) and patients (141 of 156, 90.4%) during procedures. Overall, the concern was high (Figure 6).

Discussion

The findings of this survey highlight several features related to current knowledge, management and prevention strategies associated with IR across Cath-Labs in Portugal.

Knowledge

For monitoring and surveillance, there are two categories of exposed workers:

- category A: workers who are liable to receive an effective dose of 20 mSv per year averaged over five consecutive years (100 mSv in 5 years) and of 50 mSv in any single year; an equivalent dose to the lens of the eye of 20 mSv per year averaged over five consecutive years (100 mSv in 5 years) and of 50 mSv in any single year; an equivalent dose to the extremities (hands and feet) or to the skin of 500 mSv in a year, and
- category B: workers who are not classified as A.^{3,13,14}

Knowledge of these categories is crucial for the understanding of the suitability (and safety) for any given radiation dose exposure.

In the present survey, most of the workers did not know to which category they belonged (66.0%). This is also important in defining the periodicity of dosimeter readings (monthly in category A and quarterly in category B).¹³ A total of 64.9% of the survey respondents reported monthly readings, which should be equivalent to the true percentage of workers in category A.

The use of dosimeters should be systematic, which is not the case for about 40.0% of the survey respondents. Only two-thirds of participants knew the legally established limits for workers (50 mSv in any single year), as well as the exposure that should trigger patient follow-up due to possible tissue reactions (cumulative air kerma ($K_{a,r}$) of 5000 mGy or air kerma-area product (P_{KA}) of 500 Gy cm²).¹⁵ Any pregnant or breastfeeding professional must report to the occupational health team, and may not perform functions involving a significant risk (should not exceed 1 mSv during pregnancy).^{4,13,16} The majority of participants appear to

be aware of these regulations. Certified IR training is recommended to all team members, but this appears to be lacking in the majority of survey respondents.^{1,13}

Safety

Methods and techniques to minimize radiation exposure were often used (90.4%), albeit “as low as reasonably achievable” (ALARA). The use of collimation and the reduced distance between the patient and the image detector (raised table, low detector) was widely used by the survey respondents (main operators). Collimation reduces the irradiated volume, the overall patient dose and scatter radiation, and this parameter directly affects P_{KA} .^{1,17} In addition, guidelines also advise the use of “store-fluoro” option and to avoid angled projections, techniques to minimize IR that can potentially be used more often in Portuguese Cath-Labs.^{1,17}

The use of low-dose protocols is important to minimize exposure, as was the case for 76.9% of survey respondents. Low-rate fluoroscopy at 7.5 p/s is a very effective measure to reduce both operator and patient radiation exposure, without significant effect on procedure duration, fluoroscopy time or contrast use. A reduction from 15 p/s to 7.5 p/s with a low-dose fluoroscopy mode minimizes radiation exposure by 67%.^{1,8,9,18–20} It is important to highlight that most main operators used fluoroscopy pulse rate at 7.5 p/s (55.4%), but this may not be standard practice in all Cath-Labs.

Performing a CTO was the main intervention leading to suspension of a procedure due to reaching the permitted radiation dose limit. Thereby, it is recommended that operators with experience in highly complex procedures must be aware of the importance of intensifying efforts to reduce radiation exposure for both the patient and the operator.⁸

Personal protective equipment

Knowledge of the distribution of scattered radiation levels around a patient and the effective use of protective devices are crucial for team protection.^{12,21} Personal protective devices comprise aprons (or vest/skirt), thyroid protectors, eyewear and radiation protection gloves.²² All survey respondents used aprons and thyroid protectors. Aprons should be individually tailored, because if oversized, they normally provide insufficient protection to the exposed areas. The fitting of the protective apron is often more important to reduce effective dose than the lead equivalent. Aprons do not protect the neck, and if no thyroid protection is worn, the dose at which the unprotected thyroid is irradiated may be twofold that of the effective dose.¹²

The most important factor in protecting the head and the eyes is the proper use of ceiling shields. Lead glasses/visors are an important component of the eyewear protection, proved to reduce the lens exposed dose substantially.¹² In one study, 45.0% of IC professionals had posterior subcapsular lens changes attributed to IR exposure with infrequently use of lead glasses (only 57% of IC).²³ Considering the low rates of glasses/visors used in this study, across all professionals, alongside the scarcity of lens evaluation at

occupational health, eye health may be neglected in the sample.

Protective gloves block 30.0–40.0% of the scattered radiation with some limitation of finger movement. Its use can be justified when the IC have to operate close to the irradiated area.²² In the current study, only 6.8% of main operators of the survey respondents used this type of protection.

Curtains attached to the patient table and properly placed ceiling suspended shields have been shown to dramatically reduce operator dose.²² This type of protection is common in Cath-Labs and was used by most survey respondents.

Awareness and prevention of the consequences of radiation

There are two main biological effects of radiation: deterministic or tissue effects, which cause immediate and predictable tissue lesion, like cataracts (in professionals) and skin burns (in patients); and stochastic effects, with long term carcinogenic and genetic alterations.³ Most operators seem to have an insight into the deleterious effects of IR, with high rates of concern about their own and their patients' IR exposure.

Workers in category A should have at least one yearly occupational health appointment (including ophthalmologic/lens evaluation).¹³ Only 32.0% of the participants in the survey have been checked with this periodicity, despite the estimated real percentage of operators in category A being 64.9%.

Limitations

This survey has some methodological limitations worth highlighting. First, the survey was only sent to registered APIC members. Those professionals who were not registered, and would otherwise be eligible to participate, were not considered. This leads to the second limitation, which is the relatively small participant sample. However, considering the 85 IC subspecialists in Portugal in 2021, the main operators in the survey correspond to 71.8% of the population. Third, as certain members in the intervention team can have multiple roles, this may affect how they have responded to the survey. Finally, this survey was not applied to other areas of cardiology than IC and who also work with radiation, such as electrophysiology/pacing and pediatric cardiology. As such, the survey findings may not be generalizable to all national Cath-Labs or non-APIC members and would benefit from being disseminated to a wider sample of non-APIC members.

Conclusions

Interventional cardiology professionals have reported awareness of the use and implications of IR, as demonstrated by following relevant protocols and measures to minimize radiation, as well as the frequent use of personal protective equipment. Despite this, there is room for improvement, in particular when it comes to eyewear protection. In addition, considering the low rates of dosimeter use, as well as yearly

occupational health follow-ups and the lack of knowledge of the legal limits of radiation dose, there is an urgent need to improve monitoring and certification. Similarly, the documentation and reporting of radiation protection practices should be performed in Cath-Labs to arrive at a national standardization of procedures.

Conflicts of interest

The authors have no conflicts of interest to declare.

Acknowledgments

The authors would like to thanks to the physicians at Garcia de Orta Hospital, Vanessa de Sousa e Gabriela Cardoso, who collaborated very actively in the elaboration of the applied questionnaire. Also, the authors would like to express their gratitude to all APIC researchers and colleagues who cooperate in providing data for the current survey, and their commitment in attenuate the effects of IR in catheterization laboratories.

Appendix A. Supplementary data

Supplementary material associated with this article can be found in the online version at [doi:10.1016/j.repc.2023.07.008](https://doi.org/10.1016/j.repc.2023.07.008).

References

1. Hirshfeld JW, Balter S, Brinker JA, et al. ACCF/AHA/HRS/SCAI clinical competence statement on physician knowledge to optimize patient safety and image quality in fluoroscopically guided invasive cardiovascular procedures: a report of the American College of Cardiology Foundation/American Heart Association. *J Am Coll Cardiol*. 2004;44:2259–82.
2. Siiskonen T, Ciraj-Bjelac O, Dabin J, et al. Establishing the European diagnostic reference levels for interventional cardiology. *Phys Med*. 2018;54:42–8.
3. Picano E, Vano E. The radiation issue in cardiology: the time for action is now. *Cardiovasc Ultrasound*. 2011;9:1–13.
4. Williams MC, Stewart C, Weir NW, et al. Using radiation safely in cardiology: what imagers need to know. *Heart*. 2019;105:798–806.
5. Damilakis J, Paulo G, Frija G, et al. Radiation Protection series publication 195: European study on clinical diagnostic reference levels for X-ray medical imaging (EUCLID); 2021.
6. Abbott JD. Controlling radiation exposure in interventional cardiology. *Circ Cardiovasc Interv*. 2014;7:425–8.
7. Stocker TJ, Abdel-Wahab M, Möllmann H, et al. Trends and predictors of radiation exposure in percutaneous coronary intervention: the PROTECTION VIII study. *EuroIntervention*. 2022;18:e324–32.
8. Christakopoulos G, Karpaliotis D, Alaswad K, et al. Predictors of excess patient radiation exposure during chronic total occlusion coronary intervention: insights from a contemporary multicenter registry. *Can J Cardiol*. 2017;33:478–84.
9. Wassef AWA, Hiebert B, Ravandi A, et al. Radiation dose reduction in the cardiac catheterization laboratory utilizing a novel protocol. *JACC Cardiovasc Interv*. 2014;7:550–7.
10. Uthirapathy I, Dorairaj P, Ravi S, et al. Knowledge and practice of radiation safety in the catheterization laboratory among

- interventional cardiologists – an online survey. *Indian Heart J.* 2022;74:420–3.
11. Bar O, Maccia C, Pagès P, et al. A multicentre survey of patient exposure to ionising radiation during interventional cardiology procedures in France. *EuroIntervention.* 2008;3:593–9.
12. López PO, Dauer LT, Loose R, et al. ICRP publication 139: occupational radiological protection in interventional procedures. *Ann ICRP.* 2018;47:1–118.
13. European Parliament. Council Directive 2013/59/Euratom of 5 December 2013 laying down basic safety standards for protection against the dangers arising from exposure to ionising radiation, and repealing Directives 89/618/Euratom, 90/641/Euratom, 96/29/Euratom, 97/43/Euratom and 2003/122/Euratom. *Off J Eur Commun.* 2014;L13:1–73.
14. Yukiya A. Radiation protection and safety of radiation sources: international basic safety standards. IAEA Saf Stand Relat Publ; 2014 [Part 3].
15. Stecker MS, Balter S, Towbin RB, et al. Guidelines for patient radiation dose management. *J Vasc Interv Radiol.* 2009;20 Suppl.:S263–73.
16. Dauer LT, Miller DL, Schueler B, et al. Occupational radiation protection of pregnant or potentially pregnant workers in IR: a joint guideline of the society of interventional radiology and the cardiovascular and interventional radiological society of europe. *J Vasc Interv Radiol.* 2015;26:171–81.
17. Bhatt DL. Cardiovascular intervention: a companion to Braunwald's heart disease. 1st edition United Kingdom: Elsevier Health Sciences; 2015.
18. Abdelaal E, Plourde G, MacHaalany J, et al. Effectiveness of low rate fluoroscopy at reducing operator and patient radiation dose during transradial coronary angiography and interventions. *JACC Cardiovasc Interv.* 2014;7:567–74.
19. Steffenino G, Rossetti V, Ribichini F, et al. Short communication: staff dose reduction during coronary angiography using low framing speed. *Br J Radiol.* 1996;69:860–4.
20. Chambers CE, Fetterly KA, Holzer R, et al. Radiation safety program for the cardiac catheterization laboratory. *Catheter Cardiovasc Interv.* 2011;77:546–56.
21. Valentin J. Avoidance of radiation injuries from medical interventional procedures. *Ann ICRP.* 2000;30:1–67.
22. Bartal G, Vano E, Paulo G. Get protected! Recommendations for staff in IR. *Cardiovasc Intervent Radiol.* 2021;44:871–6.
23. Papp C, Romano-Miller M, Descalzo A, et al. Results of RELID study 2014 – Buenos Aires – Argentina. *Radiat Prot Dosim.* 2017;173:212–7.